

**Table 1.** Potential causes of frailty in patients with CKD reported in the literature

Category	Type		Risk Difference (95% CI)	Patient CKD severity	Frailty Assessment method	Sample Size	Ref
Demographic profile	Advanced age	Age > 60 years	OR 4.0 (1.0-16.2)	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
		per year	OR 1.02 (1.01-1.03)	stages 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
			OR 1.03 (1.01-1.04)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
	Female gender		OR 11.3 (2.3-55.6)	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
			OR 1.55 (1.27-1.88)	stage 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
			OR 11.6 (1.7-79.1)	Elderly with stage 5D (HD)	Multidimensional frailty score	46	<sup>62</sup> JKMS
	Male gender		OR 0.49 (0.39-0.62)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM
	Non-white race		OR 1.9 (1.1-1.3)	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
	Unemployed status		OR 1.89 (1.36-2.62)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
	Higher education level		OR 0.67 (0.49-0.91) for 7 <sup>th</sup> -12 <sup>th</sup> grade, 0.53 (0.35-0.82) for >12 <sup>th</sup> grade				
Lifestyle	Smoking		RR 1.18 (1.04-1.34)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017 SJKDT

Anthropometric parameters		BMI	OR 1.2 (1.0-1.4) per 5 kg/m <sup>2</sup>	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
			OR 1.06 (1.02-1.1) per kg/m <sup>2</sup>	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
			OR 0.58 (0.38-0.88) per kg/m <sup>2</sup>	Elderly with stage 5D	Multidimensional frailty score	46	<sup>62</sup> JKMS
		Waist circumference (cm)	OR 3.84 (1.39-10.61; 3 <sup>rd</sup> tertile)	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
CKD severity		Mild	OR 2.21 (1.49-3.28)	stages 1/2	Modified Fried Phenotypes	10256	<sup>5</sup> 2009 AJM
			OR 1.48 (1.00-2.19)	Cre > 1.3 mg/dL	CHS scale	5888	<sup>6</sup> 2004 AJKD
		Moderate	OR 2.48 (1.57-3.93)	stages 3a	Modified Fried Phenotypes	10256	<sup>5</sup> 2009 AJM
		Severe	OR 5.88 (3.40-10.16)	stages 3b-5		Modified CHS scale	336
			OR 2.8 (1.3-6.3)	stage 3b			
			OR 2.1 (1.0-4.7)	stage 4			
Biological							
Cardiovascular	Hypertension	RR 1.6 (1.26-2.04)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017 SJKDT	
	Peripheral vascular disease	RR 1.58 (1.34-1.8)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017SJKDT	
		OR 1.67 (1.16-2.41)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM	
	Left ventricular dysfunction	RR 1.18 (1.03-1.36)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017SJKDT	

	Cardiac disorder (any)	OR 1.43 (1.01-1.98)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
	Endothelial dysfunction	OR 3.86 (1.00-14.88)	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
Central nervous system	Cerebrovascular Accident	RR 1.34 (1.19-1.5)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017SJKDT
		OR 1.55 (1.05-2.29)	stage 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
		OR 1.85 (1.04-3.28)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM
		OR 1.56 (1.04-2.35)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
Pulmonary	COPD	OR 2.20 (1.20-4.03)	CKD stages 1-5	Modified Fried Phenotypes	10256	<sup>5</sup> 2009 AJM
Endocrinologic/ Metabolic	Diabetes	OR 1.68 (1.16-2.45)	CKD stages 1-5	Fried Phenotypes	10256	<sup>5</sup> 2009 AJM
		OR 1.35 (1.10-1.65)	stage 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
		OR 1.52 (1.18-1.96)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM
		OR 1.44 (1.11-1.87)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
	Obesity	OR 6.63 (1.16-36.77)	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
Cancer	Cancer	OR 1.89 (1.19-2.99)	CKD stages 1-5	Modified Fried Phenotypes	10256	<sup>5</sup> 2009 AJM

	Musculoskeletal	Arthritis	OR 3.34 (2.08-5.38)	CKD stages 1-5	Modified Fried Phenotypes	10256	<sup>5</sup> 2009 AJM
	Body composition	Fat mass	OR 3.27 (1.17-9.09; 2 <sup>nd</sup> tertile) and 4.97 (1.7-14.55; 3 <sup>rd</sup> tertile)	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
		ECW to ICW ratio	OR 3.85 (1.18-10.50; 3 <sup>rd</sup> tertile)				
Psychological		Depression	OR 3.97 (2.28-6.91)	stage 5T	Fried Phenotypes	773	<sup>48</sup> Clin Transplant
Functional status		Disability	OR 5.6 (4.12-7.62)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
Vascular access		Permanent vascular access (fistula or graft)	OR 0.71 (0.51-0.98)	stage 5D (HD)	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
Laboratory Data		Creatinine < 4 mg/dL*	RR 1.46 (1.22-1.71)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017SJKDT
		eGFR (per 5 mL/min/1.73m <sup>2</sup> increase)	OR 1.44 (1.23-1.68)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM
		Albumin < 3.2 (g/dL)	OR 1.89 (1.43-2.49)	stage 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
		Lower free testosterone, (per 50% lower)	OR 1.30 (1.03-1.58)	Male stage 5D (HD)	Fried Phenotypes	440	<sup>8</sup> 2018 NDT



**Table 2.** Potential modifiers of frailty trajectories in patients with CKD reported in the literature

Category	Type	Risk Difference (95% CI)	Patient CKD severity	Frailty Assessment method	Sample Size	Ref
Ethnicity	Hispanic	Frailty scores increase 0.6 (0-1.1) per year	stage 5D (HD)	Fried Phenotypes	762	<sup>2</sup> 2017 CJASN
	Black	Frail to non-frail after transplantation (RRR 1.98 [1.07-3.67])	stage 5	Fried Phenotypes	569	<sup>53</sup> Transplantation
Biological						
Endocrinologic/ Metabolic	Diabetes	Remain frail after transplantation (RRR 2.56 [1.22-5.39])	stage 5	Fried Phenotypes	569	<sup>53</sup> Transplantation
		Frailty scores increase 0.7 (0.3-1.0) per year	stage 5D (HD)	Fried Phenotypes	762	<sup>2</sup> 2017 CJASN
Laboratory data	IL-6	Frailty scores increase 0.3 (0.1-0.4) per year				
	Serum Albumin Concentrations (g/dL)	Frailty scores decrease 1.1 (0.7-1.5) per g/dL				
	Low free testosterone (< 147 pmol/L)	Developing Frailty over 12 months (OR 1.56, 1.04-2.33)				
Dialysis course	Time of dialysis	Frail to non-frail after	stage 5	Fried Phenotypes	569	<sup>53</sup>

	(year)	transplantation (RRR 0.88 [0.78-1])				Transplantation
Healthcare utilization						
Hospitalization	Hospitalization during past year	Frailty scores increase 0.6 (0.3-0.8) per year	stage 5D (HD)	Fried Phenotypes	762	<sup>2</sup> 2017 CJASN

**Table 3.** Unadjusted associates of frailty in CKD patients

Category	Type	Prevalence (Frail vs. non-frail, %), values, or correlation	Patient CKD Severity	Frailty Assessment method	Sample Size	Ref
Demographic profile	Age (years)	$r = 0.24, p = 0.04$	stage 5D (HD)	Fried Phenotypes	74	<sup>9</sup> 2018 Clinics
		57.0 vs. 52.0	stage 5D (HD)	Fried Phenotypes	324	<sup>10</sup> 2015 CJASN
		82.5 vs. 65.4	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
		62.1 vs. 58.5	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		69.4 vs. 56.6 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
		69 vs. 59	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
		78.4 vs. 65.5	stage 5D (HD)	FRAIL scale	51	<sup>29</sup> Nephrology
		71.7 vs. 61.5	stage 5D (HD)	CHS scale	214	<sup>35</sup> JBMM
		58 vs. 53	stage 5D (HD)	Performance-based frailty	80	<sup>36</sup> J Ren Nutr
		75.3 vs. 65.1	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
		69.5 vs. 63.7	stage 2-4	Modified Fried Phenotypes	168	<sup>40</sup> HQoLO
		64.9 vs. 57.3	stages 3-5	Modified Fried	61	<sup>41</sup> HQoLO



				Phenotypes		
		62.9 vs. 55.1	stage 5D (HD)	Fried Phenotypes	146	<sup>50</sup> JAGS
		55.8 vs. 50.7	stage 5T	Fried Phenotypes	537	<sup>54</sup> Am J Transplant
	Gender (male)	56% vs. 21% (moderate/severe vs. NF/mild)	stage 5D (HD)	FRAIL scale	46	<sup>16</sup> 2017 BMCG
		55% vs. 72%	stage 5D (HD)	Performance-based frailty	80	<sup>36</sup> J Ren Nutr
		11% vs. 51%	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
		42.3% vs. 71.4%	stages 3-5	Modified Fried Phenotypes	61	<sup>41</sup> HQoLO
		51.2% vs. 68.2%	stage 5D (HD)	Self-reported frailty	1646	<sup>47</sup> AJN
		60.8% vs. 57.9%	stage 5T	Fried Phenotypes	383	<sup>51</sup> Am J Transplant
Anthropometric parameters	BMI (kg/m <sup>2</sup> )	31.5 vs. 27.6 (based on DW)	stage 5D (HD)	Fried Phenotypes	324	<sup>10</sup> 2015 CJASN
		22.53 vs. 26.16	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
		28.3 vs. 25.6	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
		28.8 vs. 24.9	stage 5D (HD)	Performance-	80	<sup>36</sup> J Ren Nutr

				based frailty		
		30.1 vs. 28.1	stage 5D (HD)	Self-reported frailty	1646	<sup>47</sup> AJN
	Waist circumference (cm)	101.0 vs. 97.7	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		103.0 vs. 93.6	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
Multimorbidity	Charlson comorbidity index	5.0 vs. 2.0	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
		5.8 vs. 5.1	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		6.6 vs. 4.3 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
	Number of comorbidities	6 vs. 5	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> Can J Diabet
Dialysis duration	Duration	70.5 vs. 162.1 (weeks)	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
		47.9 vs. 34.3 (months)	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		61.5 vs. 45.8 (months) (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
Physical examination	Diastolic blood pressure (mmHg)	75.6 vs. 80.2	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR

		72.7 vs. 82.5 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
<b>Biological</b>						
Cardiovascular	Heart Failure (%)	30% vs 12%	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
		44% vs. 11%	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
		36.4% vs. 25.3%	stage 5D (HD)	Self-reported frailty	1646	<sup>47</sup> AJN
	Peripheral vascular disease (%)	38.8% vs. 17.21%	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
		13.6% vs. 0% (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
		42.6% vs. 10.5%	stage 5D (HD)	Fried Phenotypes	146	<sup>50</sup> JAGS
	Angina (%)	34% vs. 22%	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
	Coronary heart disease (%)	27.3% vs. 5.1% (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
	Atherosclerosis (%)	40.1% vs. 30.6%	stage 5D (HD)	Self-reported frailty	1646	<sup>47</sup> AJN
Central nervous system	Cerebrovascular Disease (%)	26.4% vs. 12%	stage 5D (HD)	Fried Phenotypes	324	<sup>10</sup> 2015 CJASN

Endocrinologic/ Metabolic	Diabetes (%)		64% vs. 49%	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
			63.6% vs. 27.1% (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
			65% vs. 45%	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
			80% vs. 44%	stage 5D (HD)	FRAIL scale	51	<sup>29</sup> Nephrology
			60% vs. 36%	stage 5D (HD)	Performance-based frailty	80	<sup>36</sup> J Ren Nutr
			63% vs. 43.7%	stage 5D (HD)	Self-reported frailty	1646	<sup>47</sup> AJN
			75.4% vs. 44.7%	stage 5D (HD)	Fried Phenotypes	146	<sup>50</sup> JAGS
	Obesity (%)		64% vs. 50%	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
			51.8% vs. 23.9%	stage 5D (HD)	Fried Phenotypes	324	<sup>10</sup> 2015 CJASN
Musculoskeletal	Osteoporosis		Higher in frail patients (p = 0.01)	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
Immunological	Viral infection						
		HCV infection	21.5% vs. 2.6%	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017 SJKDT
Body Composition	Fat mass		r = 0.25, p = 0.04	stages 3-5	Modified Fried	61	<sup>3</sup> 2012

					Phenotypes		Mansur
			40.2% vs. 30.5% (severe F vs. NF) (high fat prevalence)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
			40.7% vs. 35.0%	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
			30.7 vs. 24.4 kg				
	Total mass						
		trunk mass (kg)	40.8 vs. 48.88	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> Can J Diabet
			29.4 vs. 33.5	stage 5D (HD)	FRAIL scale	44	<sup>32</sup> JPSM
		Cephalic mass (kg)	4.64 vs. 4.93				
	Lower lean body mass		57.1% vs. 14.7%	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> Can J Diabet
			45.49 vs. 53.62 kg				
			34.7 vs. 43.1 kg	stage 5D (HD)	FRAIL scale	44	<sup>32</sup> JPSM
		Trunk lean mass (kg)	17.4 vs. 22.1				
			23.05 vs. 27.98	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> Can J Diabet
		Cephalic lean mass (kg)	3.74 vs. 4.69				
			3.06 vs. 3.29	stage 5D (HD)	FRAIL scale	44	<sup>32</sup> JPSM
		Gynoid lean mass (kg)	6.64 vs. 7.91	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> Can J Diabet
	Lean tissue mass		37.2 vs. 41.4 kg	stage 5D (PD)	In-house frailty	178	<sup>15</sup> 2018 KBPR

				questionnaire		
			53.6% vs. 67.5% (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193 <sup>18</sup> 2016 KBPR
		Skeletal muscle index (kg/m <sup>2</sup> )	6.55 vs. 7.41	stage 5D (PD)	Clinical Frailty Scale	119 <sup>12</sup> 2018 PDI
		Phase angle (degree)	5.24 vs. 6.24	stage 5D (HD)	Performance-based frailty	80 <sup>36</sup> J Ren Nutr
		Over-hydration (L)	4.19 vs. 2.49 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193 <sup>18</sup> 2016 KBPR
		Bone mineral density (g/cm <sup>2</sup> )		stage 5D (HD)	FRAIL Scale	43 <sup>34</sup> JAGS
		L3	0.81 vs. 0.97			
		L4	0.73 vs. 0.92			
		Femoral neck	0.43 vs. 0.63			
		T-score				
		L3	-1.97 vs. -0.64			
		L4	-2.6 vs. -0.95			
		Femoral neck	-3.47 vs. -1.68			
	Laboratory Data	Prealbumin (mg/dL)	28.9 vs. 38.3	stage 5D (PD)	Clinical Frailty Scale	119 <sup>12</sup> 2018 PDI
		Serum albumin (g/dL)	3.6 vs. 3.9	stages 1-4	Modified CHS scale	336 <sup>7</sup> 2012 AJKD

		$r = -0.263, p = 0.025$	stage 5D (HD)	Fried Phenotypes	74	<sup>9</sup> 2018 Clinics
		3.8 vs. 4.1	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> Can J Diabet
		2.92 vs. 3.48	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
		3.61 vs. 3.85	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
		3.29 vs. 3.49 vs. 3.58 (F + D vs. F – D vs. NF )	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		3.17 vs. 3.62 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
		3.7 vs. 3.9	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
		3.5 vs. 3.9	stage 5D (HD)	FRAIL scale	51	<sup>29</sup> Nephrology
		3.7 vs. 3.9	stage 5D (HD)	CHS scale	214	<sup>35</sup> JBMM
		3.5 vs. 3.8	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
		3.2 vs. 3.4	Elderly with stage 5D	Multidimensional frailty score	46	<sup>62</sup> JKMS
	Creatinine	299 vs. 115 umol/L	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> Can J Diabet
		11.6 vs. 9.9 mg/dL (moderate/severe vs.	stage 5D (HD)	FRAIL scale	46	<sup>16</sup> 2017 BMCG

		NF/mild)				
		12.2 vs. 10.4 mg/dL (F/PF vs. NF)	stage 5D (HD)	FRAIL scale	44	<sup>32</sup> JPSM
		8.1 vs. 11.1 mg/dL	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
	eGFR (mL/min/1.73m <sup>2</sup> )	41.1 vs. 52.5 (cystatin C)	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
		18 vs. 50	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> Can J Diabet
	Albuminuria (mg/g Cre)	311.2 vs. 102	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
	Phosphate (mg/dL)	4.1 vs. 3.7	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
	Hemoglobin (g/dL)	10.35 vs. 10.97	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
		r = -0.336, p = 0.004	stage 5D (HD)	Fried Phenotypes	74	<sup>9</sup> 2018 Clinics
		12.2 vs. 13.2	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
		10.1 vs. 9.2 (moderate/severe vs. NF/mild)	stage 5D (HD)	FRAIL scale	46	<sup>16</sup> 2017 BMCG
	Total cholesterol	4.48 vs. 5.18 mmol/L	stage 5D (PD)	In-house frailty	193	<sup>18</sup> 2016 KBPR



		(severe F vs. NF)		questionnaire		
		134 vs. 148 mg/dL	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
	LDL cholesterol (mmol/L)	2.51 vs. 3.02 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
	HDL cholesterol	1.18 vs. 1.38 mmol/L (severe F vs. NF)				
		40 vs. 46 mg/dL	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
	iPTH (pg/mL)	248.8 vs. 127.9	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
	Ferritin (ng/mL)	1202 vs. 534	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
	Transferrin saturation (%)	30.1 vs. 37.1	stages 3-5	Modified Fried Phenotypes	61	<sup>3</sup> 2012 Mansur
	25-OH-D (ng/dL)	$r = -0.363, p = 0.002$	stage 5D (HD)	Fried Phenotypes	74	<sup>9</sup> 2018 Clinics
	CRP (mg/dL)	3.8 vs. 2.1	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
		1.12 vs. 0.28 (natural Log transformed)	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
	IL-6 (pg/mL)	2.45 vs. 1.58 (natural Log transformed)	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
	nPNA (g/kg/day)	1.10 vs. 1.19	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR

CKD-related complications	Counts of complications	$r = 0.666, p < 0.0001$	Elderly with unknown CKD stages	Edmonton Frail Scale	35	<sup>17</sup> 2016 Rev Rene
Residual renal function	Residual eGFR (ml/min/1.73m <sup>2</sup> )	1.54 vs. 2.46	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		0.9 vs. 2.2	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
Care modality	Assisted PD	38.6% vs. 0% (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
	Living with caregivers	45% vs. 72%	Elderly with stages 4/5 CKD	Groningen frailty indicator	65	<sup>19</sup> Ren Fail
	Renal conservative care	45% vs. 2%	Elderly with stages 4/5 CKD	Groningen frailty indicator	65	<sup>19</sup> Ren Fail
Dialysis related parameters	Kt/V	1.69 vs. 1.55 (moderate/severe vs. NF/mild)	stage 5D (HD)	FRAIL scale	46	<sup>16</sup> 2017 BMCG
		1.44 vs. 1.58	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
	Weekly total Kt/V	1.74 vs. 1.96 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
	URR	76.2% vs. 72.5%	stage 5D (HD)	FRAIL scale	46	<sup>16</sup> 2017

		(moderate/severe vs. NF/mild)				BMCG
	Daily exchange volume (L)	6.5 vs. 7.0 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
Vascular access	Catheter, AVF, and AVG use	Catheter: 61.6% vs. 17.8% AVF: 27.78% vs. 77.5% AVG: 11.11% vs. 5%	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
Microbiota	Bacterial Abundance of some genera (Mogibacteriaceae, Coriobacteriaceae, Eggerthella, Erwinia, Coprobacillus, Anaerotruncus, etc)	Higher in F group	stages 3b/4	Fried Phenotype	79	<sup>20</sup> NDT
Neurological	Quantitative EEG findings		stage 5D (HD)	FRAIL scale	46	<sup>16</sup> 2017 BMCG
	Delta wave (central, right/left TO, left frontal area)	Lower in F group				
	Delta to alpha ratio (global, central, left frontal, right/left TO)	Lower in F group				

		area)		stage 5T	Fried phenotypes	665	<sup>30</sup> JASN
		Delta/theta to alpha/beta ratio (global, central, left frontal, right/left TO area)	Lower in F group				
		Cognitive impairment					
		Prevalence	11% vs. 6.6%				
		Pre-transplant 3MS scores	93.0 vs. 96.0				
		3MS memory	20.0 vs. 21.0				
		3MS identification/associa tion	23.0 vs. 24.0				
Psychological							
Mood	Mood Change	Negative correlation	stage 5D (HD)	Edmonton Frail Scale	60	<sup>21</sup> Act Paul Enferm	
Anxiety	Hospital Anxiety and Depression Scale	Higher global, psychological and social components (women) Higher physical	stage 5D (online- HDF)	N/A	97	<sup>22</sup> NDT	

		component (men)				
Depression	Depression	38.8% vs. 12.58%	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
	Hospital Anxiety and Depression Scale	Higher global, psychological, physical component (men)	stage 5D (online-HDF)	N/A	97	<sup>22</sup> NDT
	Self-reported major depression	83% vs. 6%	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> Can J Diabet
Physical activity	Minnesota Leisure Time Activity (LTA)	95 vs. 735	stage 5D (HD)	Fried Phenotypes	68	<sup>25</sup> J Ren Nutr
	Low Physical Activity Questionnaire (LoPAQ)	280 vs. 798				
	Sitting (hours/day)	6.5 vs. 5				
	Grip strength (kg)	16.4 vs. 24.6	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
	Walk speed (m/s)	0.79 vs. 1.67				
Nutritional Status	SGA scores	5.25 vs. 5.75	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		4.2 vs. 5.3 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
	MIS scores	8.14 vs. 5.12	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR

Quality of Life			12.2 vs. 6.0 (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
			7.6 vs. 3.9	stages 3b/4	Fried Phenotype	79	<sup>20</sup> NDT
	MNA scores		18.0 vs. 22.0	Elderly with stage 5D	Multidimensional frailty score	46	<sup>62</sup> JKMS
	Kidney Disease Quality of Life (KDQoL) components						
		Mental health	43.6 vs. 48.9	stage 5D (HD)	Fried Phenotypes	151	<sup>26</sup> J Ren Nutr
		Kidney disease symptoms	67.8 vs. 79.1				
	SF-36						
		Physical functioning	46 vs. 84	stages 3-5	Modified Fried Phenotypes	61	<sup>41</sup> HQoLO
		Role physical	53.8 vs. 75				
		Bodily pain	58.4 vs. 76.5				
		General health	48.9 vs. 62				
		Vitality	58.8 vs. 77.4				
		Mental health	69.5 vs. 80.8				
Functional outcomes	Ability for basic ADL		33.33% vs. 76.4%	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
			55% vs. 91%	Elderly with stages	Groningen frailty	65	<sup>19</sup> Ren Fail

			4/5 CKD	indicator		
		Ability to transfer	38.8% vs. 84.7%	stage 5D (HD)	Fried Phenotypes	320 <sup>11</sup> 2017 J Aging Res
		Barthel Index	90 vs. 100	stage 5D (PD)	Clinical Frailty Scale	119 <sup>12</sup> 2018 PDI
Disability		≥1 disability in ADL	15% vs. 5%	CKD stages 1-4	Fried Phenotypes	336 <sup>7</sup> 2012 AJKD
		≥1 disability in IADL	60% vs. 28%			
		≥1 disability in mobility	40% vs. 18%			
	Functional status	Karnofsky scores	44.4 vs. 95.36	stage 5D (HD)	Fried Phenotypes	320 <sup>11</sup> 2017 J Aging Res
Health-care utilization		Hospitalization >= 1 time per year	90% vs. 53%	Elderly with stages 4/5 CKD	Groningen frailty indicator	65 <sup>19</sup> Ren Fail
		Hospitalization frequency (per year)	0.78 vs. 0.28 episodes	stage 5D (HD)	Fried Phenotypes	320 <sup>11</sup> 2017 J Aging Res
		Hospitalization episode count	3.31 vs. 2.12 vs. 0.9 (in 2 years) (F + D vs. F – D vs. NF )	stage 5D (PD)	In-house frailty questionnaire	178 <sup>15</sup> 2018 KBPR
			5.2 vs. 2.4 per year (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193 <sup>18</sup> 2016 KBPR
		Cardiovascular origin hospitalization count	1.4 vs. 0.5 per year (severe F vs. NF)			

	Hospital stay (days per year)	26.62 vs. 14.05 vs. 8.04 (2 years) (F + D vs. F – D vs. NF )	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		58.5 vs. 18.3 per year (severe F vs. NF)	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
Technique survival	Technique failure	42.5% vs. 35.8% vs. 13.7% (2 years) (F + D vs. F – D vs. NF )	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
Mortality	Overall mortality	20.45% vs. 12.36% (1 year)	stage 5D (HD)	Fried Phenotypes	320	<sup>11</sup> 2017 J Aging Res
		37.5% vs. 28.6% vs. 13.4% (2 years) (F + D vs. F – D vs. NF )	stage 5D (PD)	In-house frailty questionnaire	178	<sup>15</sup> 2018 KBPR
		30% vs. 10% (1 year)	Elderly with stages 4/5 CKD	Groningen frailty indicator	65	<sup>19</sup> Ren Fail



**Table 4.** Confounder-adjusted risk of complications resulting from frailty in CKD patients

Category	Type	Hazard/odds ratio, Risk Difference (95% CI), or values in F vs. NF groups	Patient CKD Severity	Frailty Assessment method	Sample Size	Ref
Physical examination	Blood pressure	Lower in Frail group (adjusted $p = 0.001$ )	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> 2019 Can J Diabet
Biological						
Cardiovascular	QRS duration	$\beta = -0.29$ , $t = -2.03$ , $p = 0.048$	stage 5D	Edmonton frailty scale	41	<sup>28</sup> PeerJ
		$\beta = -0.27$ , $t = -1.84$ , $p = 0.05$	(HD)	FRAIL scale		
Musculoskeletal	Vertebral compression fracture (any)	OR 1.8 per FRAIL score ( $p = 0.01$ )	stage 5D (HD)	FRAIL Scale	43	<sup>34</sup> JAGS
Cognitive function	3MS scores	At baseline	stage 5D	Fried Phenotypes	324	<sup>10</sup> 2015 CJASN
		1-year	(HD)			
		Pre-transplant	stage 5T	Fried phenotypes	665	<sup>30</sup> JASN
		1-4 years post-transplant				
	TMT-A	At baseline	stage 5D (HD)	Fried Phenotypes	324	<sup>10</sup> 2015 CJASN
	TMT-B	At baseline				

Body composition	Lean mass	Lower lean mass over cephalic, trunk, and 4 extremities than NF group	stage 5D (HD)	FRAIL scale	44	<sup>32</sup> JPSM
	BMD at 1 year follow up		stage 5D (HD)	FRAIL Scale	43	<sup>33</sup> PeerJ
	Total	$\beta = -0.53, t = -3.27, p < 0.01$				
	L1	$\beta = -0.4, t = -2.18, p = 0.04$				
	L4	$\beta = -0.39, t = -2.1, p = 0.046$				
	Femoral neck	$\beta = -0.5, t = -2.96, p < 0.01$				
	Average L-spine areas					
	1 year of follow up	$\beta = -0.48, t = -2.84, p < 0.01$				
	Interval changes	$\beta = -0.5, t = -3.02, p < 0.01$				
	Interval changes in L-spine Z-score percentages	$\beta = -0.45, t = -2.11, p = 0.049$				
	QUS parameters					
	SOS	1487.8 vs. 1537.8 (female) 1493.7 vs. 1542.2 (male)	stage 5D (HD)	CHS scale	214	<sup>35</sup> JBMM
	BUA	86.2 vs. 100.7 (female) 93.8 vs. 107.8 (male)				

	Stiffness index	54.0 vs. 77.7 (female) 60.9 vs. 83.6 (male)				
	Muscles					
	Quadriceps muscle area	$r = -30.28, p = 0.02$	stage 5D (HD)	Performance-based frailty	80	<sup>36</sup> J Ren Nutr
	Appendicular skeletal muscle mass index (ASMI)	Lower in Frail group (adjusted $p < 0.05$ )	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> 2019 Can J Diabet
	Appendicular fat percentage		stage 5D (HD)	FRAIL scale scores	44	<sup>32</sup> JPSM
	Left/Right lower extremity	$\beta = 0.34, t = 2.32; p = 0.03$ (left); $\beta = 0.3, t = 2.05; p = 0.048$ (right)				
	Left/Right upper extremity	$\beta = 0.37, t = 2.66; p = 0.01$ (left); $\beta = 0.43, t = 3.09; p < 0.01$ (right)				
	Sarcopenia	OR 12.2 (2.27-65.5)	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
Laboratory data	Serum albumin (g/dL)	Negative relationship ( $p = 0.01$ )	stage 5D (HD)	FRAIL scale	46	<sup>37</sup> Nephrology
Psychological	Hospital anxiety and depression scale	OR 1.21 (1.11-1.31)	stage 5D	Clinical Frailty Scale	251	<sup>42</sup> 2016 CJASN
Functional status	Physical functioning	Lower in Frail group (adjusted $p =$	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> 2019 Can

		0.004)		(EFS)		J Diabet
	Need assistance in ADL	OR 1.93 (1.01-3.68) for pre-frail OR 11.32 (5.49-23.32) for frail	stage 5D (HD)	Modified Fried Phenotypes	742	<sup>31</sup> HDI
	Barthel index scores	OR 0.89 (0.86-0.93)	stage 5D	Clinical Frailty Scale	251	<sup>42</sup> 2016 CJASN
<b>Psychological</b>						
	Delirium	Post-transplantation delirium	OR 2.05 (1.02-4.13)	stage 5T	Fried Phenotypes	893 <sup>23</sup> JASN
	Distress	Self-reported distress thermometer	$\beta = 0.35$ (0.12-0.58), $t = 3.0$ , $p = 0.003$	stage 5D (HD)	Canadian frailty score	382 <sup>38</sup> Nutr Clin Pract
<b>Fall</b>	Any fall	HR 2.1 (1.21-3.92)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017SJKDT
		OR 2.39 (1.22-4.71)	Stage 5D (HD)	Modified Fried Phenotype	762	<sup>44</sup> CKJ
	Increased numbers of falls	HR 3.09 (1.38-6.90)	stage 5D (HD)	Modified Fried Phenotype	95	<sup>45</sup> BMCN
	Time to first fall	HR 1.60 (1.16-2.20)	stage 5D (HD)	Self-reported frailty	1646	<sup>47</sup> AJN
<b>Quality of Life</b>						
	KDQoL					
	Physical health	33.7 vs. 40.7	stage 5D	Fried Phenotypes	151	<sup>26</sup> J Ren
	Effects of disease	51.6 vs. 66.8	(HD)			Nutr
	KDQoL short form					

	Physical component	Difference -6.31 (-8.16 to -4.46)	stage 5T	Fried Phenotypes	443	43 Transplanta tion
	Physical functioning	Difference -14.17 (-18.58 to -9.76)				
	Role limitations	Difference -15.37 (-22.96 to -7.78)				
	Bodily pain	Difference -9.45 (-14.33 to -4.57)				
	General health	Difference -11.76 (-15.94 to -7.59)				
	Emotional well-being	Difference -3.05 (-6.01 to -0.09)				
	Social functioning	Difference -6.19 (-10.98 to -1.41)				
	Energy	Difference -11.66 (-16.3 to -7.03)				
	Kidney disease-specific HRQoL	Difference -6.53 (-9.17 to -3.89)				
	Symptoms	Difference -5.5 (-8.2 to -2.79)				
	Effects	Difference -7.69 (-11.66 to -3.72)				
	Burden	Difference -10.19 (-15.94 to -4.44)				
	Cognitive function	Difference -5.51 (-9 to -2.02)				
	Social interaction	Difference -4.7 (-7.85 to -1.56)				
	Sleep	Difference -6.29 (-10.56 to -2.02)				
	Social support	Difference -5.69 (-9.92 to -1.47)				
	HRQoL					
	Fair/Poor HRQoL at follow-up	OR 2.79 (1.32-5.90)	stage 5D	Fried Phenotypes	233	39 J Frailty Aging
	Worse HRQOL after	RR 2.91 (1.08-7.80)				

	follow-up					
	SF-36					
	Physical components	Lower in Frail group (adjusted $p = 0.002$ )	stages 1-5	Edmonton Frail Scale	41	<sup>14</sup> 2019 Can J Diabet
		$\beta = -0.566$ , $t = -8.792$ , $p < 0.001$	stage 2-4	Modified Fried Phenotypes	168	<sup>40</sup> HQoLO
		Mean difference -1.12 (-1.47 to -0.76)	stages 3-5	Modified Fried Phenotypes	61	<sup>41</sup> HQoLO
	Mental components	Mean difference -0.75 (-1.4 to -0.16)				
		$\beta = -0.485$ , $t = -6.709$ , $p < 0.001$	stage 2-4	Modified Fried Phenotypes	168	<sup>40</sup> HQoLO
	SF-12					
	MCS	OR 0.94 (0.91-0.97)	stage 5D	Clinical Frailty Scale	251	<sup>42</sup> 2016 CJASN
	PCS	OR 0.88 (0.84-0.91)				
	Symptom scores	OR 1.23 (1.13-1.34)				
	KDQOL-SF scores 3 months after transplant		stage 5T	Fried Phenotypes	443	<sup>43</sup> Transplantation
	Physical HRQoL	0.34/month vs. 1.35/month				
	Kidney disease-specific HRQoL	2.41/month vs. 3.75 points/month				
	Effects	4.01/month vs. 7.1/month				

		Cognitive function	1.28/month vs. 2.88/month				
		Social interaction	-0.57/month vs. 1.18/month				
Graft Loss		Risk of graft loss in depressive patients	aHR 6.20 (1.67 to 22.95)	stage 5T	Fried Phenotypes	773	<sup>48</sup> Clin Transplant
Immunosuppressant use		MMF dose reduction	HR 1.29 (1.01-1.66)	stage 5T	Modified Fried Phenotypes	525	<sup>13</sup> 2015 Transplant
Dialysis access survival		Access failure	HR 2.63 (1.03-6.71)	stage 5D (HD)	FRAIL scale	51	<sup>29</sup> Nephrology
Health-care utilization		Hospitalization or mortality	HR 1.56 (1.36-1.79)	stage 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
		Hospitalization	HR 2.06 (1.18-3.58)	stage 5D (HD)	Fried Phenotypes	205	<sup>1</sup> 2017SJKDT
			aHR 1.83 (1.41-2.37)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
			HR 1.43 (1.00-2.03)	stage 5D (HD)	Fried Phenotypes	146	<sup>50</sup> JAGS
		Number of all-cause hospitalizations	beta = 0.29, p < 0.0001	stage 5D (PD)	In-house frailty questionnaire	193	<sup>18</sup> 2016 KBPR
		Number of cardiovascular hospitalizations	beta = 0.37, p < 0.0001				

	Time to first hospitalization	HR 1.26 (1.09-1.45)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM
	Early Hospital Readmission	RR 1.59 (1.17-2.17)	stage 5T	Fried Phenotypes	383	<sup>51</sup> Am J Transplant
	Longer Length of Stay					
	LOS (days)	RR 1.15 (1.03-1.29)	stage 5T	Fried Phenotypes	589	<sup>52</sup> Ann Surg
	> 2 weeks	OR 1.57 (1.06-2.33)				
		OR 2.02 (1.20-3.40) for increased frail category; OR 1.92 (1.13-3.25) for increased frail scores	stage 5	Fried Phenotypes	569	<sup>53</sup> Transplantation
	In depressive patients	aRR 1.88 (1.70-2.08)	stage 5T	Fried Phenotypes	773	<sup>48</sup> Clin Transplant
	Hospitalization frequency	Higher in Frail group (adjusted $p < 0.001$ )	stages 1-5	Edmonton Frail Scale (EFS)	41	<sup>14</sup> 2019 Can J Diabet
	Emergency department visit frequency	Higher in Frail group (adjusted $p = 0.002$ )				
	Total medical visit frequency	Higher in Frail group (adjusted $p = 0.001$ )				
<b>Mortality</b>	Overall mortality	HR 2.17 (1.01-4.65) after transplantation	stage 5T	Fried Phenotypes	537	<sup>54</sup> Am J Transplant



		HR 2.0 (1.5-2.7)	stages 1-5	Modified Fried Phenotypes	10256	<sup>5</sup> 2009 AJM
		HR 1.57 (1.25-1.97)	stage 5D (incident)	Modified Fried Phenotypes	1576	<sup>46</sup> JAMA-IM
		HR 2.24 (1.60-3.15)	stage 5D	Modified Fried Phenotypes	2275	<sup>4</sup> 2007 JASN
		HR 1.22 (1.04-1.43)	stage 5D	Clinical Frailty Scale	390	<sup>55</sup> 2015 CJASN
		HR 4.28 (1.22-14.98)	stages 4/5	PRISMA questionnaire & TUGT	104	<sup>56</sup> SJKDT
		HR 9.83 (1.80-53.7)	stage 5D (PD)	Clinical Frailty Scale	119	<sup>12</sup> 2018 PDI
		HR 2.60 (1.04-6.49)	stage 5D (HD)	Fried Phenotypes	146	<sup>50</sup> JAGS
		HR 2.08 (1.04-4.16)	stage 5D	Modified CHS scale	1658	<sup>49</sup> J Ren Nutr
		HR 1.78 (1.15-2.8) for performance-based frailty; HR 1.66 (1.06-2.6) for self-reported frailty; HR 1.95 (1.19-3.2) for both definition positivity	stage 5D (HD)	Modified Fried Phenotypes and self-reported frailty	771	<sup>57</sup> CJASN
		HR 1.66 (1.03-2.67) in general; HR 3.77 (1.10-12.92) in general obesity; HR 2.38 (1.17-4.82) in abdominal	stage 5D (HD)	Fried Phenotypes	370	<sup>59</sup> NDT

		obesity				
		HR 2.43 (1.48-3.99)	stage 5D and 5T from ANCA vasculitis	Inability to walk without help	425	<sup>60</sup> QJM
		HR 1.93 (1.58-2.36)	stage 5D and 5T from MM or amyloidosis	Inability to walk without help	1462	<sup>61</sup> CJASN
	In depressive patients	aHR 2.62 (1.03 to 6.70)	stage 5T	Fried Phenotypes	773	<sup>48</sup> Clin Transplant
	Modify the association between comorbidity and mortality	HR 0.75 (0.44-1.29) in F group vs. 1.66 (1.17-2.35) in NF group	stage 5	Fried Phenotypes	2086	<sup>58</sup> Am J Nephrol
		HR 1.93 (1.58-2.36)	stage 5D and 5T from MM or amyloidosis	Inability to walk without help	1462	<sup>61</sup> CJASN
	Post-transplant mortality	HR 2.27 (1.11-4.65) for increased frail category; OR 2.36 (1.12-4.99) for increased frail scores	stage 5	Fried Phenotypes	569	<sup>53</sup> Transplantation

Composite	Mortality or dialysis	HR 2.5 (1.4-4.4)	stages 1-4	Modified CHS scale	336	<sup>7</sup> 2012 AJKD
	Mortality or cardiovascular hospitalization	HR 23.58 (1.61-346.03)	Elderly with stage 5D (HD)	Multidimensional frailty score	46	<sup>62</sup> JKMS
	30-day post-transplant complications	$\beta$ =13.31 (5.72-20.89), p = 0.0007	stage 5T	Groningen Frailty Indicator	150	<sup>63</sup> Transplant Int

## Bibliography

1. YadlaM, JohnJ, MummadiM. A study of clinical assessment of frailty in patients on maintenance hemodialysis supported by cashless government scheme. Saudi J Kidney Dis Transplant. 2017. doi:10.4103/1319-2442.198102
2. JohansenKL, DalrympleLS, DelgadoC, et al. Factors Associated with Frailty and Its Trajectory among Patients on Hemodialysis. Clin J Am Soc Nephrol. 2017;12(7):1100-1108. doi:10.2215/CJN.12131116
3. MansurHN. Fragilidade na doença renal crônica: prevalência e fatores associados. 2012.
4. JohansenKL, ChertowGM, JinC, KutnerNG. Significance of frailty among dialysis patients. J Am Soc Nephrol. 2007;18(11):2960-2967.
5. Wilhelm-LeenER, HallYN, MKT, ChertowGM. Frailty and chronic kidney disease: the Third National Health and Nutrition Evaluation Survey. Am J Med. 2009;122(7):664-71 e2. doi:10.1016/j.amjmed.2009.01.026
6. ShlipakMG, Stehman-BreenC, FriedLF, et al. The Presence of Frailty in Elderly Persons with Chronic Renal Insufficiency. Am J Kidney Dis. 2004;43(5):861-867. doi:10.1053/j.ajkd.2003.12.049
7. RoshanravanB, KhatriM, Robinson-CohenC, et al. A prospective study of frailty in nephrology-referred patients with CKD. Am J Kidney Dis. 2012;60(6):912-921. doi:10.1053/j.ajkd.2012.05.017
8. ChiangJM, KaysenGA, SegalM, ChertowGM, DelgadoC, JohansenKL. Low testosterone is associated with frailty, muscle wasting and physical dysfunction among men receiving hemodialysis: a longitudinal analysis. Nephrol Dial Transplant. 2018. doi:10.1093/ndt/gfy252
9. DemirciogluD. The Association of Vitamin D Levels and the Frailty Phenotype Among Non-geriatric Dialysis Patients: A Cross-sectional Study. Clinics. 2018. doi:10.6061/clinics/2018/e116
10. McAdams-DemarcoMA, TanJ, SalterML, et al. Frailty and cognitive function in incident hemodialysis patients. Clin J Am Soc Nephrol. 2015;10(12):2181-2189. doi:10.2215/CJN.01960215
11. Bancul, GraterolF, BonalJ, et al. Frail Patient in Hemodialysis: A New Challenge in Nephrology-Incidence in Our Area, Barcelones Nord and Maresme. J Aging Res. 2017;2017:7624139. doi:10.1155/2017/7624139
12. KamijoY, KandaE, IshibashiY, YoshidaM. Sarcopenia and Frailty in PD: Impact on Mortality, Malnutrition, and Inflammation. Perit Dial Int. 2018;38(6):447-454. doi:10.3747/pdi.2017.00271

13. McAdams-DemarcoMA, LawA, TanJ, et al. Frailty, mycophenolate reduction, and graft loss in kidney transplant recipients. *Transplantation*. 2015;99(4):805-810. doi:10.1097/TP.0000000000000444
14. Adame PerezSI, SeniorPA, FieldCJ, JindalK, MagerDR. Frailty, Health-Related Quality of Life, Cognition, Depression, Vitamin D and Health-Care Utilization in an Ambulatory Adult Population with Type 1 or Type 2 Diabetes Mellitus and Chronic Kidney Disease: A Cross-Sectional Analysis. *Can J Diabetes*. June2018. doi:10.1016/j.jcjd.2018.06.001
15. SzetoC-C, ChanGC-K, NgJK-C, et al. Depression and Physical Frailty Have Additive Effect on the Nutritional Status and Clinical Outcome of Chinese Peritoneal Dialysis. *Kidney Blood Press Res*. 2018;43(3):914-923. doi:10.1159/000490470
16. ChaoC-T, LaiH-J, TsaiH-B, YangS-Y, HuangJ-W. Frail phenotype is associated with distinct quantitative electroencephalographic findings among end-stage renal disease patients: an observational study. *BMC Geriatr*. 2017;17(1):277. doi:10.1186/s12877-017-0673-3
17. deSousa MeiraA, Aparecido BatistaM, dePina PereiraRM, Partezani RodriguesRA, Silva FhonJR, KusumotaL. Frailty in elderly patients with chronic kidney disease under conservative treatment. *Rev da Rede Enferm do Nord*. 2016;17(3):386-392. doi:10.15253/2175-6783.2016000300012
18. NgJK-C, KwanBC-H, ChowK-M, et al. Frailty in Chinese Peritoneal Dialysis Patients: Prevalence and Prognostic Significance. *Kidney Blood Press Res*. 2016;41(6):736-745. doi:10.1159/000450563
19. MeulendijksFG, HamakerME, BoereboomFTJ, KalfA, VögtlanderNPJ, vanMunsterBC. Groningen frailty indicator in older patients with end-stage renal disease. *Ren Fail*. 2015;37(9):1419-1424.
20. MargiottaE, CaldiroliL, VettorettiS, et al. SuO004GUT MICROBIOTA COMPOSITION AND FRAILITY IN ELDERLY PATIENTS WITH CHRONIC KIDNEY DISEASE. *Nephrol Dial Transplant*. 2018;33(suppl\_1):i618-i618. doi:10.1093/ndt/gfy104.SuO004
21. DeSouza OrlandiF, GesualdoGD. Assessment of the frailty level of elderly people with chronic kidney disease undergoing hemodialysis. *ACTA Paul Enferm*. 2014;27(1):29-34. doi:10.1590/1982-0194201400007
22. SalesC, TavaresR, AmadoL, et al. SP651ANXIETY AND DEPRESSION IN END STAGE RENAL DISEASE PATIENTS AND ITS ASSOCIATION WITH CLINICAL AND LABORATORIAL DATA. *Nephrol Dial Transplant*. 2017;32(suppl\_3):iii355-iii355. doi:10.1093/ndt/gfx154.SP651
23. HaugenCE, MountfordA, WarsameF, et al. Incidence, Risk Factors, and Sequelae of Post-kidney Transplant Delirium. *J Am Soc Nephrol*.

2018;29(6):1752-1759. doi:10.1681/ASN.2018010064

24. MoffattH, MoorhouseP, MalleryL, LandryD, TennankoreK. Using the Frailty Assessment for Care Planning Tool (FACT) to screen elderly chronic kidney disease patients for frailty: the nurse experience. *Clin Interv Aging*. 2018;13:843.
25. JohansenKL, PainterP, DelgadoC, DoyleJ. Characterization of physical activity and sitting time among patients on hemodialysis using a new physical activity instrument. *J Ren Nutr*. 2015;25(1):25-30. doi:10.1053/j.jrn.2014.06.012
26. NooriN, Sharma ParpiaA, LakhaniR, JanesS, GoldsteinMB. Frailty and the Quality of Life in Hemodialysis Patients: The Importance of Waist Circumference. *J Ren Nutr*. 2018;28(2):101-109. doi:10.1053/j.jrn.2017.07.007
27. Fabrício-WehbeSCC, SchiavetoFV, VendrusculoTRP, HaasVJ, DantasRAS, RodriguesRAP. Cross-cultural adaptation and validity of the “Edmonton Frail Scale - EFS” in a Brazilian elderly sample. *Rev Lat Am Enfermagem*. 2009;17(6):1043-1049. doi:10.1590/S0104-11692009000600018
28. ChaoC-T, HuangJ-W. Frailty severity is significantly associated with electrocardiographic QRS duration in chronic dialysis patients. *PeerJ*. 2015;3:e1354. doi:10.7717/peerj.1354
29. ChaoC-T, ChiangC-K, HuangJ-W, HungK-Y. Self-reported frailty among end-stage renal disease patients: A potential predictor of dialysis access outcomes. *Nephrology (Carlton)*. 2017;22(4):333-334. doi:10.1111/nep.12961
30. ChuNM, GrossAL, ShafferAA, et al. Frailty and Changes in Cognitive Function after Kidney Transplantation. *J Am Soc Nephrol*. 2019;30(2):336-345. doi:10.1681/ASN.2018070726
31. KutnerNG, ZhangR, AllmanRM, BowlingCB. Correlates of ADL difficulty in a large hemodialysis cohort. *Hemodial Int*. 2014;18(1):70-77. doi:10.1111/hdi.12098
32. ChaoC-T, ChanD-C, HuangJ-W. Frail Phenotype Might Be Associated With Higher Appendicular but Not Truncal Fat Among End-Stage Renal Disease Patients. *J Pain Symptom Manage*. 2017;53(4):e1-e4. doi:10.1016/j.jpainsymman.2017.01.004
33. ChaoC-T, HuangJ-W, ChanD-C. Frail phenotype might herald bone health worsening among end-stage renal disease patients. *PeerJ*. 2017;5:e3542. doi:10.7717/peerj.3542
34. ChaoC-T, ChiangC-K, HuangJ-W, ChanD-C. Effect of Frail Phenotype on Bone Mass and Vertebral Compression Fracture in Individuals

Undergoing Dialysis. *J Am Geriatr Soc.* 2016;64(9):e19-21. doi:10.1111/jgs.14296

35. YonekiK, KitagawaJ, HoshiK, et al. Association between frailty and bone loss in patients undergoing maintenance hemodialysis. *J Bone Miner Metab.* January2018. doi:10.1007/s00774-017-0898-4

36. DelgadoC, DoyleJW, JohansenKL. Association of frailty with body composition among patients on hemodialysis. *J Ren Nutr.* 2013;23(5):356-362. doi:10.1053/j.jrn.2013.02.010

37. ChaoC-T, HsuY-H, ChangP-Y, et al. Simple self-report FRAIL scale might be more closely associated with dialysis complications than other frailty screening instruments in rural chronic dialysis patients. *Nephrology (Carlton).* 2015;20(5):321-328. doi:10.1111/nep.12401

38. CamilleriS, ChongS, TangvoraphonkchaiK, YoowannakulS, DavenportA. Effect of Self-Reported Distress Thermometer Score on the Maximal Handgrip and Pinch Strength Measurements in Hemodialysis Patients. *Nutr Clin Pract.* 2017;32(5):682-686. doi:10.1177/0884533617697936

39. McAdams-DeMarcoMA, YingH, OlorundareI, et al. Frailty and Health-Related Quality of Life in End Stage Renal Disease Patients of All Ages. *J frailty aging.* 2016;5(3):174-179.

40. LeeSJ, SonH, ShinSK. Influence of frailty on health-related quality of life in pre-dialysis patients with chronic kidney disease in Korea: a cross-sectional study. *Health Qual Life Outcomes.* 2015;13(1):70.

41. MansurHN, ColugnatiFA, GrincenkovFR, BastosMG. Frailty and quality of life: a cross-sectional study of Brazilian patients with pre-dialysis chronic kidney disease. *Heal Qual Life Outcomes.* 2014;12:27. doi:10.1186/1477-7525-12-27

42. IyasereOU, BrownEA, JohanssonL, et al. Quality of Life and Physical Function in Older Patients on Dialysis: A Comparison of Assisted Peritoneal Dialysis with Hemodialysis. *Clin J Am Soc Nephrol.* 2016;11(3):423-430. doi:10.2215/CJN.01050115

43. McAdams-DeMarcoMA, OlorundareIO, YingH, et al. Frailty and Postkidney Transplant Health-Related Quality of Life. *Transplantation.* 2018;102(2):291-299. doi:10.1097/TP.0000000000001943

44. KutnerNG, ZhangR, HuangY, WasseH. Falls among hemodialysis patients: potential opportunities for prevention? 2014. doi:10.1093/ckj/sfu034

45. McAdams-DeMarcoMA, SureshS, LawA, et al. Frailty and falls among adult patients undergoing chronic hemodialysis: a prospective cohort study. *BMC Nephrol.* 2013;14:224. doi:10.1186/1471-2369-14-224

46. BaoY, DalrympleL, ChertowGM, KaysenGA, JohansenKL. Frailty, dialysis initiation, and mortality in end-stage renal disease. *Arch Intern Med.* 2012;172(14):1071-1077. doi:10.1001/archinternmed.2012.3020
47. DelgadoC, ShiehS, GrimesB, et al. Association of Self-Reported Frailty with Falls and Fractures among Patients New to Dialysis. *Am J Nephrol.* 2015;42(2):134-140. doi:10.1159/000439000
48. KonelJM, WarsameF, YingH, et al. Depressive symptoms, frailty, and adverse outcomes among kidney transplant recipients. *Clin Transplant.* 2018;32(10):e13391. doi:10.1111/ctr.13391
49. LeeS-Y, YangDH, HwangE, et al. The prevalence, association, and clinical outcomes of frailty in maintenance dialysis patients. *J Ren Nutr.* 2017;27(2):106-112.
50. McAdams-DeMarcoMA, LawA, SalterML, et al. Frailty as a novel predictor of mortality and hospitalization in individuals of all ages undergoing hemodialysis. *J Am Geriatr Soc.* 2013;61(6):896-901. doi:10.1111/jgs.12266
51. McAdams-DeMarcoMA, LawA, SalterML, et al. Frailty and Early Hospital Readmission After Kidney Transplantation. *Am J Transplant.* 2013;13(8):2091-2095. doi:10.1111/ajt.12300
52. McAdams-DeMarcoMA, KingEA, LuoX, et al. Frailty, Length of Stay, and Mortality in Kidney Transplant Recipients: A National Registry and Prospective Cohort Study. *Ann Surg.* 2017;266(6):1084-1090. doi:10.1097/SLA.0000000000002025
53. ChuNM, DengA, YingH, et al. Dynamic Frailty Before Kidney Transplantation-Time of Measurement Matters. *Transplantation.* February2019. doi:10.1097/TP.0000000000002563
54. McAdams-DeMarcoMA, LawA, KingE, et al. Frailty and mortality in kidney transplant recipients. *Am J Transplant.* 2015;15(1):149-154. doi:10.1111/ajt.12992
55. AlfaadhelTA, SorokaSD, KiberdBA, LandryD, MoorhouseP, TennankoreKK. Frailty and mortality in dialysis: evaluation of a clinical frailty scale. *Clin J Am Soc Nephrol.* 2015;10(5):832-840. doi:10.2215/CJN.07760814
56. AliH, AbdelazizT, AbdelaalF, BaharaniJ. Assessment of prevalence and clinical outcome of frailty in an elderly predialysis cohort using simple tools. *Saudi J Kidney Dis Transpl.* 2018;29(1):63-70. doi:10.4103/1319-2442.225175
57. JohansenKL, DalrympleLS, GliddenD, et al. Association of Performance-Based and Self-Reported Function-Based Definitions of Frailty with



Mortality among Patients Receiving Hemodialysis. Clin J Am Soc Nephrol. 2016;11(4):626-632. doi:10.2215/CJN.03710415

58. Pérez FernándezM, Martínez MiguelP, YingH, et al. Comorbidity, Frailty, and Waitlist Mortality among Kidney Transplant Candidates of All Ages. Am J Nephrol. 2019;49(2):103-110. doi:10.1159/000496061
59. FitzpatrickJ, SozioSM, JaarBG, et al. Frailty, body composition and the risk of mortality in incident hemodialysis patients: the Predictors of Arrhythmic and Cardiovascular Risk in End Stage Renal Disease study. Nephrol Dial Transplant. 2019;34(2):346-354. doi:10.1093/ndt/gfy124
60. RomeuM, CouchoudC, DelarozierJC, et al. Survival of patients with ANCA-associated vasculitis on chronic dialysis: data from the French REIN registry from 2002 to 2011. QJM. 2014;107(7):545-555. doi:10.1093/qjmed/hcu043
61. DecourtA, GondouinB, DelarozierJC, et al. Trends in Survival and Renal Recovery in Patients with Multiple Myeloma or Light-Chain Amyloidosis on Chronic Dialysis. Clin J Am Soc Nephrol. 2016;11(3):431-441. doi:10.2215/CJN.06290615
62. LeeSW, LeeA, YuMY, et al. Is Frailty a Modifiable Risk Factor of Future Adverse Outcomes in Elderly Patients with Incident End-Stage Renal Disease? J Korean Med Sci. 2017;32(11):1800-1806. doi:10.3346/jkms.2017.32.11.1800
63. SchopmeyerL, ElMoumniM, Nieuwenhuijs-MoekeGJ, BergerSP, BakkerSJL, PolRA. Frailty has a significant influence on postoperative complications after kidney transplantation-a prospective study on short-term outcomes. Transpl Int. 2019;32(1):66-74. doi:10.1111/tri.13330