

# Week 12: Discrete scalar dynamical systems

May 20, 2024

# First order linear difference equations

We consider the first order linear homogeneous difference equation with constant coefficient

$$x_{k+1} - ax_k = 0, \quad k \in \mathbb{N} \quad (1)$$

where  $a \in \mathbb{R}$  is given, and the unknown is the sequence of real numbers  $(x_k)_{k \geq 0}$ .

Its characteristic equation is  $r - a = 0$  with the real root  $r = a$ . We associate the sequence  $a^k$  and write the general solution

$$x_k = ca^k, \quad k \in \mathbb{N},$$

with  $c \in \mathbb{R}$  an arbitrary constant.

# First order linear difference equations

Now we consider the first order linear non-homogeneous difference equation

$$x_{k+1} - ax_k = b_k, \quad k \in \mathbb{N} \quad (2)$$

where  $(b_k)_{k \geq 0}$  is a given sequence of real numbers.

It can be proved that

its general solution is the sum between a particular solution of it and the general solution of the linear homogeneous difference equation associated  $x_{k+1} - ax_k = 0$ .

## Second order linear homogeneous difference equations

We consider the second order linear homogeneous difference equation with constant coefficients

$$x_{k+2} + a_1 x_{k+1} + a_2 x_k = 0, \quad k \in \mathbb{N} \quad (3)$$

where  $a_1, a_2 \in \mathbb{R}$  are given, and the unknown is the sequence of real numbers  $(x_k)_{k \geq 0}$ . We present the characteristic equation method to find its general solution.

First note that, if we look for solutions of the form  $x_k = r^k$  we get  $r^{k+2} + a_1 r^{k+1} + a_2 r^k = 0$  for all  $k \in \mathbb{N}$ . Thus, we must have  $r^2 + a_1 r + a_2 = 0$ .

# The characteristic equation method

$$x_{k+2} + a_1 x_{k+1} + a_2 x_k = 0, \quad k \in \mathbb{N} \quad (4)$$

*Step 1.* Write the characteristic equation  $r^2 + a_1 r + a_2 = 0$  and find its roots  $r_1, r_2 \in \mathbb{C}$ .

*Step 2.* According to the nature of the roots we associate two sequences following the rules:

If  $r_{1,2} \in \mathbb{R}$  and  $r_1 \neq r_2$  then we associate  $r_1^k$  and  $r_2^k$ .

If  $r_1 = r_2 \in \mathbb{R}$  then we associate  $r_1^k$  and  $kr_1^k$ .

If  $r_1 = \overline{r_2} \in \mathbb{C} \setminus \mathbb{R}$  then we associate the real part and, respectively, the imaginary part of  $r_1^k$ .

*Step 3.* We write the general solution as a linear combination with arbitrary coefficients of the two sequences found at Step 2.

# Scalar discrete dynamical systems

For a given map  $f : \mathbb{R} \rightarrow \mathbb{R}$ , starting with an initial value  $x_0 = \eta \in \mathbb{R}$ , we construct a sequence  $(x_k)_{k \geq 0}$  such that

$$x_{k+1} = f(x_k).$$

This is said to be the positive orbit of the initial state  $\eta$ ; or the sequence of iterates of  $f$  that starts with  $\eta$ .

**Notation:**  $f^2 = f \circ f$ ,  $f^3 = f \circ f \circ f$  and  $f^k$  = the  $k$  times composition of  $f$  with itself. Then  $x_k = f^k(\eta)$ .

We are interested in studying the long-term behavior of each orbit.  
We will always assume at least that  $f$  is continuous.

The simplest sequence is a constant one and it is obtained when the initial value is a fixed point of  $f$ , i.e.  $f(\eta) = \eta$ .

## Fixed points - an exercise

We consider the logistic map  $f_\lambda : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f_\lambda(x) = \lambda x(1 - x)$ , where  $\lambda \in [1, 4]$  is a fixed parameter. Our aim is to find the fixed points of  $f$ .  $\diamond$

We have to solve the equation  $x = f_\lambda(x) \iff x = \lambda x(1 - x) \iff \lambda x^2 - (\lambda - 1)x = 0$ . This equation has two roots, which means that  $f_\lambda$  has two fixed points  $\eta_1^* = 0$  and  $\eta_2^* = \frac{\lambda-1}{\lambda}$ .  $\square$

In particular, this implies that the unique solution of the IVP

- (a)  $x_{k+1} = \lambda x_k(1 - x_k)$ ,  $x_0 = 0$  is  $x_k = 0$  for all  $k \geq 0$ .
- (b)  $x_{k+1} = 2x_k(1 - x_k)$ ,  $x_0 = 0.5$  is  $x_k = 0.5$  for all  $k \geq 0$ .

Graphically, the fixed points of  $f$  are found at the intersection between the graph of  $f$  and the first bisectrix (i.e. the line  $y = x$ ).

# Scalar discrete dynamical systems

Another simple behavior is when **the sequences is convergent**, i.e. there exists  $\eta^* \in \mathbb{R}$  such that  $x_k \rightarrow \eta^*$  when  $k \rightarrow \infty$ .

In this case, using also the continuity of  $f$ , we have  $x_{k+1} \rightarrow \eta^*$  and  $f(x_k) \rightarrow f(\eta^*)$  when  $k \rightarrow \infty$ .

Since  $x_{k+1} = f(x_k)$  we deduce that  $\eta^* = f(\eta^*)$ . Thus, **The limit of a convergent sequence of iterates of  $f$  is a fixed point of  $f$ .**

We say that a fixed point  $\eta^*$  is an **attractor** of  $f$  when there exists  $\rho > 0$  such that for each  $\eta$  with  $|\eta - \eta^*| \leq \rho$  we have  $f^k(\eta) \rightarrow \eta^*$  when  $k \rightarrow \infty$ . **The basin of attraction** of an attractor  $\eta^*$  is defined as

$$A_{\eta^*} = \{\eta \in \mathbb{R} : f^k(\eta) \rightarrow \eta^* \text{ when } k \rightarrow \infty\}.$$

When  $A_{\eta^*} = \mathbb{R}$  we say that  $\eta^*$  is a **global attractor** of  $f$ .



# A linear map

We consider  $f(x) = ax$ , where  $a \in \mathbb{R} \setminus \{-1, 0, 1\}$  and the associated linear difference equation

$$x_{k+1} = ax_k.$$

It is easy to see that the only fixed point of  $f$  is  $\eta^* = 0$  and  $x_k = f^k(\eta) = a^k \eta$  for any  $\eta \in \mathbb{R}$ ,  $k \in \mathbb{N}$ .

If  $|a| < 1$  then  $\eta^* = 0$  is a global attractor of  $f$ .

If  $|a| > 1$  then  $f^k(\eta)$  is unbounded for any  $\eta \neq 0$ .

# The linearization method

## Theorem

Assume that  $f$  is a  $C^1$  function and let  $\eta^*$  be a fixed point of  $f$ .

If  $|f'(\eta^*)| < 1$  then  $\eta^*$  is an attractor.

If  $|f'(\eta^*)| > 1$  then  $\eta^*$  is not an attractor.

Exercise: Apply the linearization method to the fixed points of the logistic map  $f_\lambda(x) = \lambda x(1 - x)$  where  $\lambda \in [1, 4]$  is a parameter.  $\diamond$

We found that  $f_\lambda$  has two fixed points  $\eta_1^* = 0$  and  $\eta_2^* = \frac{\lambda-1}{\lambda}$ .

We compute the derivative of  $f_\lambda(x) = \lambda x - \lambda x^2$  and obtain

$$f'_\lambda(x) = \lambda - 2\lambda x.$$

Then  $|f'_\lambda(\eta_1^*)| = \lambda$  and  $|f'_\lambda(\eta_2^*)| = |\lambda - 2(\lambda - 1)| = |2 - \lambda|$ .

The above theorem only assures that:

If  $\lambda \in (1, 3)$  then  $\eta_1^* = 0$  is not an attractor, but  $\eta_2^*$  is an attractor.

If  $\lambda \in (3, 4]$  then neither of the two fixed points is an attractor.

## The cobweb or stair-step diagram

The cobweb diagram is an intuitive geometric way of showing the behavior of the orbits  $x_k = f^k(\eta)$  of a discrete dynamical system.

The algorithm works as follows for  $k = 0, 1, 2, \dots$ :

Beginning with  $x_k$ , move vertically to find the point on the graph of the map  $f$  that corresponds to it, i.e. the point of coordinates  $(x_k, f(x_k))$ .

Then move horizontally to a point on the line  $y = x$ , i.e the point of coordinates  $(f(x_k), f(x_k))$ .

Repeat as many times as needed.

The cobweb diagram for the logistic map can be better visualized here:

<http://sites.saintmarys.edu/~sbroad/example-logistic-cobweb.html>

<https://www.geogebra.org/m/gHYqKMSJ>

We proved that, when  $\lambda \in (1, 3)$  the fixed point  $\eta_2^* = \frac{\lambda-1}{\lambda}$  is an attractor. It is important to find its basin of attraction  $A_2$ .

After we have seen, it seems that  $A_2 = (0, 1)$ .

In fact, this is mathematically proved. (see Elaydi: Discrete chaos, 2008)

## Periodic points

We will employ again the notation  $f^k = f \circ f \circ \dots \circ f$ , the  $k$  times composition of the map  $f$  with itself. We start by noting that a fixed point of  $f$  is also a fixed point of  $f^k$  for any  $k \in \mathbb{N}$ .

### Definition

Let  $\eta^* \in \mathbb{R}$  and  $p \in \mathbb{N}$ ,  $p \geq 2$ . We say that  $\eta^*$  is a  $p$ -periodic point of  $f$  when  $\eta^*$  is a fixed point of  $f^p$  but  $\eta^*$  is not a fixed point of  $f^{p-1}, \dots, f^2, f$ .

**Remark.** If  $\eta^*$  is a  $p$ -periodic point of  $f$  then the unique solution of the IVP  $x_{k+1} = f(x_k)$ ,  $x_0 = \eta^*$  is a sequence whose first  $p$  terms are  $\eta^*, f(\eta^*), \dots, f^{p-1}(\eta^*)$  and then they are repeated. Such an orbit is also called a  $p$ -cycle.

**Exercise:** Check that  $\{\frac{1+\sqrt{5}}{2}, \frac{1-\sqrt{5}}{2}\}$  is a 2-cycle of the map  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = 1 - 2x^2$ .  $\diamond$

We compute  $f\left(\frac{1+\sqrt{5}}{2}\right) = \frac{1-\sqrt{5}}{2}$  and  $f^2\left(\frac{1+\sqrt{5}}{2}\right) = f\left(\frac{1-\sqrt{5}}{2}\right) = \frac{1+\sqrt{5}}{2}$ .  $\square$

## Periodic points - an exercise

Exercise: Find the fixed points and the 2-periodic points of  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x) = 1 - 2x^2$ .

To find the fixed points we solve the equation  $x = f(x) \Leftrightarrow 2x^2 + x - 1 = 0 \Leftrightarrow (2x - 1)(x + 1) = 0$ .

Thus,  $f$  has two fixed points:  $\eta_1^* = 1/2$  and  $\eta_2^* = 1$ .

To find the fixed points we first compute  $f^2 = f \circ f$ , then find the fixed points of  $f^2$ , i.e. solve the equation  $x = f^2(x)$ . We will use that the fixed points of  $f$  are also fixed points of  $f^2$ .

We have

$$f^2(x) = f(f(x)) = 1 - 2f(x)^2 = 1 - 2(1 - 2x^2)^2 = 1 - 2(1 - 4x^2 + 4x^4). \\ \text{Then } f^2(x) = -8x^4 + 8x^2 - 1 \text{ for all } x \in \mathbb{R}.$$

The equation  $x = f^2(x) \Leftrightarrow 8x^4 - 8x^2 + x + 1 = 0$ . This is a polynomial equation of degree 4. We have that the fixed points of  $f$  ( $1/2$  and  $1$ ) are roots of this equation. Since

$8x^4 - 8x^2 + x + 1 = (2x^2 + x - 1)(4x^2 - 2x - 1)$ , the 2-periodic points are the roots of  $4x^2 - 2x - 1 = 0$ . Thus, they are  $\frac{1+\sqrt{5}}{2}$  and  $\frac{1-\sqrt{5}}{2}$ .

# Attracting cycle

## Definition

Let  $\eta^*$  be a  $p$ -periodic point of  $f$ . We say that the corresponding  $p$ -cycle is an attracting cycle of  $f$  when  $\eta^*$  is an attracting fixed point of  $f^p$ . The basin of attraction of the fixed point  $\eta^*$  of  $f^p$  is said to be the basin of attraction of the  $p$ -cycle.

**Remark 1.** Let  $\eta^*$  be a  $p$ -periodic point of  $f$  such that the corresponding  $p$ -cycle is an attracting cycle of  $f$ . Let  $\eta$  be in the basin of attraction of the fixed point  $\eta^*$  of  $f^p$ . Then the unique solution of the IVP  $x_{k+1} = f(x_k)$ ,  $x_0 = \eta$  is a sequence that can be split into  $p$  convergent sub-sequences. The limits are the terms of the attracting  $p$ -cycle, i.e.  $\eta^*, f(\eta^*), \dots, f^{p-1}(\eta^*)$ .

**Remark 2.** What we see in the cobweb-diagram for the logistic map with  $\lambda = 3.15$  (for example) suggests us that there exists an attracting 2-cycle having the interval  $(0, 1)$  as its basin of attraction.

Note that it is very difficult (impossible) to detect (using the cobweb diagram) a cycle which is not attracting.

# The linearization method for a 2-cycle

## Theorem

*Let  $\{\eta^*, f(\eta^*)\}$  be a 2-cycle of  $f$ .*

*If  $|f'(\eta^*)f'(f(\eta^*))| < 1$  then the cycle is an attractor.*

*If  $|f'(\eta^*)f'(f(\eta^*))| > 1$  then the cycle is not an attractor.*

Proof. Using the definition of an attracting cycle and the linearization method for fixed points, the following statements are valid.

If  $|(f^2)'(\eta^*)| < 1$  then the cycle is an attractor.

If  $|(f^2)'(\eta^*)| > 1$  then the cycle is not an attractor.

The proof is done after we notice that

$$(f^2)'(x) = \frac{d}{dx}f(f(x)) = f'(f(x))f'(x) \text{ for all } x \in \mathbb{R}. \quad \square$$

Exercise. Prove that neither the fixed points, nor the 2-cycle of  $f(x) = 1 - 2x^2$  are attractors.  $\diamond$

# Exercises

1. We consider the map

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = \frac{1}{50}x(100 - x).$$

- (a) Find its fixed points and study their stability.
- (b) Using the stair-step diagram estimate the basin of attraction of the attractor fixed point.
- (c) If  $(x_k)_{k \geq 0}$  represent the number of fish in some lake at month  $k$  and

$$x_{k+1} = \frac{1}{50}x_k(100 - x_k), \quad x_0 = \eta$$

try to predict the fate of the fish in the case  $\eta = 80$  and also in the case  $\eta = 10$ .  $\diamond$



# Exercises

2. Using the stair-step diagram, estimate the basin of attraction for each of the fixed points (if there is any which is an attractor) of the map

$$f : (0, \infty) \rightarrow \mathbb{R}, \quad f(x) = \frac{x^2 + 5}{2x}.$$

All the exercises have been solved during the lecture.