• Linear Gaussian Estimation

assumption: discrete, linear, time varying

• motion model:

$$x_k = A_{k-1} x_{k-1} + v_k + w_k$$

(x_{k-1}: random; v_k: input, deterministic; w_k: process noise, random)

$$y_k = C_k x_k + n_k$$

$x_0 \in \mathbb{R}^N \sim \mathcal{N}(\check{x}_0, \check{P}_0)$

$w_k \in \mathbb{R}^N \sim \mathcal{N}(0, Q_k)$

$n_k \in \mathbb{R}^M \sim \mathcal{N}(0, R_k)$

• batch linear-Gaussian estimation problem
- Bayesian
- Maximum a Posteriori

▲ Maximum a Posteriori

• $\hat{x} = \underset{x}{\arg\max}\ P(x \mid v, y)$

$x = x_{0:K} = (x_0, x_1, \ldots, x_K)$

$v = (\check{x}_0, v_{1:K}) = (\check{x}_0, v_1, \ldots, v_K)$ } for prior

$y = y_{0:K} = (y_0, \ldots, y_K)$ } for posterior

• Bayes' rule:

$\hat{x} = \underset{x}{\arg\max}\ P(x \mid v, y)$ (does not affect y)

$= \underset{x}{\arg\max} \dfrac{P(y \mid x, v) P(x \mid v)}{P(y \mid v)}$ → does not affect x

$= \underset{x}{\arg\max}\ P(y \mid x) P(x \mid v)$

$P(A \mid B) = \dfrac{P(A \cap B)}{P(B)}$

$P(B \mid A) = \dfrac{P(A \cap B)}{P(A)}$

$\therefore P(A \cap B) = P(B \mid A) P(A)$

$\therefore P(A \mid B) = \dfrac{P(B \mid A) P(A)}{P(B)}$ ★

$P(A \mid BC) = \dfrac{P(A \cap B \cap C)}{P(B \cap C)}$

$P(C \mid AB) = \dfrac{P(A \cap B \cap C)}{P(A \cap B)}$

$\therefore P(A \cap B \cap C) = P(A \cap B) P(C \mid A \cap B) = P(A \mid B) P(B) P(C \mid A \cap B)$

$\therefore P(A \mid BC) = \dfrac{P(A \mid B) P(B) P(C \mid A \cap B)}{P(B) P(C \mid B)} = \dfrac{P(A \mid B) P(C \mid A \cap B)}{P(C \mid B)}$

• assume each w_k, n_k are NOT correlated:

$$P(y \mid x) = \prod_{k=0}^{K} P(y_k \mid x_k)$$

• Bayes' Rule:

$$P(x \mid v) = P(x_0 \mid \check{x}_0) \prod_{k=0}^{K} P(x_k \mid x_{k-1}, v_k)$$

$\therefore P(x_0 \mid \check{x}_0) = \dfrac{1}{\sqrt{(2\pi)^N \det \check{P}_0}} \times \exp\left(-\frac{1}{2}(x_0 - \check{x}_0)^T \check{P}_0^{-1} (x_0 - \check{x}_0)\right)$

$\therefore P(x_k \mid x_{k-1}, v_k) = \dfrac{1}{\sqrt{(2\pi)^N \det Q_k}} \times \exp\left(-\frac{1}{2}(x_k - (A_{k-1}x_{k-1} + v_k))^T Q_k^{-1} (x_k - (A_{k-1}x_{k-1} + v_k))\right)$

$\therefore P(y_k \mid x_k) = \dfrac{1}{\sqrt{(2\pi)^M \det R_k}} \times \exp\left(-\frac{1}{2}(y_k - C_k x_k)^T R_k^{-1} (y_k - C_k x_k)\right)$

• logarithm

$\ln(P(y \mid x) P(x \mid v))$

$= \ln P(x_0 \mid \check{x}_0) + \sum_{k=1}^{K} \ln P(x_k \mid x_{k-1}, v_k) + \sum_{k=1}^{K} \ln P(y_k \mid x_k)$

where

$\ln P(x_0 \mid \check{x}_0) = -\frac{1}{2}(x_0 - \check{x}_0)^T \check{P}_0^{-1}(x_0 - \check{x}_0) + -\frac{1}{2}\ln((2\pi)^N \det \check{P}_0)$

$\ln P(x_k \mid x_{k-1}, v_k) = -\frac{1}{2}(x_k - A_{k-1}x_{k-1} - v_k)^T Q_k^{-1}(x_k - A_{k-1}x_{k-1} - v_k) + -\frac{1}{2}\ln((2\pi)^N \det Q_k)$

$\ln P(y_k \mid x_k) = -\frac{1}{2}(y_k - C_k x_k)^T R_k^{-1}(y_k - C_k x_k) + -\frac{1}{2}\ln((2\pi)^M \det R_k)$

• J, cost

$J_{v,k}(x) = \begin{cases} \frac{1}{2}(x_0 - \check{x}_0)^T \check{P}_0^{-1}(x_0 - \check{x}_0) \\ \frac{1}{2}(x_k - A_{k-1}x_{k-1} - v_k)^T Q_k^{-1}(x_k - A_{k-1}x_{k-1} - v_k) \end{cases}$

$J_{y,k}(x) = \frac{1}{2}(y_k - C_k x_k)^T R_k^{-1}(y_k - C_k x_k)$

Mahalanobis distance

$$J(x) = \sum_{k=0}^{K}(J_{v,k}(x) + J_{y,k}(x))$$

$$\therefore x = \underset{x}{\arg\min}\ J(x)$$

• from previous

$J_{v,k}(x) = \begin{cases} \frac{1}{2}(x_0 - \check{x}_0)^T \check{P}_0^{-1}(x_0 - \check{x}_0) \\ \frac{1}{2}(x_k - A_{k-1}x_{k-1} - v_k)^T Q_k^{-1}(x_k - A_{k-1}x_{k-1} - v_k) \end{cases}$

$J_{y,k}(x) = \frac{1}{2}(y_k - C_k x_k)^T R_k^{-1}(y_k - C_k x_k)$

let $z = \begin{bmatrix} \check{x}_0 \\ v_1 \\ \vdots \\ v_K \\ \hline y_0 \\ y_1 \\ \vdots \\ y_K \end{bmatrix}$, $x = \begin{bmatrix} x_0 \\ \vdots \\ x_K \end{bmatrix}$

& $H = \begin{bmatrix} 1 & & \\ -A_0 & 1 & \\ & \ddots & \ddots \\ & -A_{K-1} & 1 \\ \hline C_0 & & \\ & C_1 & \\ & & \ddots \\ & & C_K \end{bmatrix}$

$W = \begin{bmatrix} \check{P}_0 & & & \\ & Q_1 \cdots Q_K & & \\ & & R_0 & \\ & & & R_1 \cdots R_K \end{bmatrix}$

$$\therefore J(x) = \frac{1}{2}(z - Hx)^T W^{-1} (z - Hx)$$

$P(z \mid x) = \eta \exp\left(-\frac{1}{2}(z - Hx)^T W^{-1}(z - Hx)\right)$ (↳ normalization constant)

$\therefore \left.\dfrac{\partial J(x)}{\partial x^T}\right|_{\hat{x}} = -H^T W^{-1}(z - H\hat{x}) = 0$

$$\Rightarrow (H^T W^{-1} H)\hat{x} = H^T W^{-1} z$$

▲ Bayesian Inference

compute full Bayesian Posterior

$P(x \mid v, y)$

• $x_k = A_{k-1}x_{k-1} + v_k + w_k$

lift → $x = A(v + w)$

where

$A = \begin{bmatrix} 1 & & & & \\ A_0 & 1 & & & \\ A_1 A_0 & A_1 & 1 & & \\ \vdots & \vdots & \vdots & \ddots & \\ A_{K-2}\cdots A_0 & A_{K-2}\cdots A_1 & A_{K-2}\cdots A_2 & \cdots & 1 \\ A_{K-1}\cdots A_0 & A_{K-1}\cdots A_1 & A_{K-1}\cdots A_2 & \cdots A_{K-1} & 1 \end{bmatrix}$

• $\check{x} = E[x] = E[A(v + w)] = E[Av + Aw] = Av$

• $\check{P} = E[(x - E[x])(x - E[x])^T] = AQA^T$

$Q = E[ww^T] = \text{diag}(\check{P}_0, Q_1, \ldots, Q_K)$

• prior ∴ ⇒

$P(x \mid v) = \mathcal{N}(\check{x}, \check{P}) = \mathcal{N}(Av, AQA^T)$

• $y_k = C_k x_k + n_k$

lift → $y = Cx + n$

where

$C = \text{diag}(C_0, C_1, \ldots, C_K)$

• joint density of prior lifted state & measurements is now:

$$P(x, y \mid v) = \mathcal{N}\left(\begin{bmatrix} \check{x} \\ C\check{x} \end{bmatrix}, \begin{bmatrix} \check{P} & \check{P}C^T \\ C\check{P} & C\check{P}C^T + R \end{bmatrix}\right)$$

$R = E[nn^T] = \text{diag}(R_0, R_1, \ldots, R)$

• $P(x, y \mid v) = P(x \mid y, v) P(y \mid v)$ — we only care about this ($P(x \mid y, v)$)

- $P(x, y) = \mathcal{N}\left(\begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}, \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{bmatrix}\right)$
- $P(x, y) = P(x \mid y) P(y)$

w/ schur complement

$\begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{bmatrix} = \begin{bmatrix} I & \Sigma_{xy}\Sigma_{yy}^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} \Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx} & 0 \\ 0 & \Sigma_{yy} \end{bmatrix} \begin{bmatrix} I & 0 \\ \Sigma_{yy}^{-1}\Sigma_{yx} & I \end{bmatrix}$

$\Rightarrow \begin{bmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{bmatrix}^{-1} = \begin{bmatrix} I & 0 \\ -\Sigma_{yy}^{-1}\Sigma_{yx} & I \end{bmatrix} \begin{bmatrix} (\Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx})^{-1} & 0 \\ 0 & \Sigma_{yy}^{-1} \end{bmatrix} \begin{bmatrix} I & -\Sigma_{xy}\Sigma_{yy}^{-1} \\ 0 & I \end{bmatrix}$

- PDF of $P(x, y)$

recall: $P(x) = (2\pi)^{-\frac{k}{2}} \det(\Sigma)^{-\frac{1}{2}} \exp\left(-\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)\right)$

$\Rightarrow P(x, y) = (2\pi)^{-\frac{k}{2}} \det(\Sigma)^{-\frac{1}{2}} \exp(Z)$

$Z = \left(\begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}\right)^T \begin{pmatrix} \Sigma_{xx} & \Sigma_{xy} \\ \Sigma_{yx} & \Sigma_{yy} \end{pmatrix}^{-1} \left(\begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}\right)$

$= \left(\begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}\right)^T \begin{bmatrix} I & 0 \\ -\Sigma_{yy}^{-1}\Sigma_{yx} & I \end{bmatrix} \begin{bmatrix} (\Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx})^{-1} & 0 \\ 0 & \Sigma_{yy}^{-1} \end{bmatrix} \begin{bmatrix} I & -\Sigma_{xy}\Sigma_{yy}^{-1} \\ 0 & I \end{bmatrix} \left(\begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} \mu_x \\ \mu_y \end{bmatrix}\right)$

$= [x - \mu_x - \Sigma_{xy}\Sigma_{yy}^{-1}(y - \mu_y)]^T (\Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx})^{-1} \times [x - \mu_x - \Sigma_{xy}\Sigma_{yy}^{-1}(y - \mu_y)] + (y - \mu_y)^T \Sigma_{yy}^{-1}(y - \mu_y)$

sum of two quadratic terms

} $P(x, y) = P(x \mid y) P(y)$

$P(x \mid y) = \mathcal{N}(\mu_x + \Sigma_{xy}\Sigma_{yy}^{-1}(y - \mu_y),\ \Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx})$

$P(y) = \mathcal{N}(\mu_y, \Sigma_{yy})$

▲ on $(H^T W^{-1} H)\hat{x} = H^T W^{-1} z$

• unique solution:

$\hat{x} = (H^T W^{-1} H)^{-1} H^T W^{-1} z$ — need to be invertible

w/ $\hat{x} \in \mathbb{R}^{K+1}$

rank $(H^T W^{-1} H) = K+1$

⇒ rank $(H^T H)$ = rank (H^T)

• rank (H^T): — we hv $\check{x}_0$ / — we dn't hv $\check{x}_0$

✓ - we hv $\check{x}_0$

rank H^T

$= \text{rank} \left[\begin{array}{cccc|cccc} 1 & -A_0^T & & & C_0^T & & & \\ & 1 & -A_1^T & & & C_1^T & & \\ & & \ddots & -A_{K-1}^T & & & C_2^T & \\ & & & 1 & & & & C_K^T \end{array}\right]$

⇒ full rank. ∴ $Q_k \succ 0, R_k \succ 0$ ⇒ hv sol.

✓ - we dn't hv $\check{x}_0$

rank H^T

$= \text{rank} \left[\begin{array}{c|c} -A_0^T & C_0^T \\ \hline 1 \quad -A_1^T & C_1^T \\ \quad 1 \ \ddots \ -A_{K-1}^T & \quad C_2^T \\ \qquad\quad 1 & \qquad C_K^T \end{array}\right]$

$\Rightarrow \text{rank} \left[\begin{array}{c|c} 1 \quad -A_1^T & C_1^T \\ \quad 1 \ \ddots \ -A_{K-1}^T & \quad C_2^T \\ \qquad\quad 1 & \qquad C_K^T \\ \hline -A_0^T & C_0^T \end{array}\right]$

$\Rightarrow \text{rank} \left[\begin{array}{c|c} 1 \quad -A_1^T & C_1^T \\ \quad 1 \ \ddots \ -A_{K-1}^T & \quad C_2^T \ \ddots \\ \qquad\quad 1 & \qquad C_K^T \\ \hline & C_0^T \ A_0^T C_1^T \ A_0^T A_1^T C_2^T \cdots A_0^T \cdots A_{K-1}^T C_K^T \end{array}\right]$ → make sure this full rank

• rank $[C_0^T \ A_0^T C_1^T \ A_0^T A_1^T C_2^T \cdots A_0^T \cdots A_{K-1}^T C_K^T] = N$

from Cayley-Hamilton theorem: any power of $A \in \mathbb{R}^{N \times N}$, power $k \geq N$

$\Rightarrow (A^T)^{k-1} = a_0 I^T + a_1 A^T + a_2 A^T A^T + \cdots + a_{N-1}(A^T)^{N-1}$

Assume $A_0 = A_1 = A_2 = \cdots = A_{K-1} = A$ & $k >> N$

we first hv

$(A^T)^{k-1} C^T = a_0 I^T C^T + a_1 A^T C^T + a_2 A^T A^T C^T + \cdots + a_{N-1}(A^T)^{N-1} C^T$

rank $[C_0^T \ A_0^T C_1^T \ A_0^T A_1^T C_2^T \cdots A_0^T \cdots A_{K-1}^T C_K^T]$

⇒ rank $[C^T \ A^T C^T \ A^T A^T C^T \cdots (A^T)^K C^T]$

= rank $[C^T \ A^T C^T \cdots (A^T)^{N-1} C^T]$

• $\mathcal{O} = \begin{bmatrix} C \\ CA \\ \vdots \\ CA^{(N-1)} \end{bmatrix}$

• rank $\mathcal{O} = N$ (observability)

∴ $Q_k \succ 0$, $R_k \succ 0$, rank $\mathcal{O} = N$
⇒ hv sol.

$$P(x, y \mid v) = P(x \mid y, v) P(y \mid v)$$

$\therefore P(x \mid y, v)$

recall $P(x \mid y) = \mathcal{N}(\mu_x + \Sigma_{xy}\Sigma_{yy}^{-1}(y - \mu_y),\ \Sigma_{xx} - \Sigma_{xy}\Sigma_{yy}^{-1}\Sigma_{yx})$

&

$P(x, y \mid v) = \mathcal{N}\left(\begin{bmatrix} \check{x} \\ C\check{x} \end{bmatrix}, \begin{bmatrix} \check{P} & \check{P}C^T \\ C\check{P} & C\check{P}C^T + R \end{bmatrix}\right)$

$= \mathcal{N}\left(\check{x} + \check{P}C^T(C\check{P}C^T + R)^{-1}(y - C\check{x}),\ \check{P} - \check{P}C^T(C\check{P}C^T + R)^{-1}(C\check{P})\right)$

thru Sherman-Morrison-Woodbury Identity

$= \mathcal{N}\left((\check{P}^{-1} - C^T R^{-1} C)^{-1}(\check{P}^{-1}\check{x} + C^T R^{-1} y),\ (\check{P}^{-1} + C^T R^{-1} C)^{-1}\right)$

i.e., $\hat{x} = (\check{P}^{-1} - C^T R^{-1} C)^{-1}(\check{P}^{-1}\check{x} + C^T R^{-1} y)$

$\Rightarrow (\check{P}^{-1} - C^T R^{-1} C)\hat{x} = \check{P}^{-1}\check{x} + C^T R^{-1} y$

$\Rightarrow ((AQA^T)^{-1} - C^T R^{-1} C)\hat{x} = (AQA^T)^{-1} Av + C^T R^{-1} y$

$\Rightarrow (A^{-T} Q^{-1} A^{-1} - C^T R^{-1} C)\hat{x} = A^{-T} Q^{-1} A^{-1} Av + C^T R^{-1} y$

$\Rightarrow (A^{-T} Q^{-1} A^{-1} - C^T R^{-1} C)\hat{x} = A^{-T} Q^{-1} v + C^T R^{-1} y$

$A^{-1} = \begin{bmatrix} 1 & & & & \\ -A_0 & 1 & & & \\ & -A_1 & 1 & & \\ & & -A_2 & \ddots & \\ & & & & 1 \\ & & & -A_{K-1} & 1 \end{bmatrix}$, $z = \begin{bmatrix} v \\ y \end{bmatrix}$, $H = \begin{bmatrix} A^{-1} \\ C \end{bmatrix}$, $W = \begin{bmatrix} Q & \\ & R \end{bmatrix}$

$$\Rightarrow (H^T W^{-1} H)\hat{x} = H^T W^{-1} z$$

▲ solving $(H^T W^{-1} H)\hat{x} = H^T W^{-1} z$

• Cholesky Decomposition

$H^T W^{-1} H = LL^T$ — Cholesky Decomp.

- first decompose this
- $Ax = b$
  $LL^T x = b$
  ⇒ $Ly = b$ (↓ forward substitution)
- $L^T x = y$ (↑ backward substitution)
- solve x!

$\therefore (H^T W^{-1} H)\hat{x} = H^T W^{-1} z$

$\Rightarrow LL^T \hat{x} = H^T W^{-1} z$

$\Rightarrow Ld = H^T W^{-1} z$

$\Rightarrow L^T \hat{x} = d$

• let $L = \begin{bmatrix} L_0 & & & & \\ L_{10} & L_1 & & & \\ & L_{21} & L_2 & & \\ & & \ddots & \ddots & \\ & & & L_{K-1,K-2} & L_{K-1} \\ & & & & L_{K,K-1} & L_K \end{bmatrix}$

LL^T

$L_0 L_0^T = \underbrace{\check{P}_0^{-1} + C_0^T R_0^{-1} C_0}_{I_0} + A_0^T Q_1^{-1} A_0$

$L_{10} L_0^T = -Q_1^{-1} A_0$

$L_1 L_1^T = -L_{10} L_{10}^T + Q_1^{-1} + C_1^T R_1^{-1} C_1 + A_1^T Q_2^{-1} A_1$

$L_{21} L_1^T = -Q_2^{-1} A_1$

$\vdots$

$L_{K-1} L_{K-1}^T = \underbrace{-L_{K-1,K-2} L_{K-1,K-2}^T + Q_{K-1}^{-1} + C_{K-1}^T R_{K-1}^{-1} C_{K-1}}_{I_{K-1}} + A_{K-1}^T Q_K^{-1} A_{K-1}$

$L_{K,K-1} L_{K-1}^T = -Q_K^{-1} A_{K-1}$

$L_K L_K^T = \underbrace{-L_{K,K-1} L_{K,K-1}^T + Q_K^{-1} + C_K^T R_K^{-1} C_K}_{I_K}$

$I_0, I_1, \ldots, I_{K-1}, I_K$ ⇒ Information Matrix

Basically solve L_0 → solve L_{10} → solve L_1 ⋮ $O(N(K+1))$

forward

• $Ld = H^T W^{-1} z$ get $d = [d_0\ d_1 \ldots d_K]^T$

$L_0 d_0 = \underbrace{\check{P}_0^{-1}\check{x}_0 + C_0^T R_0^{-1} y_0}_{q_0} - A_0^T Q_1^{-1} v_1$

$L_1 d_1 = \underbrace{-L_{10} d_0 + Q_1^{-1} v_1 + C_1^T R_1^{-1} y_1}_{q_1} - A_1^T Q_2^{-1} v_2$

$\vdots$

$L_{K-1} d_{K-1} = \underbrace{-L_{K-1,K-2} d_{K-2} + Q_{K-1}^{-1} v_{K-1} + C_{K-1}^T R_{K-1}^{-1} y_{K-1}}_{q_{K-1}} - A_{K-1}^T Q_K^{-1} v_K$

$L_K d_K = \underbrace{-L_{K,K-1} d_{K-1} + Q_K^{-1} v_K + C_K^T R_K^{-1} y_K}_{q_K}$

Similarly, solve d_0 → solve d_1 → solve d_2 ⋮ $O(N(K+1))$

backward

• $L^T \hat{x} = d$

$L_K^T \hat{x}_K = d_K$

$L_{K-1}^T \hat{x}_{K-1} = -L_{K,K-1}^T \hat{x}_K + d_{K-1}$

$\vdots$

$L_1^T \hat{x}_1 = -L_{21}^T \hat{x}_2 + d_1$

$L_0^T \hat{x}_0 = -L_{10}^T \hat{x}_1 + d_0$

Similarly: solve $\hat{x}_K$ → solve $\hat{x}_{K-1}$ → solve $\hat{x}_{K-2}$ ⋮ $O(N(K+1))$

▲ Forward

$L_{K-1} L_{K-1}^T = I_{K-1} + A_{K-1}^T Q_K^{-1} A_{K-1}$

$L_{K-1} d_{K-1} = q_{K-1} - A_{K-1}^T Q_K^{-1} v_K$

$L_{K,K-1} L_{K-1}^T = -Q_K^{-1} A_{K-1}$

$I_K = -L_{K,K-1} L_{K,K-1}^T + Q_K^{-1} + C_K^T R_K^{-1} C_K$

$q_K = -L_{K,K-1} d_{K-1} + Q_K^{-1} v_K + C_K^T R_K^{-1} y_K$

▲ Backward

$L_{K-1}^T \hat{x}_{K-1} = -L_{K,K-1}^T \hat{x}_K + d_{K-1}$

→ Kalman filter

△ solving $(H^TW^{-1}H)\hat{x} = H^TW^{-1}z$

- **Rauch-Tung-Striebel**

from above:

$L_{k-1}L_{k-1}^T = I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1}$

$L_{k,k-1}L_{k-1}^T = -Q_k^{-1}A_{k-1}$

$I_k = -L_{k,k-1}L_{k,k-1}^T + Q_k^{-1} + C_k^TR_k^{-1}C_k$

$\therefore L_{k,k-1} = -Q_k^{-1}A_{k-1}L_{k-1}^{-T}$

$L_{k,k-1}L_{k,k-1}^T = Q_k^{-1}A_{k-1}L_{k-1}^{-T}L_{k-1}^{-1}A_{k-1}^TQ_k^{-T}$

$= Q_k^{-1}A_{k-1}(L_{k-1}L_{k-1}^T)^{-1}A_{k-1}^TQ_k^{-T}$

$\therefore I_k = -L_{k,k-1}L_{k,k-1}^T + Q_k^{-1} + C_k^TR_k^{-1}C_k$

$= Q_k^{-1} - Q_k^{-1}A_{k-1}(I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})A_{k-1}^TQ_k^{-T} + C_k^TR_k^{-1}C_k$

$= (A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1} + C_k^TR_k^{-1}C_k$

from Sherman-Morrison-Woodbury

$I_k = (A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1} + C_k^TR_k^{-1}C_k$

$\hat{P}_{k,f} = I_k^{-1}$

$\check{P}_{k,f} = A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k$

$\hat{P}_{k,f}^{-1} = \check{P}_{k,f}^{-1} + C_k^TR_k^{-1}C_k$

△ inducing KF

let us define

$K_k = \hat{P}_{k,f}C_k^TR_k^{-1}$

$= (\check{P}_{k,f}^{-1} + C_k^TR_k^{-1}C_k)^{-1}C_k^TR_k^{-1}$ } SMW

$= \check{P}_{k,f}C_k^T(C_k\check{P}_{k,f}C_k^T + R_k)^{-1}$

which is the Kalman gain that we are familiar of

- $\check{P}_{k,f}^{-1} = \hat{P}_{k,f}^{-1} - C_k^TR_k^{-1}C_k$

$= \hat{P}_{k,f}^{-1}[1 - \hat{P}_{k,f}C_k^TR_k^{-1}C_k)$

$= \hat{P}_{k,f}^{-1}(1 - K_kC_k)$

$\Rightarrow \hat{P}_{k,f} = (1 - K_kC_k)\check{P}_{k,f}$

- now,

from above,

$$\begin{cases} L_{k,k-1}L_{k-1}^T = -Q_k^{-1}A_{k-1} \\ L_{k-1}d_{k-1} = q_{k-1} - A_{k-1}^TQ_k^{-1}v_k \\ L_{k-1}L_{k-1}^T = I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1} \end{cases}$$

$L_{k,k-1} = (-Q_k^{-1}A_{k-1})L_{k-1}^{-T}$

$d_{k-1} = L_{k-1}^{-1}(q_{k-1} - A_{k-1}^TQ_k^{-1}v_k)$

$\Rightarrow L_{k,k-1}d_{k-1} = (-Q_k^{-1}A_{k-1})L_{k-1}^{-T}L_{k-1}^{-1} \times (q_{k-1} - A_{k-1}^TQ_k^{-1}v_k)$

$\Rightarrow L_{k,k-1}d_{k-1} = (-Q_k^{-1}A_{k-1})(L_{k-1}L_{k-1}^T)^{-1}(q_{k-1} - A_{k-1}^TQ_k^{-1}v_k)$ (1)

$\because q_k = -L_{k,k-1}d_{k-1} + Q_k^{-1}v_k + C_k^TR_k^{-1}y_k$ (2)

combining (1) & (2)

$q_k = Q_k^{-1}A_{k-1}(I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1}q_{k-1}$
$+ (Q_k^{-1} - Q_k^{-1}A_{k-1}(I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1}A_{k-1}^TQ_k^{-1})v_k$
$+ C_k^TR_k^{-1}y_k$ thru SMW

$= (A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1}A_{k-1}I_{k-1}^{-1}q_{k-1}$
$+ (A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1}v_k$
$+ C_k^TR_k^{-1}y_k$

$= (A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1}(A_{k-1}I_{k-1}^{-1}q_{k-1} + v_k)$
$+ C_k^TR_k^{-1}y_k$

$= (\check{P}_{k,f})^{-1}(A_{k-1}\hat{P}_{k-1,f}\hat{P}_{k-1,f}^{-1}\hat{x}_{k-1,f} + v_k)$
$+ C_k^TR_k^{-1}y_k$ let $q_k = \hat{P}_{k,f}^{-1}\hat{x}_{k,f}$

$\Rightarrow \hat{P}_{k,f}^{-1}\hat{x}_{k,f} = \check{P}_{k,f}^{-1}(A_{k-1}\hat{x}_{k-1,f} + v_k) + C_k^TR_k^{-1}y_k$

$\Rightarrow \hat{P}_{k,f}^{-1}\hat{x}_{k,f} = \check{P}_{k,f}^{-1}\check{x}_{k,f} + C_k^TR_k^{-1}y_k$

$\Rightarrow \hat{x}_{k,f} = \hat{P}_{k,f}\check{P}_{k,f}^{-1}\check{x}_{k,f} + \hat{P}_{k,f}C_k^TR_k^{-1}y_k$

$= (1 - K_kC_k)\check{x}_{k,f} + K_ky_k$

$\Rightarrow \hat{x}_{k,f} = \check{x}_{k,f} + K_k(y_k - C_k\check{x}_{k,f})$

end of "forward"

- backward

from above

$L_{k-1}^T\hat{x}_{k-1} = -L_{k,k-1}^T\hat{x}_k + d_{k-1}$

$\Rightarrow \hat{x}_{k-1} = (L_{k-1}^T)^{-1}(-L_{k,k-1}^T\hat{x}_k + d_{k-1})$

$= L_{k-1}^{-T}L_{k-1}^{-1}L_{k-1}(-L_{k,k-1}^T\hat{x}_k + d_{k-1})$

$= (L_{k-1}L_{k-1}^T)^{-1}L_{k-1}(-L_{k,k-1}^T\hat{x}_k + d_{k-1})$

$\Rightarrow$ from

$L_{k-1}L_{k-1}^T = I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1}$

$L_{k-1}d_{k-1} = q_{k-1} - A_{k-1}^TQ_k^{-1}v_k$

$L_{k,k-1}L_{k-1}^T = -Q_k^{-1}A_{k-1}$

$= (L_{k-1}L_{k-1}^T)^{-1}(-L_{k-1}L_{k,k-1}^T\hat{x}_k + L_{k-1}d_{k-1})$

$= (I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1} \times (-(L_{k,k-1}L_{k-1}^T)^T\hat{x}_k + q_{k-1} - A_{k-1}^TQ_k^{-1}v_k)$

$= (I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1} \times (A_{k-1}^TQ_k^{-1}\hat{x}_k + q_{k-1} - A_{k-1}^TQ_k^{-1}v_k)$

$= (I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1} \times (A_{k-1}^TQ_k^{-1}(\hat{x}_k - v_k) + q_{k-1})$

$= (I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1}A_{k-1}^TQ_k^{-1}(\hat{x}_k - v_k)$
$+ (I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1}q_{k-1}$

$(I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1}A_{k-1}^TQ_k^{-1}(\hat{x}_k - v_k)$
$+ (I_{k-1} + A_{k-1}^TQ_k^{-1}A_{k-1})^{-1}q_{k-1}$ } thru SMW

$= I_{k-1}^{-1}A_{k-1}^T(A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1}(\hat{x}_k - v_k)$
$+ (I_{k-1}^{-1} - I_{k-1}^{-1}A_{k-1}^T(A_{k-1}I_{k-1}^{-1}A_{k-1}^T + Q_k)^{-1}A_{k-1}I_{k-1}^{-1})q_{k-1}$

$= \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}(\hat{x}_k - v_k)$
$+ (\hat{P}_{k-1,f} - \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}A_{k-1}\hat{P}_{k-1,f})\hat{P}_{k-1,f}^{-1}\hat{x}_{k-1,f}$

$= \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}(\hat{x}_k - v_k)$
$+ \hat{x}_{k-1,f} - \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}A_{k-1}\hat{P}_{k-1,f}\hat{P}_{k-1,f}^{-1}\hat{x}_{k-1,f}$

$= \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}(\hat{x}_k - v_k)$
$+ \hat{x}_{k-1,f} - \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}\check{x}_{k,f}$

$= \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}(\hat{x}_k - \check{x}_{k,f} - v_k) + \hat{x}_{k-1,f}$

$= \hat{P}_{k-1,f}A_{k-1}^T\check{P}_{k,f}^{-1}(\hat{x}_k - \check{x}_{k,f}) + \hat{x}_{k-1,f}$

Non-linear Estimations

Recall

J, cost

$$J_{v,k}(x) = \begin{cases} \frac{1}{2}(x_0 - \check{x}_0)^T\check{P}_0^{-1}(x_0 - \check{x}_0) \\ \frac{1}{2}(x_k - A_{k-1}x_{k-1} - v_k)^TQ_k^{-1}(x_k - A_{k-1}x_{k-1} - v_k) \end{cases}$$

$J_{y,k}(x) = \frac{1}{2}(y_k - C_kx_k)^TR_k^{-1}(y_k - C_kx_k)$

Mahalanobis distance

$$J(x) = \sum_{k=0}^{K}(J_{v,k}(x) + J_{y,k}(x))$$

$\therefore x = \underset{x}{\operatorname{argmin}}\ J(x)$

- here the system, i.e.,
  - the prediction model
  - the measurement model

  is no longer linear

$$e_{v,k}(x) = \begin{cases} \check{x}_0 - x_0 & , k=0 \\ f(x_{k-1}, v_k, 0) - x_k & , k=1 \sim K \end{cases}$$

$e_{y,k}(x) = y_k - g(x_k, 0)$, $k = 0 \sim K$

$J_{v,k}(x) = \frac{1}{2}e_{v,k}(x)^TW_{v,k}^{-1}e_{v,k}(x)$

$J_{y,k}(x) = \frac{1}{2}e_{y,k}(x)^TW_{y,k}^{-1}e_{y,k}(x)$

$\Rightarrow J(x) = \sum_{k=0}^{K}(J_{v,k}(x) + J_{y,k}(x))$

$\Rightarrow$ • let $e(x) = \begin{bmatrix} e_v(x) \\ e_y(x) \end{bmatrix}$

$e_v(x) = \begin{bmatrix} e_{v,0}(x) \\ \vdots \\ e_{v,K}(x) \end{bmatrix}$

$e_y(x) = \begin{bmatrix} e_{y,0}(x) \\ \vdots \\ e_{y,K}(x) \end{bmatrix}$

- let $W = \operatorname{diag}(W_v, W_y)$

  $W_v = \operatorname{diag}(W_{v,0}, \dots W_{v,K})$

  $W_y = \operatorname{diag}(W_{y,0}, \dots W_{y,K})$

- $J(x) = \frac{1}{2}e(x)^TW^{-1}e(x)$

$\Rightarrow$ let $u(x) = Le(x)$

$L^TL = W^{-1}$

$J(x) = \frac{1}{2}e(x)^TL^TLe(x)$

$= \frac{1}{2}[Le(x)]^TLe(x)$

$= \frac{1}{2}u(x)^Tu(x)$

$\Rightarrow \hat{x} = \underset{x}{\operatorname{argmin}}\ J(x)$

Solving $\hat{x} = \underset{x}{\operatorname{argmin}}\ J(x)$

- Newton's Method

@ update step

$J(x_{op} + \delta x)$

$\approx J(x_{op}) + \left(\frac{\partial J(x)}{\partial x}\Big|_{x_{op}}\right)\delta x + \frac{1}{2}\delta x^T\left(\frac{\partial^2 J(x)}{\partial x\partial x^T}\Big|_{x_{op}}\right)\delta x$

$\frac{\partial J(x_{op} + \delta x)}{\partial \delta x}$

$= \left(\frac{\partial J(x)}{\partial x}\Big|_{x_{op}}\right) + \delta x^{*T}\left(\frac{\partial^2 J(x)}{\partial x\partial x^T}\Big|_{x_{op}}\right) = 0$

$\Rightarrow \left(\frac{\partial^2 J(x)}{\partial x\partial x^T}\Big|_{x_{op}}\right)\delta x^* = -\left(\frac{\partial J(x)}{\partial x}\Big|_{x_{op}}\right)^T$ (δx^* → update step)

then

$x_{op} \leftarrow x_{op} + \delta x^*$

- Gauss-Newton Method

from above

- $J(x) = \frac{1}{2}u(x)^Tu(x)$
- Jacobian

  $\frac{\partial J(x)}{\partial x}\Big|_{x_{op}} = u(x_{op})^T\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)$

- Hessian

  $\frac{\partial^2 J(x)}{\partial x\partial x^T}\Big|_{x_{op}}$

  $= \left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)^T\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right) + \sum_{i=1}^{M} u_i(x_{op})\left(\frac{\partial^2 u_i(x)}{\partial x\partial x^T}\Big|_{x_{op}}\right)$ (neglect this)

  $\approx \left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)^T\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)$

$\therefore \left(\frac{\partial^2 J(x)}{\partial x\partial x^T}\Big|_{x_{op}}\right)\delta x^* = -\left(\frac{\partial J(x)}{\partial x}\Big|_{x_{op}}\right)^T$

$\Rightarrow \left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)^T\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)\delta x^* = -\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)^Tu(x_{op})$

△ GN method

recall:

- $\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)^T\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)\delta x^* = -\left(\frac{\partial u(x)}{\partial x}\Big|_{x_{op}}\right)^Tu(x_{op})$
- $J(x) = \frac{1}{2}e(x)^TW^{-1}e(x)$

  $= \frac{1}{2}e(x)^TL^TLe(x)$

  $= \frac{1}{2}[Le(x)]^TLe(x)$

  $= \frac{1}{2}u(x)^Tu(x)$

- $$e_{v,k}(x) = \begin{cases} \check{x}_0 - x_0 & , k=0 \\ f(x_{k-1}, v_k, 0) - x_k & , k=1 \sim K \end{cases}$$

  $e_{y,k}(x) = y_k - g(x_k, 0)$, $k = 0 \sim K$

NOW:

- for any function $f(x)$, perturbed by δx @ a we can write Taylor expansion: (2nd order)

$f(a + \delta x)$

$= f(a) + \frac{\partial f(x)}{\partial x}\Big|_{x=a}\delta x + \frac{1}{2}\delta x^T\frac{\partial^2 f(x)}{\partial x\partial x^T}\Big|_{x=a}\delta x$

$\therefore$ w/ 1st order

$$e_{v,k}(x_{op} + \delta x) \approx \begin{cases} e_{v,0}(x_{op,0}) + \frac{\partial e_{v,0}(x)}{\partial x}\Big|_{x_{op,0}}\delta x, & k=0 \\ e_{v,k}(x_{op,k}) + \frac{\partial e_{v,k}(x)}{\partial x}\Big|_{x_{op,k}}\delta x, & k\neq 0 \end{cases}$$

$$\Rightarrow \begin{cases} ① \ \check{x}_0 - x_{op,0} - 1\cdot\delta x & , k=0 \\ ② \ f(x_{op,k-1}, v_k, 0) - x_{op,k} + F_{k-1}\delta x_{k-1} - \delta x_k, & k\neq 0 \end{cases}$$

$F_{k-1} = \frac{\partial f(x_{k-1}, v_k, w_k)}{\partial x_{k-1}}\Big|_{x_{op,k-1}, v_k, 0}$

$e_{y,k}(x_{op} + \delta x) \approx e_{y,k}(x_{op}) + \frac{\partial e_{y,k}(x)}{\partial x}\Big|_{x_{op,k}}$

$\Rightarrow$ ③ $y_k - g(x_{op,k}, 0) - G_k\delta x_k$

$G_k = \frac{\partial g(x_k, n_k)}{\partial x_k}\Big|_{x_{op,k}, 0}$

THEN

$J(x) = \frac{1}{2}e(x)^TW^{-1}e(x)$

- stack all $e(x)$'s

$$e(x_{op}) = \begin{bmatrix} e_{v,0}(x_{op}) \\ e_{v,1}(x_{op}) \\ \vdots \\ e_{v,K}(x_{op}) \\ \hline e_{y,0}(x_{op}) \\ e_{y,1}(x_{op}) \\ \vdots \\ e_{y,K}(x_{op}) \end{bmatrix}$$

- WHILE in iterative method, variable is δx

$$\delta x = \begin{bmatrix} \delta x_0 \\ \delta x_1 \\ \vdots \\ \delta x_K \end{bmatrix}$$

- ALSO, recall GN:

$J(x)^TJ(x)\delta x = -J(x)^Tf(x)$

HERE

$$\therefore J(x) = \begin{bmatrix} \frac{\partial f_1(x)}{\partial x_1} & \dots & \frac{\partial f_1(x)}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n(x)}{\partial x_1} & \dots & \frac{\partial f_n(x)}{\partial x_n} \end{bmatrix}$$

$$\text{"}f(x)\text{"} = e(x_{op}) = \begin{bmatrix} e_{v,0}(x_{op}) \\ e_{v,1}(x_{op}) \\ \vdots \\ e_{v,K}(x_{op}) \\ \hline e_{y,0}(x_{op}) \\ e_{y,1}(x_{op}) \\ \vdots \\ e_{y,K}(x_{op}) \end{bmatrix}$$

$$\text{"}x\text{"} = \delta x = \begin{bmatrix} \delta x_0 \\ \delta x_1 \\ \delta x_2 \\ \vdots \\ \delta x_K \end{bmatrix}$$

"$J(x)$"

$= J$

$$= \begin{bmatrix} \frac{\partial e_{v,0}(x_{op})}{\partial \delta x_0} & \frac{\partial e_{v,0}(x_{op})}{\partial \delta x_1} & \dots \\ \frac{\partial e_{v,1}(x_{op})}{\partial \delta x_0} & \frac{\partial e_{v,1}(x_{op})}{\partial \delta x_1} & \dots \\ \vdots & \frac{\partial e_{v,2}(x_{op})}{\partial \delta x_2} & \ddots \\ \hline \frac{\partial e_{y,0}(x_{op})}{\partial \delta x_0} & \frac{\partial e_{y,0}(x_{op})}{\partial \delta x_1} & \dots \\ \frac{\partial e_{y,1}(x_{op})}{\partial \delta x_0} & \frac{\partial e_{y,1}(x_{op})}{\partial \delta x_1} & \dots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

$$= \begin{bmatrix} 1 & & & \\ F_0 & 1 & & \\ & F_1 & \ddots & \\ & & F_{K-1} & 1 \\ \hline -G_0 & & & \\ & -G_1 & & \\ & & \ddots & \end{bmatrix}$$

$(J^TW^{-1}J)\delta x = -J^TW^{-1}e(x_{op})$

remember $u(x) = Le(x)$ previously, I dropped "L"

@ iteration step i

solve δx

THEN

$x_{op} \leftarrow x_{op} + \delta x$

★ solving the linear system @ i, has similar formulation w/ LGE

★ NLGE ⟷ LGE

- iterative linearization
- solve the linear system locally @ i
- solve the linear system globally

★ $\delta x^k = \underset{\delta x}{\operatorname{argmin}} \sum \|e - J\delta x\|_\Lambda^2$

Marginalization – Sliding Window

recall the linear system

$H\delta x = b$

ALSO

recall Schur complement

$X = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}$

$S = C - B^TA^{-1}C$

→ Schur complement of A in X

- The core idea of marginalization in SWF:

- $H\delta x = b$ (we want this out: δx_a)

$\Rightarrow \begin{bmatrix} \Lambda_a & \Lambda_b \\ \Lambda_b^T & \Lambda_c \end{bmatrix}\begin{bmatrix} \delta x_a \\ \delta x_b \end{bmatrix} = \begin{bmatrix} g_a \\ g_b \end{bmatrix}$

$\Rightarrow \begin{bmatrix} \Lambda_a & \Lambda_b \\ \Lambda_b^T - \Lambda_b^T\Lambda_a^{-1}\Lambda_a & \Lambda_c - \Lambda_b^T\Lambda_a^{-1}\Lambda_b \end{bmatrix}\begin{bmatrix} \delta x_a \\ \delta x_b \end{bmatrix} = \begin{bmatrix} g_a \\ g_b - \Lambda_b^T\Lambda_a^{-1}g_a \end{bmatrix}$

$\Rightarrow \begin{bmatrix} \Lambda_a & \Lambda_b \\ 0 & \Lambda_c - \Lambda_b^T\Lambda_a^{-1}\Lambda_b \end{bmatrix}\begin{bmatrix} \delta x_a \\ \delta x_b \end{bmatrix} = \begin{bmatrix} g_a \\ g_b - \Lambda_b^T\Lambda_a^{-1}g_a \end{bmatrix}$

$\Rightarrow (\Lambda_c - \Lambda_b^T\Lambda_a^{-1}\Lambda_b)\delta x_b = g_b - \Lambda_b^T\Lambda_a^{-1}g_a$

→ we get marginalized state!

ALSO! sparse → dense

- in pose graph optimization
- Generally H matrix has following structure



→ this is sparse!

$\Rightarrow \begin{bmatrix} B & E \\ E^T & C \end{bmatrix}\begin{bmatrix} \Delta x_c \\ \Delta x_p \end{bmatrix} = \begin{bmatrix} v \\ w \end{bmatrix}$ (Δx_c: camera pose, Δx_p: 3D pts)

$\Rightarrow \begin{bmatrix} B - EC^{-1}E^T & E - EC^{-1}C \\ E^T & C \end{bmatrix}\begin{bmatrix} \Delta x_c \\ \Delta x_p \end{bmatrix} = \begin{bmatrix} v - EC^{-1}w \\ w \end{bmatrix}$

$\Rightarrow \begin{bmatrix} B - EC^{-1}E^T & 0 \\ E^T & C \end{bmatrix}\begin{bmatrix} \Delta x_c \\ \Delta x_p \end{bmatrix} = \begin{bmatrix} v - EC^{-1}w \\ w \end{bmatrix}$

$$\begin{cases} (B - EC^{-1}E^T)\Delta x_c = v - EC^{-1}w \\ E^T\Delta x_c + C\Delta x_p = w \end{cases}$$

→ C^{-1} (diagnol) ⇒ easy to solve then solve Δx_p!