

Bayesian leave-one-out cross-validation for non-factorizable normal models*

Paul-Christian Bürkner[†]

Jonah Gabry[‡]

Aki Vehtari[§]

February 28, 2019

Abstract

Cross-validation can be used to measure a model’s predictive accuracy for the purpose of model comparison, averaging, or selection. Standard leave-one-out cross-validation (LOO-CV) requires the likelihood to be factorizable, but many important models in temporal and spatial statistics do not have this property. We derive how to compute and validate both exact and approximate LOO-CV for Bayesian non-factorizable models with a multivariate normal likelihood.

Keywords: cross-validation, Pareto-smoothed importance-sampling, non-factorizable models.

1. Introduction

In the absence of new data, cross-validation is a general approach for evaluating a statistical model’s predictive accuracy for the purpose of model comparison, averaging, or selection (Geisser and Eddy, 1979; Hoeting et al., 1999; Ando and Tsay, 2010; Vehtari and Ojanen, 2012). One widely used variant of cross-validation is *leave-one-out cross-validation* (LOO-CV), where observations are left out one at a time and then predicted based on the model fit to the remaining data. Predictive accuracy is evaluated by first computing the expected log predictive density of the left-out observation and then taking the sum of these values over all observations to obtain the expected log predictive density (ELPD) as a single measure of predictive accuracy. Unfortunately, exact LOO-CV is costly, as it requires fitting the model as many times as there are observations in the data. Depending on the size of the data, complexity of the model, and estimation method, this can be practically infeasible as it simply requires too much computation time. For this reason, approximate versions of LOO-CV have been developed (Gelfand et al., 1992; Vehtari et al., 2017b), most recently using Pareto-smoothed importance-sampling (PSIS, Vehtari et al. (2017b,a)).

A standard assumption of any such LOO-CV approach is that the joint likelihood of the model over all observations has to be factorizable. That is, the observations have to be pairwise conditionally independent given the model parameters. However, many important models do not have this property. Particularly in temporal and spatial statistics it is common to fit models with multivariate normal likelihoods that have structured covariance matrices such that the likelihood does not factorize. This is typically due to the fact that observations depend on other observations

*We thank Daniel Simpson for useful discussion and the Academy of Finland (grants 298742, 313122) for partial support of this work.

[†]Department of Psychology, University of Münster, Germany.

[‡]Institute for Social and Economic Research and Policy, Columbia University.

[§]Department of Computer Science, Aalto University, Finland.

from different time periods or different spatial units in addition to the dependence on the model parameters.

In this short paper we show how equations derived in Sundararajan and Keerthi (2001) can be repurposed and combined with PSIS to allow for performing efficient approximate LOO-CV for *any* multivariate normal Bayesian model with an invertible covariance matrix, regardless of whether or not the likelihood factorizes. We also provide equations for computing exact LOO-CV for these models, which can be used to validate the approximation. Throughout, a Bayesian model specification and estimation via Markov chain Monte Carlo (MCMC) is assumed. In online supplementary material we provide R code demonstrating how to carry out the approximation described in the paper as well as comparisons between the approximate results and results from exact LOO-CV.¹

2. Pointwise log-likelihood for non-factorizable normal models

When computing *exact* LOO-CV for a Bayesian model we need to compute the log leave-one-out predictive densities $\log p(y_i | y_{-i})$ for every response value y_i , $i = 1, \dots, N$, where y_{-i} denotes all response values except observation i . This requires fitting the model N times. For *approximate* LOO-CV using only a single model fit, we instead calculate the pointwise log-likelihood (the log-predictive density evaluated at each data point), without leaving out any observations, and then apply an importance sampling correction (Gelfand et al., 1992; Vehtari et al., 2017b).

The pointwise log-likelihood is straightforward to compute for *factorizable* models in which response values are conditionally independent given the model parameters θ and the likelihood can be written in the familiar form

$$p(y | \theta) = \prod_{i=1}^N p(y_i | \theta). \quad (1)$$

When $p(y)$ can be factorized in this way, the conditional pointwise log-likelihood can be obtained easily by computing $\log p(y_i | \theta)$ for each i .

The situation is more complicated for *non-factorizable* models in which response values are not conditionally independent. When there is residual dependence even after accounting for the model parameters, the conditional pointwise log-likelihood has the general form $\log p(y_i | y_{-i}, \theta)$. Computing this pointwise log-likelihood for non-factorizable models is often impossible, but there is a large class of multivariate normal models for which an analytical solution is available.

Sundararajan and Keerthi (2001) provide equations for the predictive mean and standard deviation for a zero-mean Gaussian process model with prior covariance K and residual standard deviation σ ,

$$y \sim N(0, K + \sigma^2 I), \quad (2)$$

¹Supplemental materials available at <https://mc-stan.org/loo/articles/loo2-non-factorizable.html>.

where I is the identity matrix of appropriate dimension and $C = K + \sigma^2 I$ is the covariance matrix of the model. These equations were not traditionally intended to be used for LOO-CV of Gaussian process models because, in most cases, Gaussian processes are combined with a factorizable likelihood so that simpler equations for univariate distributions can be applied. But the derivations of Sundararajan and Keerthi's equations make no use of the special form of C for Gaussian process models and thus immediately generalize to the case of an arbitrary invertible covariance matrix C .

For such models the LOO predictive mean and standard deviation can be computed using the equations from Sundararajan and Keerthi (2001) as follows:

$$\begin{aligned}\mu_{\tilde{y},-i} &= y_i - \bar{c}_{ii}^{-1} g_i \\ \sigma_{\tilde{y},-i} &= \sqrt{\bar{c}_{ii}^{-1}},\end{aligned}\tag{3}$$

where $g_i = [C^{-1}y]_i$ and $\bar{c}_{ii} = [C^{-1}]_{ii}$. The log predictive density of the i th observation is

$$\log p(y_i | y_{-i}, \theta) = -\frac{1}{2} \log(2\pi) - \frac{1}{2} \log \sigma_{-i}^2 - \frac{1}{2} \frac{(y_i - \mu_{-i})^2}{\sigma_{-i}^2},\tag{4}$$

and expressing this same equation in terms of g_i and \bar{c}_{ii} , we obtain²:

$$\log p(y_i | y_{-i}, \theta) = -\frac{1}{2} \log(2\pi) + \frac{1}{2} \log \bar{c}_{ii} - \frac{1}{2} \frac{g_i^2}{\bar{c}_{ii}}.\tag{5}$$

Evaluating equation (5) for each y_i provides the pointwise log-likelihood values required for the PSIS-LOO-CV approximation.

3. Approximate LOO-CV for non-factorizable normal models

The conditional pointwise log-likelihood is the only required input to the PSIS-LOO-CV algorithm from Vehtari et al. (2017b) and thus Sundararajan and Keerthi's repurposed equations allow for approximate LOO-CV for *any* model that can be expressed conditionally in terms of a multivariate normal with invertible covariance matrix C , including those where the likelihood does not factorize. For a Bayesian model fit using MCMC the procedure is as follows:

1. Fit the model using MCMC obtaining S samples from the posterior distribution of the parameters θ .
2. For each of the S draws of θ , compute the pointwise log-likelihood value for each of the N observations in y using (5). The results can be stored in an $S \times N$ matrix.
3. Run the PSIS algorithm from Vehtari et al. (2017b) on the $S \times N$ matrix obtained in step 2. For convenience the `loo` R package (Vehtari et al., 2018) provides this functionality.

In the supplementary materials we demonstrate this method by computing approximate LOO-CV for a lagged simultaneously autoregressive (SAR) model fit to spatially correlated crime data.

²Note that Vehtari et al. (2016) has a typo in the corresponding Equation 34.

4. Validation using exact LOO-CV

In order to validate the approximate LOO-CV procedure, and also in order to allow exact computations to be made for a small number of leave-one-out folds for which the Pareto- k diagnostic (Vehtari et al., 2017a) indicates an unstable approximation, we need to consider how we might do *exact* LOO-CV for a non-factorizable model. Here we will provide the necessary equations and in the supplementary materials we provide code for comparing the exact and approximate versions.

In the case of a Gaussian process that has the marginalization property, exact LOO-CV is relatively straightforward: when refitting the model we can simply drop the one row and column of the covariance matrix C corresponding to the held out observation. But this does not hold in general for multivariate normal models, and to keep the original prior we may need to maintain the full covariance matrix C even when one of the observations is left out.

The general solution is to model y_i as a missing observation and estimate it along with all of the other model parameters. For a multivariate normal model $\log p(y_i | y_{-i})$ can be computed as follows. First, we model y_i as missing and denote the corresponding parameter y_i^{mis} . Then, we define

$$y_{\text{mis}(i)} = (y_1, \dots, y_{i-1}, y_i^{\text{mis}}, y_{i+1}, \dots, y_N). \quad (6)$$

to be the same as the full set of observations y but replacing y_i with the *parameter* y_i^{mis} .

Second, we compute the LOO predictive means and standard deviations using the equations from section 2, but replacing y with $y_{\text{mis}(i)}$ in the computation of $\mu_{\tilde{y}, -i}$:

$$\mu_{\tilde{y}, -i} = y_{\text{mis}(i)} - \bar{c}_{ii}^{-1} g_i, \quad (7)$$

where in this case we have

$$g_i = [C^{-1} y_{\text{mis}(i)}]_i. \quad (8)$$

The conditional log predictive density is then computed with the above $\mu_{\tilde{y}, -i}$ and the left out observation y_i :

$$\log p(y_i | y_{-i}, \theta) = -\frac{1}{2} \log(2\pi) - \frac{1}{2} \log \sigma_{\tilde{y}, -i}^2 - \frac{1}{2} \frac{(y_i - \mu_{\tilde{y}, -i})^2}{\sigma_{\tilde{y}, -i}^2}. \quad (9)$$

Finally, the leave-one-out predictive distribution can be estimated as

$$p(y_i | y_{-i}) \approx \frac{1}{S} \sum_{s=1}^S p(y_i | y_{-i}, \theta_{-i}^{(s)}), \quad (10)$$

where $\theta_{-i}^{(s)}$ are draws from the posterior distribution $p(\theta | y_{\text{mis}(i)})$.

5. Conclusion

We have provided equations that enable both approximate and exact LOO-CV for non-factorizable multivariate normal Bayesian models. Although exact LOO-CV is usually impractical, our exact

LOO-CV procedure can be used to validate the more efficient PSIS-LOO-CV approximation, as we show in the supplementary materials.

The primary motivation for this paper is to enable approximate LOO-CV for models that cannot be factorized at all, but our approach also works for *any* Bayesian model that can be expressed in terms of a multivariate normal likelihood. Therefore it may also be useful for models that are factorizable but for which the factorized representation is difficult to compute or not available to the researcher for some other reason.

References

- Ando, T. and R. Tsay (2010). Predictive likelihood for Bayesian model selection and averaging. *International Journal of Forecasting* 26(4), 744–763.
- Geisser, S. and W. F. Eddy (1979). A predictive approach to model selection. *Journal of the American Statistical Association* 74(365), 153–160.
- Gelfand, A., D. Dey, and H. Chang (1992, 01). Model determination using predictive distributions with implementation via sampling-based methods. *Bayesian Statistics* 4, 147–167.
- Hoeting, J. A., D. Madigan, A. E. Raftery, and C. T. Volinsky (1999, 11). Bayesian model averaging: a tutorial. *Statist. Sci.* 14(4), 382–417.
- Sundararajan, S. and S. S. Keerthi (2001). Predictive approaches for choosing hyperparameters in Gaussian processes. *Neural Computation* 13(5), 1103–1118.
- Vehtari, A., J. Gabry, Y. Yao, and A. Gelman (2018). **loo**: *Efficient Leave-One-Out Cross-Validation and WAIC for Bayesian Models*. R package version 2.0.0.
- Vehtari, A., A. Gelman, and J. Gabry (2017a). Pareto smoothed importance sampling. *arXiv preprint*.
- Vehtari, A., A. Gelman, and J. Gabry (2017b). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing* 27(5), 1413–1432.
- Vehtari, A., T. Mononen, V. Tolvanen, T. Sivula, and O. Winther (2016). Bayesian leave-one-out cross-validation approximations for Gaussian latent variable models. *Journal of Machine Learning Research* 17(103), 1–38.
- Vehtari, A. and J. Ojanen (2012). A survey of bayesian predictive methods for model assessment, selection and comparison. *Statistics Surveys* 6, 142–228.