Tracking biological ageing to support healthier careers in emergency services

CRICOS No. 00213J

Firefighters showed a greater decline in Hannum age acceleration than paramedics over 12 months.

-2.33 years

01 Study and analysis

Emergency responder professions differ in their exposures. We therefore examined whether epigenetic ageing changed differently by profession.

332 participants at baseline



Psychosocial assessments
+ saliva DNA methylation

Four epigenetic clocks

Horvath	**Hannum [1]**
PhenoAge	**GrimAge**

MethylationEPICv2 data were analysed with mixed-effects models: Timepoint × Profession and participant random intercepts.

Adjusted for sex, ethnicity, trauma exposure, BMI and estimated cell composition.

02 Hannum change over time

Only Hannum showed evidence that trajectories differed by profession. EMD did not differ significantly from paramedics.




Figure 1. Model-estimated Hannum trajectories. EMD = emergency medical dispatchers. Adjusted analysis: 464 observations from 330 participants; p-values test individual Timepoint × Profession coefficients.

03 Biological processes

We then explored biological processes linked to differences in methylation change over time between firefighters and paramedics.




Figure 2. GOstats used a heat-response reference universe and CpGs selected at nominal p < 0.05.

04 Interpretation

Possible immune recovery

Blood studies link Hannum to immune cell composition. Therefore, recovery from recent immune activation is plausible [3].

Psychological factors

Psychological symptoms and positive factors showed no association across the four clocks. However, subtle changes in DNAm-based ageing over 12 months may have limited the detection of associations.

Exploratory pathways

GOmeth tests of time and the omnibus Timepoint × Profession interaction found no GO/KEGG terms at FDR < 0.05 [2].

Next step

Longer follow-up with exposure measurements, inflammatory markers and repeated methylation assessments could distinguish transient responses from sustained epigenetic change.

Paul Ruiz Pinto[a,b], Mackenzie Rubens[a,b], Dagmar Bruenig[a,b], Jane Shakespeare-Finch[c], Caroline Nievergelt[d,e,f], Marcus Ising[g], Divya Mehta[a,b]

a Centre for Genomics and Personalised Health, Faculty of Health, Queensland University of Technology, Brisbane, Australia
b School of Biomedical Sciences, Faculty of Health, Queensland University of Technology, Brisbane, Australia
c School of Psychology and Counselling, Faculty of Health, Queensland University of Technology, Brisbane, Australia
d Department of Psychiatry, University of California San Diego, La Jolla, USA
e VA San Diego Health Care System, San Diego, USA
f VA Center of Excellence for Stress and Mental Health, VA San Diego Health Care System, San Diego, USA
g Max Planck Institute of Psychiatry, Munich, Germany

We thank the study participants, participating emergency-service organisations, Jessica May Adams and Zoe Christofis. We acknowledge QUT's administrative support.

[1] Hannum G et al. Molecular Cell. 2013;49:359-367. [2] Maksimovic J et al. Genome Biology. 2021;22:173.
[3] Zhang Z et al. Aging Cell. 2024;23:e14071.

Funding: NHMRC Ideas Grant 2004536

Contact: paul.ruizpinto@qut.edu.au; divya.mehta@qut.edu.au

Download poster

Connect on LinkedIn