

# Formulation for The Merry Movie Montage

Paula Fermín Cueto

December 2021

Sets:

|           |   |
|-----------|---|
| $A$       | Set of activities (either a movie, a wildcard or no movie). $A = \{\emptyset, *, A, B, C, D, E, F, G\}$ |
| $M$       | Set of movies. $M = \{A, B, C, D, E, F, G\}$  |
| $T$       | Set of time periods. $T = 1 \dots T_{max}$ .  |
| $G$       | Set of groups of elves wathing movies. $G = \{1, 2, 3\}$ .  |
| $P$       | Set of movie permutations.  |
| $P_{all}$ | Set of movie permutations that must be watched by all teams.  |

Parameters:

|               |  |
|---------------|--|
| $ P $         | Number of permutations of $ M $ movies. $ P  =  M ! = 7! = 5,040$ .                  |
| $ P_{all} $   | Number of permutations starting with $AB$ . $ P_{all}  =  M - 2 ! = 5! = 120$ .      |
| $T_{max}$     | Upper bound on the length of the longest schedule.                                   |
|               | $T_{max} = \frac{1}{ G }( M  - 1)!(2 M  - 1) + \frac{2}{ G }( M  - 2)! M  = 3,680$ . |
| $\omega_{pk}$ | Movie in position $k \in 1 \dots  M $ of permutation $p \in P$ .                     |

Decision variables:

|                |   |
|----------------|---|
| $x_{gta}$      | $= \begin{cases} 1 & \text{if group } g \in G \text{ performs activity } a \in A \text{ at time } t \in T. \\ 0 & \text{otherwise.} \end{cases}$                |
| $\delta_{pgt}$ | $= \begin{cases} 1 & \text{if permutation } p \in P \text{ is found in group } g \in G \text{ starting at time } t \in T. \\ 0 & \text{otherwise.} \end{cases}$ |
| $\gamma_{pg}$  | $= \begin{cases} 1 & \text{if permutation } p \in P \text{ is found in group } g \in G. \\ 0 & \text{otherwise.} \end{cases}$                                   |
| $\ell$         | Duration of the longest schedule of the three teams.  |

$$\min_{x, \delta, \gamma, \ell} \quad \ell \tag{1}$$

$$\text{s.t.} \quad \ell \geq \sum_{\substack{t \in T, \\ a \in A \setminus \{\emptyset\}}} x_{gta} \quad \forall g \in G. \tag{2}$$

$$\sum_{\substack{a \in A \\ t + |M| - 1}} x_{gta} = 1 \quad \forall g \in G, t \in T. \tag{3}$$

$$\sum_{s=t} x_{gs*} \leq 1 \quad \forall g \in G, t = 1 \dots T_{max} - |M| + 1. \tag{4}$$

$$\sum_{t \in T} x_{gt*} \leq 2 \quad \forall g \in G. \quad (5)$$

$$\delta_{pgt} \leq x_{gk\omega_{pk}} \quad \forall g \in G, p \in P, t = 1 \dots T_{max} - |M| + 1, k = p \dots p + |M| - 1. \quad (6)$$

$$\gamma_{pg} \leq \sum_{t=1}^{T_{max}-|M|+1} \delta_{pgt} \quad \forall g \in G, p \in P. \quad (7)$$

$$\sum_{g \in G} \gamma_{pg} \geq 1 \quad \forall p \in P \setminus P_{\text{all}}. \quad (8)$$

$$\gamma_{pg} = 1 \quad \forall p \in P_{\text{all}}, g \in G. \quad (9)$$

$$x_{gt\emptyset} \leq x_{g,t+1,\emptyset} \quad \forall g \in G, t = 1 \dots T_{max} - 1. \quad (10)$$

$$x_{gtm} + x_{g,t+1,m} \leq 1 \quad \forall g \in G, m \in M, t = 1 \dots T_{max} - 1. \quad (11)$$

$$x_{gta} \in \{0, 1\} \quad \forall g \in G, t \in T, a \in A. \quad (12)$$

$$\delta_{pgt} \in \{0, 1\} \quad \forall p \in P, g \in G, t = 1 \dots T_{max} - |M| + 1. \quad (13)$$

$$\gamma_{pg} \in \{0, 1\} \quad \forall p \in P, g \in G. \quad (14)$$

$$\ell \in \mathbb{N}. \quad (15)$$