

Modular Forms Modulo 2:


Governing Fields for the Hecke Algebra

Paul Dubois

University College of London

March 25, 2020

Modular Forms



Complex vector space of modular forms of weight n : M_n

For $f \in M_n$: $f(z) = \sum_{n=0}^{\infty} c(n)q^n$ with $q^n = e^{2\pi inz}$

Reduction Modulo 2

$$\left\{ f \in M_n \mid f = \sum_{n \in \mathbb{N}} c(n) q^n \text{ s.t. } c(n) \in \mathbb{N} \right\}$$

\parallel

$$\mathbb{F}_2[\Delta, E_2, E_3]$$

$$\overline{\Delta} \rightsquigarrow \Delta$$

$$\overline{E_2} \rightsquigarrow 1$$

$$\overline{E_3} \rightsquigarrow 1$$

$$\overline{M_n} \rightsquigarrow \mathbb{F}_2[\Delta]$$

Modular Forms Modulo 2

$$\mathcal{F} = \langle \Delta^k \mid k \text{ odd} \rangle_{\mathbb{F}_2} = \langle \Delta, \Delta^3, \Delta^5, \dots \rangle_{\mathbb{F}_2}$$

$$\begin{aligned}\Delta(q) &= q \prod_{n=1}^{\infty} (1 - q^n)^{24} \\ &= \sum_{n=0}^{\infty} \tau(n) q^n \\ &\equiv \sum_{m=0}^{\infty} q^{(2m+1)^2} \pmod{2}\end{aligned}$$

Hecke Operators

With

$$f(q) = \sum_{n \in \mathbb{N}} c(n) q^n$$

We define

$$\overline{T_m} | f(q) = \sum_{n \in \mathbb{N}} \gamma(n) q^n$$

Where

$$\gamma(n) = \sum_{a|(n,m), a \geq 1} a^{2k-1} c\left(\frac{mn}{a^2}\right)$$

Hecke Operators Modulo 2

With

$$f(q) = \sum_{n \in \mathbb{N}} c(n)q^n$$

We define

$$\overline{T_p} | f(q) = \sum_{n \in \mathbb{N}} \gamma(n)q^n$$

Where

$$\gamma(n) = \begin{cases} c(np) & \text{if } p \nmid n \\ c(np) + c(n/p) & \text{if } p \mid n \end{cases} \quad \text{and } p \text{ an odd prime.}$$

Examples

	Δ^1	Δ^3	Δ^5	Δ^7	Δ^9	Δ^{11}	Δ^{13}
T_3	0	Δ	0	Δ^5	Δ^3	Δ^9	Δ^7
T_5	0	0	Δ	Δ^3	0	0	Δ^9
T_7	0	0	0	Δ	0	0	Δ^3
T_{11}	0	Δ	0	Δ^5	Δ^3	$\Delta + \Delta^9$	Δ^7
T_{13}	0	0	Δ	Δ^3	0	0	$\Delta + \Delta^9$
T_{17}	0	0	0	0	Δ	Δ^3	Δ^5
T_{19}	0	Δ	0	Δ^5	Δ^3	$\Delta + \Delta^9$	Δ^7

$$\begin{array}{ccc}
 f = \Delta^k & & T_p|f = \Delta^m + \dots \\
 \updownarrow & & \updownarrow \\
 f = q^k + \dots & \xrightarrow{T_p} & T_p|f = q^m + \dots
 \end{array}$$

The Hecke Algebra

$$\begin{aligned} A &= \mathbb{F}_2 [T_3, T_5, T_7, T_{11}, T_{13}, \dots] \\ &= \mathbb{F}_2 [T_p \mid p \in \mathbb{P}] \end{aligned}$$

$$A = \mathbb{F}_2 [[T_3, T_5]]$$

$$T_p = \sum_{i+j \geq 1} a_{ij}(p) T_3^i T_5^j$$

Examples

$$x = T_3 \quad y = T_5$$

$$\text{i.e. } x^a y^b = T_3^a T_5^b$$

$$T_3 = x^1 y^0 = x$$

$$T_5 = x^0 y^1 = y$$

$$T_7 = x^1 y^1 + x^3 y^1 + x^3 y^3 + x^5 y^1 + x^1 y^7 + x^1 y^9 + x^7 y^3 + x^7 y^5 + x^9 y^3 + x^{11} y^1 + x^3 y^{11} + x^5 y^9 + x^{13} y^1 + x^3 y^{13} + x^5 y^{11} + x^9 y^7 + x^{11} y^5 + x^{13} y^3 + x^3 y^{15} + x^7 y^{11} + x^9 y^9 + x^{13} y^5 + x^{15} y^3 + \dots$$

$$T_{11} = x^1 y^0 + x^1 y^2 + x^3 y^0 + x^1 y^4 + x^3 y^2 + x^5 y^0 + x^1 y^6 + x^3 y^4 + x^7 y^2 + x^1 y^{10} + x^3 y^8 + x^7 y^4 + x^9 y^2 + x^{11} y^2 + x^3 y^{12} + x^5 y^{10} + x^7 y^8 + x^{11} y^4 + x^{13} y^2 + x^9 y^8 + x^{17} y^0 + \dots$$

Dirichlet Density Theorem

Intuition: *Animation*

(see <https://pauldubois98.github.io/DirichletDensityTheoremAnimation/>)

Theorem (Dirichlet's Density Theorem)

Let $n \in \mathbb{N}^$, $a \in \mathbb{N}$ such that $\gcd(a, n) = 1$.*

If $S = \{p \in \mathbb{P} \mid p \equiv a \pmod{n}\}$, then S has density $1/\varphi(n)$.

Chebotarev Density Theorem

Theorem (Chebotarev Density Theorem)

With L/K an extension of Galois group $G = \text{Gal}(L/K)$.

Let C be a conjugacy class in G .

Then, the proportion of unramified primes ideals \mathfrak{p} in K that have Frobenius element $\text{Frob}_{L/K}(\mathfrak{p}) = C$ is $|C|/|G|$.

$L = \mathbb{Q}(\zeta_n)$ and $K = \mathbb{Q}$ give Dirichlet's density theorem.

Frobenian Maps

With K a number field, P the set of primes in K . $f : P \rightarrow \Omega$ is Frobenian if there exists

- ▶ a set $S \subset P$
- ▶ a field M extending K
- ▶ a class function $\phi : \text{Gal}(M/K) \rightarrow \Omega$

such that:

$$f(\mathfrak{p}) = \phi(\text{Frob}_{M/K}(\mathfrak{p})) \quad \forall \mathfrak{p} \in P \setminus S$$

The maps a_{ij}

$$T_p = \sum_{i+j \geq 1} a_{ij}(p) T_3^i T_5^j$$

$a_{ij} : p \mapsto a_{ij}(p)$ is Frobenian

M_{ij} denotes a governing field for a_{ij}

G_{ij} denotes a governing group for a_{ij}

Known identities

- ▶ $a_{10}(p) = 1 \iff p \equiv 3 \pmod{8}$
- ▶ $a_{01}(p) = 1 \iff p \equiv 5 \pmod{8}$
- ▶ $a_{11}(p) = 1 \iff p \equiv 7 \pmod{8}$
- ▶ $a_{20}(p) = 1 \iff \exists a, b \in \mathbb{Z} \text{ and } b \text{ odd, such that}$
 $p = a^2 + 8b^2, p \equiv 3 \pmod{8}$
- ▶ $a_{02}(p) = 1 \iff \exists a, b \in \mathbb{Z} \text{ and } b \text{ odd, such that}$
 $p = a^2 + 16b^2, p \equiv 3 \pmod{8}$

Known Governing Fields

$$M_{01} = \mathbb{Q}(\zeta_8)$$

$$M_{02} = \mathbb{Q}(\zeta_8, \sqrt[4]{2})$$

$$M_{11} = \mathbb{Q}(\zeta_8, \sqrt{\zeta_8}) = \mathbb{Q}(\zeta_{16})$$

$$M_{02} = \mathbb{Q}(\zeta_8, \sqrt{1+i})$$

$$M_{01} = \mathbb{Q}(\zeta_8)$$

$$\text{Note } \mathbb{Q}(\zeta_8) = \mathbb{Q}(i, \sqrt{2})$$

Potential New Governing Fields

$$M_{03} \stackrel{?}{=} \mathbb{Q} \left(\zeta_8, \sqrt[4]{2}, \sqrt{\alpha} \right)$$

where:

$$\alpha = -\frac{3136435454775881\sqrt[4]{2}}{562949953421312} + \frac{4208721080340285\sqrt{2}}{2251799813685248} +$$
$$\frac{3672578267558083 \cdot \sqrt[4]{2}^3}{562949953421312} + \frac{3582104167901087}{281474976710656}$$


Potential New Governing Fields

$$M_{05} \stackrel{?}{=} \mathbb{Q} \left(\zeta_8, \sqrt[4]{2}, \sqrt{\alpha}, \sqrt{\beta} \right)$$

where α remains and

$$\begin{aligned} \beta = & - \frac{8282936156772053\alpha^{\frac{13}{2}}}{1125899906842624} - \frac{1240182980093567\alpha^6}{562949953421312} \\ & - \frac{336382584949535\alpha^{\frac{9}{2}}}{2199023255552} - \frac{6445823996745319\alpha^4}{140737488355328} \\ & - \frac{4638634719581101\alpha^{\frac{5}{2}}}{35184372088832} - \frac{2954723016803317\alpha^2}{70368744177664} \\ & - \frac{5142889464378747\sqrt[4]{2}}{140737488355328} - \frac{4198844765367981\sqrt{\alpha}}{1125899906842624} \\ & + \dots \end{aligned}$$

Governing Fields Extension Graph




Governing Groups

$$\begin{array}{llll} G_{01} & \cong & D_4 & \cong C_2 \times C_2 \\ G_{02} & \cong & D_8 & \\ G_{03} & \cong & D_{16} & \\ G_{04} & \cong & D_{16} & \\ G_{05} & \cong & D_{32} & \\ G_{06} & \cong & D_{32} & \\ G_{07} & \cong & D_{32} & \end{array}$$

$$\begin{array}{llll} G_{10} & \cong & D_4 & \cong C_2 \times C_2 \\ G_{20} & \cong & D_8 & \\ G_{30} & \cong & D_{16} & \\ G_{40} & \cong & D_{16} & \\ G_{50} & \cong & D_{32} & \\ G_{60} & \cong & D_{32} & \\ G_{70} & \cong & D_{32} & \end{array}$$

Governing Groups Extension Graph



Diagonal Governing Groups

Conjecture (Diagonal Governing Groups Conjecture)

For all $k \in \mathbb{N}^$, there exists a field M_{0k} such that M_{0k} is a governing field for a_{0k} , and $G_{0k} = \text{Gal}(M_{0k}/\mathbb{Q})$ is dihedral.
For all $k \in \mathbb{N}^*$, there exists a field M_{k0} such that M_{k0} is a governing field for a_{k0} , and G_{k0} is dihedral.*

Moreover $M_{k0} \neq M_{0k}$ in general, but $G_{k0} \cong G_{0k}$.

Thank you