


# Mid-PhD Defense

Paul Dubois

TheraPanacea  
MICS, CentraleSupélec  
Institut du Cancer de Montpellier

21st June 2023

# Automatic Dose Optimization for Radiotherapy



# Problem Formulation

## IMRT

Bixel values:

$$x_{i,j}^{\theta} \geq 0, \text{ for } \theta \in \Theta \text{ and } 1 \leq i, j \leq 20^1$$

usually concatenated to a single bixels-value vector  $x$ .

Dose calculation:

$$\mathbf{y} = L\mathbf{x} \text{ with } L \text{ (pre-calculated) dose-influence (DI) matrix}$$

---

<sup>1</sup>20x20 is a typical bixel discretization

# Problem Formulation

IMRT (bis)

Objective for *maximum* constraint  $c$  on structure  $s$ , dose  $d$ :

$$f_c(\mathbf{y}) = \frac{1}{|\mathcal{V}|} \sum_{v \in \mathcal{V}} (\mathbf{y}_v - d)_+^2$$

(reverse sign for minimal constraint).


Final objective:

$$f(\mathbf{y}) = \sum_{c \in \mathcal{C}} w_c f_c(\mathbf{y})$$

with  $w_c$  the weight of constraint  $c$ .

# Problem Optimization

## Optimizer review



# Problem Optimization

## Optimizer review (bis)

