

# Exercises Set 5

Paul Dubois

September 25, 2023

## Abstract

Only the questions with a \* are compulsory (but do all of them!).

## 1 Change of Basis

Let  $\mathcal{B} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  be the standard canonical basis for  $\mathbb{R}^3$ .

Suppose we have another basis  $\mathcal{B}' = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$  for  $\mathbb{R}^3$  and let  $Q$  be the matrix whose columns are the coordinates of

$$\mathbf{u}_1 = \begin{pmatrix} 0.5 \\ -1 \\ 1 \end{pmatrix}, \mathbf{u}_2 = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}, \text{ and } \mathbf{u}_3 = \begin{pmatrix} -0.25 \\ 0.5 \\ 0 \end{pmatrix}$$

with respect to the standard basis. That is,  $Q = [\mathbf{u}_1 \quad \mathbf{u}_2 \quad \mathbf{u}_3]$ .

Let  $\mathbf{v} = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}$  (in the standard basis). Express  $\mathbf{v}$  in the basis  $\mathcal{B}'$ .