

Exercises Set 5

Paul Dubois

September 25, 2023

Abstract

Only the questions with a * are compulsory (but do all of them!).

1 Change of Basis

Let $\mathcal{B} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ be the standard canonical basis for \mathbb{R}^3 .

Suppose we have another basis $\mathcal{B}' = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ for \mathbb{R}^3 and let Q be the matrix whose columns are the coordinates of

$$\mathbf{u}_1 = \begin{pmatrix} 0.5 \\ -1 \\ 1 \end{pmatrix}_{\mathcal{B}}, \mathbf{u}_2 = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}_{\mathcal{B}}, \text{ and } \mathbf{u}_3 = \begin{pmatrix} -0.25 \\ 0.5 \\ 0 \end{pmatrix}_{\mathcal{B}}$$

with respect to the standard basis. That is, $Q = [\mathbf{u}_1 \ \mathbf{u}_2 \ \mathbf{u}_3]$.

Let $\mathbf{v} = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}_{\mathcal{B}'}$. Express \mathbf{v} in the standard basis \mathcal{B} .

Let $\mathbf{w} = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}_{\mathcal{B}}$. Express \mathbf{w} in the basis \mathcal{B}' .

2 Variance