

NotebookOptimization_Solutions

August 28, 2025

```
[1]: import numpy as np
import matplotlib.pyplot as plt
import plotly.graph_objects as go
```

1 Finding minimum value of a function


1.1 One dimensional function

$$f(x) = x^2 - x + 3$$

```
[2]: def f(x):
    return x**2 - x + 3
```

Simple plot of the function. Note that if the function had many (> 2 parameters, then plotting would not be possible).

```
[3]: xs = np.linspace(-4, 5, 100)
ys = f(xs)
plt.plot(xs, ys)
plt.axis('off')
plt.axhline(0, color='black', lw=0.5)
plt.axvline(0, color='black', lw=0.5)
plt.show()
```



1.1.1 Method 1: Grid search


```
[4]: x_vals = [-1.75, -1.25, -0.75, -0.25, 0.25, 0.75, 1.25, 1.75, 2.25, 2.75, 3.25]
     for x in x_vals:
         print(f'f({x}) = {f(x)}')
```

```
f(-1.75) = 7.8125
f(-1.25) = 5.8125
f(-0.75) = 4.3125
f(-0.25) = 3.3125
f(0.25) = 2.8125
f(0.75) = 2.8125
f(1.25) = 3.3125
f(1.75) = 4.3125
f(2.25) = 5.8125
f(2.75) = 7.8125
f(3.25) = 10.3125
```

More visual version:

```
[5]: x_vals = [-1.75, -1.25, -0.75, -0.25, 0.25, 0.75, 1.25, 1.75, 2.25, 2.75, 3.25]
     for x in x_vals:
         plt.scatter(x, f(x), color='red')
         plt.plot([x, x], [f(x), 0], color='black', lw=0.5, linestyle='--')
```

```
plt.plot(xs, ys)
plt.axis('off')
plt.axhline(0, color='black', lw=0.5)
plt.axvline(0, color='black', lw=0.5)
plt.show()
```



Minimum value of $f(x)$ found is 2.8125 at $x = 0.25$ and $x = 0.75$.

1.1.2 Method 2: Dichotomy

```
[6]: a = -5
b = 5
for i in range(6):
    print(f"a: f({a}) = {f(a)} \t b: f({b}) = {f(b)}", end='\t\t')
    if f(a)>f(b):
        a = (a+b)/2
        print(f'set a = {a}')
    else:
        b = (a+b)/2
        print(f'set b = {b}')

x = ((a+b)/2)
print(f"best guess: x = {x}")
```

```

a: f(-5) = 33      b: f(5) = 23          set a = 0.0
a: f(0.0) = 3.0    b: f(5) = 23          set b = 2.5
a: f(0.0) = 3.0    b: f(2.5) = 6.75      set b = 1.25
a: f(0.0) = 3.0    b: f(1.25) = 3.3125   set b = 0.625
a: f(0.0) = 3.0    b: f(0.625) = 2.765625 set a = 0.3125
a: f(0.3125) = 2.78515625 b: f(0.625) = 2.765625 set a = 0.46875
best guess: x = 0.546875


```

More visual version

```

[7]: a = -5
      b = 5
      plt.plot([a,b], [1, 1], color='green')
      plt.scatter([a,b], [1, 1], color='green')
      for i in range(6):
          if f(a)>f(b):
              a = (a+b)/2
          else:
              b = (a+b)/2
          plt.plot([a,b], [-i, -i], color='green')
          plt.scatter([a,b], [-i, -i], color='green')
      x = ((a+b)/2)
      plt.scatter([x], [-6], color='black')
      plt.plot(xs, ys)
      plt.axis('off')
      plt.axhline(0, color='black', lw=0.5)
      plt.axvline(0, color='black', lw=0.5)
      plt.show()

```



1.1.3 Method 3: Gradient descent

We first calculate the derivative $f'(x) = 2x - 1$

```
[8]: def df(x):
      return 2*x-1
```

```
[9]: lr = 0.8
x = -0.5
l = [x]
for _ in range(7):
    print(f"f({x:.4f}) = {f(x):.4f} \t f'({x:.4f}) = {df(x):.4f}")
    x = x - lr * df(x)
    l.append(x)
print(f'Final guess: x = {x:.4f}')

xs = np.linspace(min(l)-0.5, max(l+[2])+0.5, 100)
ys = f(xs)
```

$f(-0.5000) = 3.7500$	$f'(-0.5000) = -2.0000$
$f(1.1000) = 3.1100$	$f'(1.1000) = 1.2000$
$f(0.1400) = 2.8796$	$f'(0.1400) = -0.7200$
$f(0.7160) = 2.7967$	$f'(0.7160) = 0.4320$
$f(0.3704) = 2.7668$	$f'(0.3704) = -0.2592$

```

f(0.5778) = 2.7560      f'(0.5778) = 0.1555
f(0.4533) = 2.7522      f'(0.4533) = -0.0933
Final guess: x = 0.5280

```

```

[10]: lr = 0.3
x = -0.5
l = [x]
for _ in range(7):
    print(f"f({x:.4f}) = {f(x):.4f} \t f'({x:.4f}) = {df(x):.4f}")
    x = x - lr * df(x)
    l.append(x)
print(f'Final guess: x = {x:.4f}')

xs = np.linspace(min(l)-0.5, max(l+[2])+0.5, 100)
ys = f(xs)

```

```

f(-0.5000) = 3.7500      f'(-0.5000) = -2.0000
f(0.1000) = 2.9100      f'(0.1000) = -0.8000
f(0.3400) = 2.7756      f'(0.3400) = -0.3200
f(0.4360) = 2.7541      f'(0.4360) = -0.1280
f(0.4744) = 2.7507      f'(0.4744) = -0.0512
f(0.4898) = 2.7501      f'(0.4898) = -0.0205
f(0.4959) = 2.7500      f'(0.4959) = -0.0082
Final guess: x = 0.4984

```

Let's try other values for learning rate

```

[11]: for lr in [0.3, 0.01, 1, 2]:
    print(f'\t learning rate: {lr}')
    x = -1
    l = [x]
    for _ in range(7):
        print(f"f({x:.2f}) = {f(x):.2f} \t f'({x:.2f}) = {df(x):.2f}")
        x = x - lr * df(x)
        l.append(x)
    print(f'Final guess: x = {x:.2f} \n\n\n')

    xs = np.linspace(min(l)-0.5, max(l+[2])+0.5, 100)
    ys = f(xs)

```

```

\t learning rate: 0.3
f(-1.00) = 5.00      f'(-1.00) = -3.00
f(-0.10) = 3.11      f'(-0.10) = -1.20
f(0.26) = 2.81      f'(0.26) = -0.48
f(0.40) = 2.76      f'(0.40) = -0.19
f(0.46) = 2.75      f'(0.46) = -0.08
f(0.48) = 2.75      f'(0.48) = -0.03
f(0.49) = 2.75      f'(0.49) = -0.01
Final guess: x = 0.50

```

```

learning rate: 0.01
f(-1.00) = 5.00      f'(-1.00) = -3.00
f(-0.97) = 4.91      f'(-0.97) = -2.94
f(-0.94) = 4.83      f'(-0.94) = -2.88
f(-0.91) = 4.74      f'(-0.91) = -2.82
f(-0.88) = 4.66      f'(-0.88) = -2.77
f(-0.86) = 4.59      f'(-0.86) = -2.71
f(-0.83) = 4.52      f'(-0.83) = -2.66
Final guess: x = -0.80

```

```

learning rate: 1
f(-1.00) = 5.00      f'(-1.00) = -3.00
f(2.00) = 5.00       f'(2.00) = 3.00
f(-1.00) = 5.00      f'(-1.00) = -3.00
f(2.00) = 5.00       f'(2.00) = 3.00
f(-1.00) = 5.00      f'(-1.00) = -3.00
f(2.00) = 5.00       f'(2.00) = 3.00
f(-1.00) = 5.00      f'(-1.00) = -3.00
Final guess: x = 2.00

```

```

learning rate: 2
f(-1.00) = 5.00      f'(-1.00) = -3.00
f(5.00) = 23.00      f'(5.00) = 9.00
f(-13.00) = 185.00   f'(-13.00) = -27.00
f(41.00) = 1643.00   f'(41.00) = 81.00
f(-121.00) = 14765.00 f'(-121.00) = -243.00
f(365.00) = 132863.00 f'(365.00) = 729.00
f(-1093.00) = 1195745.00 f'(-1093.00) = -2187.00
Final guess: x = 3281.00

```

Visually:

```


[12]: for lr in [0.8, 0.3, 0.01, 1, 1.25]:
        x = -1
        l = [x]
        for _ in range(7):
            x = x - lr * df(x)
            l.append(x)



```



```

plt.plot(l, [f(x) for x in l], color='grey')
plt.scatter(l, [f(x) for x in l], color='black')
xs = np.linspace(min(l)-0.5, max(l+[2])+0.5, 100)
ys = f(xs)
plt.plot(xs, ys)
plt.axis('off')
plt.axhline(0, color='black', lw=0.5)
plt.axvline(0, color='black', lw=0.5)
plt.show()

```







1.2 Two dimensional function


$$f(x, y) = (x + y - 1)^2 + \frac{1}{5}(x - y)^2$$

```
[13]: def f(x, y):  
       return (x+y-1)**2 + 0.2*(x-y)**2
```

Plotting the function as a 2D surface. Plotting log of the function helps to see the minimum more clearly.

```
[14]: # interactive  
x = np.linspace(-1, 2, 100)  
y = np.linspace(-1, 2, 100)  
X, Y = np.meshgrid(x, y)  
Z = f(X, Y)  
Zlog = np.log(Z)  
go.Figure(data=[go.Surface(z=Z)])
```

```
[15]: # not interactive  
x = np.linspace(-1, 2, 100)  
y = np.linspace(-1, 2, 100)  
X, Y = np.meshgrid(x, y)  
Z = f(X, Y)  
Zlog = np.log(Z)  
  
# plot a heatmap of the function  
plt.figure(figsize=(10,5))  
plt.subplot(1,2,1)  
plt.imshow(Z, extent=(-1, 2, -1, 2))  
plt.colorbar()  
plt.title('$f(x,y)$')  
plt.xlabel('x')  
plt.ylabel('y')  
plt.subplot(1,2,2)  
plt.imshow(Zlog, extent=(-1, 2, -1, 2))  
plt.colorbar()  
plt.title('$\log(f(x,y))$')  
plt.xlabel('x')  
plt.ylabel('y')  
plt.show()
```



1.2.1 Method 1: Grid search

```
[16]: for x in np.arange(-1, 2.5, 0.5):
      for y in np.arange(-1, 2.5, 0.5):
          print(f'f({x}, {y}) = {f(x, y)}')
```

```
f(-1.0, -1.0) = 9.0
f(-1.0, -0.5) = 6.3
f(-1.0, 0.0) = 4.2
f(-1.0, 0.5) = 2.7
f(-1.0, 1.0) = 1.8
f(-1.0, 1.5) = 1.5
f(-1.0, 2.0) = 1.8
f(-0.5, -1.0) = 6.3
f(-0.5, -0.5) = 4.0
f(-0.5, 0.0) = 2.3
f(-0.5, 0.5) = 1.2
f(-0.5, 1.0) = 0.7
f(-0.5, 1.5) = 0.8
f(-0.5, 2.0) = 1.5
f(0.0, -1.0) = 4.2
f(0.0, -0.5) = 2.3
f(0.0, 0.0) = 1.0
f(0.0, 0.5) = 0.3
f(0.0, 1.0) = 0.2
f(0.0, 1.5) = 0.7
f(0.0, 2.0) = 1.8
f(0.5, -1.0) = 2.7
f(0.5, -0.5) = 1.2
f(0.5, 0.0) = 0.3
```

```

f(0.5, 0.5) = 0.0
f(0.5, 1.0) = 0.3
f(0.5, 1.5) = 1.2
f(0.5, 2.0) = 2.7
f(1.0, -1.0) = 1.8
f(1.0, -0.5) = 0.7
f(1.0, 0.0) = 0.2
f(1.0, 0.5) = 0.3
f(1.0, 1.0) = 1.0
f(1.0, 1.5) = 2.3
f(1.0, 2.0) = 4.2
f(1.5, -1.0) = 1.5
f(1.5, -0.5) = 0.8
f(1.5, 0.0) = 0.7
f(1.5, 0.5) = 1.2
f(1.5, 1.0) = 2.3
f(1.5, 1.5) = 4.0
f(1.5, 2.0) = 6.3
f(2.0, -1.0) = 1.8
f(2.0, -0.5) = 1.5
f(2.0, 0.0) = 1.8
f(2.0, 0.5) = 2.7
f(2.0, 1.0) = 4.2
f(2.0, 1.5) = 6.3
f(2.0, 2.0) = 9.0

```

Note that it becomes computationally expensive to search for the minimum in a 2D grid. For higher dimensions, this method is not feasible.


More visually:

```

[17]: # plot a heatmap of the function
plt.figure(figsize=(10,5))
plt.subplot(1,2,1)
plt.imshow(Z, extent=(-1, 2, -1, 2))
for x in np.arange(-1, 2.5, 0.5):
    for y in np.arange(-1, 2.5, 0.5):
        plt.scatter(x, y, color='red')
plt.colorbar()
plt.title('$f(x,y)$')
plt.xlabel('x')
plt.ylabel('y')
plt.subplot(1,2,2)
plt.imshow(Zlog, extent=(-1, 2, -1, 2))
for x in np.arange(-1, 2.5, 0.5):
    for y in np.arange(-1, 2.5, 0.5):
        plt.scatter(x, y, color='red')
plt.colorbar()

```

```
plt.title('$\log(f(x,y))$')
plt.xlabel('x')
plt.ylabel('y')
plt.show()
```



1.2.2 Method 2: Dichotomy

```
[18]: a, b = -10, 10
      c, d = -10, 10
      for _ in range(10):
          f1 = f(a, c)
          f2 = f(a, d)
          f3 = f(b, c)
          f4 = f(b, d)
          if f1 <= f2 and f1 <= f3 and f1 <= f4:
              b = (a+b)/2
              d = (c+d)/2
          if f2 <= f1 and f2 <= f3 and f2 <= f4:
              b = (a+b)/2
              c = (c+d)/2
          if f3 <= f2 and f3 <= f1 and f3 <= f4:
              a = (a+b)/2
              d = (c+d)/2
          if f4 <= f2 and f4 <= f3 and f4 <= f1:
              a = (a+b)/2
              c = (c+d)/2
          print(f'f({(a+b)/2:.2f}, {(c+d)/2:.2f}) = {f((a+b)/2, (c+d)/2):.4f} \t {a:.2f} < x < {b:.2f} \t {c:.2f} < y < {d:.2f} \t ')

```

```
print(f'Best guess: x = {(a+b)/2:.2f}, y = {(c+d)/2:.2f}')
```

f(-2.50, 2.50) = 6.0000	-5.00 < x < 0.00	0.00 < y < 5.00
f(-1.25, 1.25) = 2.2500	-2.50 < x < 0.00	0.00 < y < 2.50
f(-0.62, 0.62) = 1.3125	-1.25 < x < 0.00	0.00 < y < 1.25
f(-0.31, 0.94) = 0.4531	-0.62 < x < 0.00	0.62 < y < 1.25
f(-0.16, 0.78) = 0.3164	-0.31 < x < 0.00	0.62 < y < 0.94
f(-0.08, 0.86) = 0.2236	-0.16 < x < 0.00	0.78 < y < 0.94
f(-0.04, 0.82) = 0.1956	-0.08 < x < 0.00	0.78 < y < 0.86
f(-0.02, 0.84) = 0.1800	-0.04 < x < 0.00	0.82 < y < 0.86
f(-0.01, 0.83) = 0.1734	-0.02 < x < 0.00	0.82 < y < 0.84
f(-0.00, 0.83) = 0.1699	-0.01 < x < 0.00	0.83 < y < 0.84

Best guess: x = -0.00, y = 0.83

More visual version

```
[19]: a, b = -5, 5
      c, d = -5, 5


      fig = plt.figure(figsize=(15,5))
      x = np.linspace(a,b, 100)
      y = np.linspace(c,d, 100)
      X, Y = np.meshgrid(x, y)
      Z = f(X, Y)
      Zlog = np.log(Z)
      plt.subplot(1,3,1)
      plt.imshow(Z, extent=(a, b, c, d))
      plt.title('$f(x,y)$')
      plt.xlabel('x')
      plt.ylabel('y')
      plt.subplot(1,3,2)
      plt.imshow(Zlog, extent=(a, b, c, d))
      plt.title('$\log(f(x,y))$')
      plt.xlabel('x')
      plt.ylabel('y')

      ax = plt.subplot(1,3,3)
      rectangles = []
      for _ in range(10):
          rectangle = plt.Rectangle((a, c), b-a, d-c, facecolor='green', alpha=0.1)
          ax.add_patch(rectangle)
          f1 = f(a, c)
          f2 = f(a, d)
          f3 = f(b, c)
          f4 = f(b, d)
          if f1<=f2 and f1<=f3 and f1<=f4:
              b = (a+b)/2
```

```

    d = (c+d)/2
    if f2<=f1 and f2<=f3 and f2<=f4:
        b = (a+b)/2
        c = (c+d)/2
    if f3<=f2 and f3<=f1 and f3<=f4:
        a = (a+b)/2
        d = (c+d)/2
    if f4<=f2 and f4<=f3 and f4<=f1:
        a = (a+b)/2
        c = (c+d)/2
plt.plot((a+b)/2, (c+d)/2, '+', color='black')
plt.axis('equal')
plt.show()

```



1.2.3 Method 3: Gradient descent

We start by calculating the partial derivatives of the function with respect to x and y .

$$\frac{\partial f}{\partial x}(x,y) = 2(x+y-1) + \frac{2}{5}(x-y)$$

$$\frac{\partial f}{\partial y}(x,y) = 2(x+y-1) - \frac{2}{5}(x-y)$$

```

[20]: def dfx(x, y):
        return 2*(x+y-1) + 0.4*(x-y)
    def dfy(x, y):
        return 2*(x+y-1) - 0.4*(x-y)

```

```

[21]: lr = 0.6
    x = 2.5
    y = 3
    lx = [x]
    ly = [y]
    for _ in range(7):
        x = x - lr * dfx(x,y)

```



```

y = y - lr * dfy(x,y)
lx.append(x)
ly.append(y)
print(f'f(x,y) = {f(x,y):.2f} \t (x,y) = ({x:.2f}, {y:.2f})')

```

```

f(x,y) = 7.20      (x,y) = (-2.78, 2.55)
f(x,y) = 0.85      (x,y) = (-0.02, 0.10)
f(x,y) = 0.25      (x,y) = (1.11, 0.09)
f(x,y) = 0.04      (x,y) = (0.63, 0.56)
f(x,y) = 0.01      (x,y) = (0.39, 0.58)
f(x,y) = 0.00      (x,y) = (0.47, 0.49)
f(x,y) = 0.00      (x,y) = (0.52, 0.48)

```

```

[22]: lr = 0.6
for lr in [0.3, 0.1, 0.01, 1.25]:
    print(f'learning rate: {lr}')
    x = 2.5
    y = 3
    lx = [x]
    ly = [y]
    for _ in range(7):
        x = x - lr * dfx(x,y)
        y = y - lr * dfy(x,y)
        lx.append(x)
        ly.append(y)
        print(f'f(x,y) = {f(x,y):.2f} \t (x,y) = ({x:.2f}, {y:.2f})')
    print('\n\n\n')

```

```

learning rate: 0.3
f(x,y) = 0.68      (x,y) = (-0.14, 1.51)
f(x,y) = 0.32      (x,y) = (-0.16, 1.10)
f(x,y) = 0.16      (x,y) = (0.03, 0.90)
f(x,y) = 0.07      (x,y) = (0.18, 0.77)
f(x,y) = 0.03      (x,y) = (0.28, 0.68)
f(x,y) = 0.01      (x,y) = (0.35, 0.62)
f(x,y) = 0.01      (x,y) = (0.40, 0.58)

```

```

learning rate: 0.1
f(x,y) = 8.14      (x,y) = (1.62, 2.22)
f(x,y) = 3.29      (x,y) = (1.08, 1.72)
f(x,y) = 1.36      (x,y) = (0.74, 1.39)
f(x,y) = 0.58      (x,y) = (0.54, 1.17)
f(x,y) = 0.27      (x,y) = (0.43, 1.02)
f(x,y) = 0.14      (x,y) = (0.36, 0.92)
f(x,y) = 0.08      (x,y) = (0.33, 0.84)

```

```

learning rate: 0.01
f(x,y) = 18.72    (x,y) = (2.41, 2.91)
f(x,y) = 17.27    (x,y) = (2.33, 2.82)
f(x,y) = 15.93    (x,y) = (2.25, 2.74)
f(x,y) = 14.69    (x,y) = (2.17, 2.66)
f(x,y) = 13.55    (x,y) = (2.09, 2.58)
f(x,y) = 12.50    (x,y) = (2.02, 2.51)
f(x,y) = 11.53    (x,y) = (1.95, 2.44)

```

```

learning rate: 1.25
f(x,y) = 112.80      (x,y) = (-8.50, 13.50)
f(x,y) = 324.80      (x,y) = (-7.50, -9.50)
f(x,y) = 1804.80     (x,y) = (36.50, -51.50)
f(x,y) = 5196.80     (x,y) = (32.50, 40.50)
f(x,y) = 28876.80    (x,y) = (-143.50, 208.50)
f(x,y) = 83148.80    (x,y) = (-127.50, -159.50)
f(x,y) = 462028.80   (x,y) = (576.50, -831.50)

```

A more visual version of the gradient descent method

```

[23]: for lr in [0.6, 0.3, 0.1, 0.01, 1.25]:
    x = 2.5
    y = 3
    lx = [x]
    ly = [y]
    for _ in range(7):
        x = x - lr * dfx(x,y)
        y = y - lr * dfy(x,y)
        lx.append(x)
        ly.append(y)


    # plot a heatmap of the function with the line search
    x = np.linspace(-3, 4, 100)
    y = np.linspace(-3, 4, 100)
    X, Y = np.meshgrid(x, y)
    Z = f(X, Y)
    Zlog = np.log(Z)

```


```

plt.figure(figsize=(10,5))
plt.subplot(1,2,1)
plt.imshow(Z, extent=(-3, 4, -3, 4))
plt.plot(lx, ly, color='grey')
plt.scatter(lx, ly, color='black')
plt.colorbar()
plt.title('$f(x,y)$')
plt.xlabel('x')
plt.ylabel('y')
plt.subplot(1,2,2)
plt.imshow(Zlog, extent=(-3, 4, -3, 4))
plt.plot(lx, ly, color='grey')
plt.scatter(lx, ly, color='black')
plt.colorbar()
plt.title('$\log(f(x,y))$')
plt.xlabel('x')
plt.ylabel('y')
plt.suptitle(f'Learning rate: {lr}')
plt.show()


```





Learning rate: 0.3



Learning rate: 0.1





2 Linear regression

We want to find the line of best fit for the following points: - $P_1 = (0, 1)$ - $P_2 = (3, 8)$ - $P_4 = (5, 12)$
 - $P_5 = (8, 17)$

The line of best fit is given by the equation $y = ax + b$ where a is the slope and b is the y -intercept.
Let's define the loss function as the sum of squared errors:

$$\mathcal{L}(a, b) = \sum_{i=1}^n (y_i - (ax_i + b))^2$$

so

$$\mathcal{L}(a, b) = (1 - (0a + b))^2 + (8 - (3a + b))^2 + (12 - (5a + b))^2 + (17 - (8a + b))^2$$

We can find the minimum of the loss function by taking the partial derivatives with respect to a and b and setting them to zero.

With respect to a :

$$\frac{\partial \mathcal{L}}{\partial a}(a, b) = -2(1 - (0a + b))0 + -2(8 - (3a + b))3 + -2(12 - (5a + b))5 + -2(17 - (8a + b))8$$

$$\frac{\partial \mathcal{L}}{\partial a}(a, b) = -6(-3a - b + 8) + -10(-5a - b + 12) + -16(-8a - b + 17)$$

$$\frac{\partial \mathcal{L}}{\partial a}(a, b) = 18a + 6b - 48 + 50a + 10b - 120 + 128a + 16b - 272$$

$$\frac{\partial \mathcal{L}}{\partial a}(a, b) = 196a + 32b - 440$$

With respect to b :

$$\frac{\partial \mathcal{L}}{\partial b}(a, b) = -2(1 - (0a + b)) + -2(8 - (3a + b)) + -2(12 - (5a + b)) + -2(17 - (8a + b))$$

$$\frac{\partial \mathcal{L}}{\partial b}(a, b) = -2(1 - b) - 2(-3a - b + 8) - 2(-5a - b + 12) - 2(-8a - b + 17)$$

$$\frac{\partial \mathcal{L}}{\partial b}(a, b) = 2b - 2 + 6a + 2b - 16 + 10a + 2b - 24 + 16a + 2b - 34$$

$$\frac{\partial \mathcal{L}}{\partial b}(a, b) = 8b - 76 + 32a$$

Setting the partial derivatives to zero:

$$196a + 32b - 440 = 0 \quad \text{and} \quad 8b - 76 + 32a = 0$$

$$\text{So } b = 9.5 - 4a$$

$$\text{Hence, } 196a + 32(9.5 - 4a) - 440 = 0 \implies 196a + 304 - 128a - 440 = 0 \implies 68a = 136 \implies a = 2$$

$$\text{Finally, } b = 9.5 - 4(2) = 1.5$$

Hence, the line of best fit is $y = 2x + 1.5$