

# Math 797W Algebraic geometry. Homework 3

Paul Hacking

October 1, 2012

- (1) Let  $X$  be an irreducible topological space and  $U \subset X$  an open subset. Show that  $U$  is irreducible.
- (2) Let  $U = \mathbb{A}_{x,y}^2 \setminus \{(0,0)\} \subset X = \mathbb{A}_{x,y}^2$ . Show that the prevariety  $U$  is not isomorphic to an affine variety. [Hint: First show  $\mathcal{O}_U(U) := \mathcal{O}_X(U) = k[x,y]$ . So, writing  $i: U \rightarrow X$  for the inclusion, the  $k$ -algebra homomorphism  $i^*: \mathcal{O}_X(X) \rightarrow \mathcal{O}_U(U)$  is an isomorphism. Now it follows that  $U$  cannot be affine (why?)]
- (3) Let  $n \in \mathbb{Z}$ . Let  $X = X(n)$  be the prevariety defined as follows:  $X$  is the union of two open affine sets  $U_1 = \mathbb{A}_{x_1,y_1}^2$  and  $U_2 = \mathbb{A}_{x_2,y_2}^2$  glued via the isomorphism

$$U_1 \supset (x_1 \neq 0) \xrightarrow{\sim} (y_2 \neq 0) \subset U_2, (x_1, y_1) \mapsto (x_1^n y_1, x_1^{-1}).$$

- (a) Compute  $\mathcal{O}_X(X)$ .
- (b) Show that  $X(0) \simeq \mathbb{P}^1 \times \mathbb{A}^1$  and  $X(-1) \simeq \mathbb{P}^2 \setminus \{P\}$ , where  $P \in \mathbb{P}^2$  is a point.
- (c) In class we defined a morphism  $p: X \rightarrow \mathbb{P}^1$ , given by

$$U_1 \rightarrow V_0 = \mathbb{A}_{z_{01}}^1, \quad (x_1, y_1) \mapsto x_1$$

$$U_2 \rightarrow V_1 = \mathbb{A}_{z_{10}}^1, \quad (x_2, y_2) \mapsto y_2$$

where  $V_i = (Z_i \neq 0) \subset \mathbb{P}_{(Z_0:Z_1)}^1$  and  $z_{ij} = Z_j/Z_i$ . The morphism  $p$  is a locally trivial fibration with fiber  $\mathbb{A}^1$ . Describe this morphism geometrically for  $n = -1$  in terms of the identification  $X(-1) \simeq \mathbb{P}^2 \setminus \{P\}$ .

- (4) Let  $Y = Z(f) \subset \mathbb{A}_{x_1, \dots, x_n}^n$  where  $f \in k[x_1, \dots, x_n]$  is irreducible. We say the hypersurface  $Y$  is *smooth* at a point  $P \in Y$  if  $\frac{\partial f}{\partial x_i}(P) \neq 0$  for some  $i$ . [This definition relies on an algebraic version of the implicit function theorem. The partial derivatives are defined formally via  $k$ -linearity, the Leibniz rule, and  $\frac{\partial x_i}{\partial x_j} = \delta_{ij}$ .] Now let  $Y = Z(y^2 - f(x)) \subset \mathbb{A}_{x,y}^2$  where  $f(x) \in k[x]$  is a polynomial of degree  $m$ . (We assume  $f(x) \in k[x]$  is not a square, so that  $y^2 - f(x)$  is irreducible.)

- (a) Show that  $X$  is smooth at each point iff  $f(x)$  has no multiple roots.
- (b) Let  $\bar{Y} \subset \mathbb{P}^2$  denote the closure of  $Y$  in

$$\mathbb{P}_{(X:Y:Z)}^2 = (Z \neq 0) \cup (Z = 0) = \mathbb{A}_{x,y}^2 \cup \mathbb{P}_{(X:Y)}^1,$$

where  $x = X/Z$ ,  $y = Y/Z$ . Show that  $\bar{Y} \setminus Y$  is a single point  $P$  for  $m \geq 3$  and  $\bar{Y}$  is not smooth at  $P$  for  $m \geq 4$ .

- (c) Let  $n = -(m + \delta)/2 \in \mathbb{Z}$ , where  $\delta = 0$  or  $1$ ,  $m \equiv \delta \pmod{2}$ . Let  $Z$  be the isomorphic image of  $Y$  in  $X(n)$  under the map

$$\mathbb{A}_{x,y}^2 \xrightarrow{\sim} U_1 = \mathbb{A}_{x_1,y_1}^2 \subset X(n), \quad (x, y) \mapsto (x_1, y_1).$$

Let  $\bar{Z}$  denote the closure of  $Z$  in  $X(n)$ . Show that  $|\bar{Z} \setminus Z| = 2 - \delta$  and  $\bar{Z}$  is smooth at each point of  $\bar{Z} \setminus Z$ .

- (d) Assume  $k = \mathbb{C}$ . Show that  $\bar{Z}$  is compact for the Euclidean topology (although  $X(n)$  is not compact).

Remark: In particular, the hyperelliptic curve  $\bar{Z}$  with affine open set  $Y \simeq Z \subset \bar{Z}$  does *not* embed in  $\mathbb{P}^2$  as the closure of  $Y$  for  $m \geq 4$ . (In fact, a smooth projective plane curve  $W = Z(F) \subset \mathbb{P}^2$  is never hyperelliptic if  $d = \deg F > 3$ .)

- (5) Let  $X = Z(f) \subset \mathbb{A}_{x,y}^2$ , where  $f \in k[x, y]$  is irreducible. Let  $P \in X$  be a point and suppose that  $X$  is smooth at  $P$ . Suppose WLOG that  $P = (0, 0)$  and  $\frac{\partial f}{\partial y}(P) \neq 0$ .

- (a) Show that the maximal ideal  $m_{X,P} \subset \mathcal{O}_{X,P}$  is generated by  $x$ . [Hint: Clearly the maximal ideal is generated by  $x$  and  $y$ . Use the equation  $f$  to eliminate  $y$ .]

- (b) Show that  $\bigcap_{n \geq 1} m_{X,P}^n = (0)$ . [Hint: Suppose  $0 \neq g \in \mathcal{O}_{X,P}$  and  $g \in m_{X,P}^n$ . Thus  $g = x^n \cdot h$ ,  $h \in \mathcal{O}_{X,P}$ . Write  $g = \frac{a}{b}$ ,  $h = \frac{c}{d}$ , where  $a, b, c, d \in k[x, y]$  and  $b(P), d(P) \neq 0$ . There are  $\alpha, \beta \in k[x, y]$  and  $\gamma \in k[x]$  such that  $\alpha f + \beta a = \gamma$  (because  $f, a$  are coprime in  $k(x)[y]$ , cf. HW1 Q11). Write  $\gamma = x^l \cdot \delta$ ,  $\delta(0) \neq 0$ . Then  $(\beta a - \gamma)d \equiv (\beta b c x^n - \delta d x^l) \equiv 0 \pmod{f}$ . Now show by contradiction that  $n \leq l$ .]
- (c) Using (a) and (b), deduce that every nonzero element  $0 \neq g \in k(X)$  can be written uniquely in the form  $g = x^n \cdot u$ , where  $n \in \mathbb{Z}$  and  $u \in \mathcal{O}_{X,P}$ ,  $u(P) \neq 0$ . (In particular,  $g \in \mathcal{O}_{X,P}$  if  $n \geq 0$  and  $g^{-1} \in \mathcal{O}_{X,P}$  if  $n \leq 0$ .)
- (d) Let  $0 \neq g_0, \dots, g_n \in k(X)$ . Show that the assignment

$$Q \mapsto (g_0(Q) : g_1(Q) : \dots : g_n(Q))$$

extends to a morphism

$$g : U \rightarrow \mathbb{P}^n$$

defined on some open neighborhood  $U$  of  $P \in X$ . [Hint: Use (c) and “clear denominators”.]

- (e) Give an example of a nonzero element  $0 \neq g \in k(\mathbb{A}^2)$  such that  $g \notin \mathcal{O}_{\mathbb{A}^2,0}$  and  $g^{-1} \notin \mathcal{O}_{\mathbb{A}^2,0}$ .
- (6) For  $n = 1$  there is a morphism  $\pi : X(1) \rightarrow \mathbb{A}^2$  given by

$$U_1 \rightarrow \mathbb{A}^2, \quad (x_1, y_1) \mapsto (x_1 y_1, y_1)$$

$$U_2 \rightarrow \mathbb{A}^2, \quad (x_2, y_2) \mapsto (x_2, x_2 y_2).$$

The morphism  $\pi$  is called the *blowup* of the point  $0 \in \mathbb{A}^2$ . The closed set  $C = \pi^{-1}(0) \subset X$  is irreducible and isomorphic to  $\mathbb{P}^1$ , and  $\pi$  restricts to an isomorphism  $X \setminus C \xrightarrow{\sim} \mathbb{A}^2 \setminus \{0\}$ .

Now suppose  $n \geq 1$ . Let  $\pi : X(n) \rightarrow \mathbb{A}^{n+1}$  be the morphism defined by

$$U_1 \rightarrow \mathbb{A}^{n+1}, \quad (x_1, y_1) \mapsto (x_1^n y_1, x_1^{n-1} y_1, \dots, y_1)$$

$$U_2 \rightarrow \mathbb{A}^{n+1}, \quad (x_2, y_2) \mapsto (x_2, x_2 y_2, \dots, x_2 y_2^n).$$

- (a) Check that  $\pi$  is a well defined morphism.

- (b) Show that  $C := \pi^{-1}(0)$  is irreducible and isomorphic to  $\mathbb{P}^1$ .
- (c) Let  $Y = \pi(X) \subset \mathbb{A}_{z_0, \dots, z_n}^{n+1}$ . Show that  $Y$  is the affine variety defined by the prime ideal  $J \subset k[z_0, \dots, z_n]$  generated by the  $2 \times 2$  minors of the matrix

$$\begin{pmatrix} z_0 & z_1 & \cdots & z_{n-1} \\ z_1 & z_2 & \cdots & z_n \end{pmatrix}$$

In particular, in the case  $n = 2$ ,  $Y$  is the hypersurface

$$Z(z_0 z_2 - z_1^2) \subset \mathbb{A}_{z_0, z_1, z_2}^3$$

studied in HW1 Q9. [Hint: Compare HW2 Q8.  $Y \subset \mathbb{A}^{n+1}$  is the affine cone over the rational normal curve of degree  $n$  in  $\mathbb{P}^n$ .]

- (d) Show that  $X \setminus C \rightarrow Y \setminus \{0\}$  is an isomorphism.
- (e) Show that  $\pi^*: k[Y] = \mathcal{O}_Y(Y) \rightarrow \mathcal{O}_X(X)$  is an isomorphism. [Hint:  $\overline{\pi(X)} = Y$  implies  $\pi^*$  is injective (why?). Now use your explicit computation of  $\mathcal{O}_X(X)$  from Q3(a) to show  $\pi^*$  is surjective.]

Remark: Note that  $0 \in Y$  is a singular point for  $n > 1$  (singular means not smooth). The morphism  $\pi: X \rightarrow Y$  is a *resolution* of the singularity  $0 \in Y$ .