

697B Example Sheet 5

Paul Hacking

14 October 2010

- (1) Show that the discriminant of $ay^2 + by + c$ is $a(4ac - b^2)$ and the discriminant of $y^3 + ay + b$ is $4a^3 + 27b^2$.
- (2) (Weak Bézout theorem) Let $f(x, y)$ and $g(x, y)$ be polynomials of degree n and m with no common factors. Let $X = (f(x, y) = 0) \subset \mathbb{C}_{x,y}^2$ and $Y = (g(x, y) = 0) \subset \mathbb{C}_{x,y}^2$. In this question we will use the resultant to show that $|X \cap Y| \leq mn$.
 - (a) Show that $X \cap Y$ is finite.
 - (b) Show that after a linear change of coordinates x, y we may assume that no two points in $X \cap Y$ have the same x coordinate and that $f(x, y) = y^n + a_1y^{n-1} \cdots + a_n$, $g(x, y) = y^m + b_1y^{m-1} + \cdots + b_m$, where the a_i, b_i are polynomials in x of degree at most i .
 - (c) With coordinates as in (b), show that the resultant $r(x) := R(f, g)$ is a nonzero polynomial of degree at most mn , and deduce that $|X \cap Y| \leq mn$.
- (3) Let $f(x, y) = y^3 - xy + 2$ and $X = (f(x, y) = 0) \subset \mathbb{C}_{x,y}^2$.
 - (a) Let $F: X \rightarrow Y$ be a non constant holomorphic map between Riemann surfaces X and Y , and $P \in X$ a point. Let z be a local coordinate at P and w a local coordinate at $F(P)$ and let $z \mapsto f(z)$ be the map F in the charts given by z, w . Then the *ramification index* of F at P is $e_P := \nu_P(f(z))$, and we say P is a ramification point if $e_P > 1$. (Note: After an analytic change of the local coordinate z , we may assume that F is given by $z \mapsto z^{e_P}$.) Find the ramification points and ramification indices of the map $p: X \rightarrow \mathbb{C}_x^1$ given by $(x, y) \mapsto x$. [Hint: The locus of ramification points is given by $(f = \frac{\partial f}{\partial y} = 0) \subset \mathbb{C}_{x,y}^2$ (why?).]

- (b) Let $\mathbb{C}^2 \subset \mathbb{P}^2$ be the inclusion given by $(x, y) \mapsto (x : y : 1)$ and let $\overline{X} \subset \mathbb{P}^2$ be the closure of X . Find the singular points of \overline{X} and describe their type.
 - (c) Describe the normalization $n: \tilde{X} \rightarrow \overline{X} \subset \mathbb{P}_{\mathbb{C}}^2$ explicitly. (Hint: Solve $f(x, y) = 0$ for x .)
 - (d) Show that the map $p: X \rightarrow \mathbb{C}^1$ extends to a holomorphic map $\tilde{p}: \tilde{X} \rightarrow \mathbb{C} \cup \{\infty\}$. Describe the ramification of \tilde{p} over ∞ .
- (4) (Degree) Let $F: X \rightarrow Y$ be a non constant holomorphic map between compact Riemann surfaces X and Y . Recall the definition of ramification points and ramification indices from Q3(a) above.
- (a) Show that F is surjective.
 - (b) Show that the number of ramification points of F is finite. (Hint: Use compactness of X and argue by contradiction.)
 - (c) Show that the function

$$f: Y \rightarrow \mathbb{Z}, \quad Q \mapsto \sum_{P \in F^{-1}(Q)} e_P.$$

is constant, say equal to d . Thus every fiber $F^{-1}(Q)$ of F consists of d points (provided we count with multiplicities), and all but finitely many fibers consist of d distinct points. The number d is called the *degree* of F . (Hint: Show that f is locally constant and use connectedness of Y .)

- (d) Let X be a compact Riemann surface and f a meromorphic function on X . In class we showed that $\sum \nu_P(f) = 0$. Use (c) to give a different proof of this fact. (Hint: Consider the holomorphic map $F: X \rightarrow \mathbb{P}^1$ defined by f .)
- (5) (Riemann–Hurwitz formula) Let $F: X \rightarrow Y$ be a non constant holomorphic map between compact Riemann surfaces X and Y . In this question we will prove the Riemann–Hurwitz formula

$$2g(X) - 2 = d(2g(Y) - 2) + \sum_{P \in X} (e_P - 1)$$

where $g(X)$ and $g(Y)$ are the genus of X and Y , and d is the degree of F (see Q4).

- (a) Let ω be a nonzero meromorphic differential on Y . (Actually, the existence of ω is not obvious, but we will assume it here.) We can define the *pullback* $F^*\omega$ of ω , a meromorphic differential on X , as follows: if $P \in X$ is a point, let z be a local coordinate at P and w a local coordinate at $F(P)$, and let $z \mapsto f(z)$ be the map F in the charts given by z, w . Write $\omega = g(w)dw$ in the chart given by w , then $F^*(\omega) = g(f(z))f'(z)dz$ in the chart given by z . Show that $F^*(\omega)$ is well defined.
 - (b) Show that $\nu_P(F^*(\omega)) = e_P \cdot \nu_{F(P)}(\omega) + (e_P - 1)$.
 - (c) Deduce the Riemann–Hurwitz formula (Hint: Use Poincaré–Hopf and Q4.)
 - (d) Use Riemann–Hurwitz to show that $g(X) \geq g(Y)$. Show that if $g(X) = g(Y)$ then either F is an isomorphism ($d = 1$), or $d > 1$ and $g(X) = 0$ or $g(X) = 1$. Give examples to show that these cases really do occur.
 - (e) Compute the genus of the curve \tilde{X} from HW2Q4 and HW3Q8 using Riemann–Hurwitz (this method is faster than the one described in HW3Q8).
 - (f) Check your answer to Q3 using Riemann–Hurwitz.
- (6) Suppose $F: \mathbb{C} \cup \{\infty\} \rightarrow \mathbb{C} \cup \{\infty\}$ is given by $z \mapsto f(z) = p(z)/q(z)$, where p, q are polynomials with no common factors. Then the degree d of F is the maximum of the degrees of p and q by HW1Q2(c). Show that the set of ramification points of F is given by $(p'q - q'p = 0) \subset \mathbb{C}$ together with ∞ if $p'q - q'p$ has degree $< 2d - 2$. What is the maximum possible number of ramification points of F ?