

SIMULATION OF DIFFUSION IN HETEROGENEOUS MEDIA

by

Paul Ionele

A dissertation submitted to the University of Ontario Institute
of Technology in accordance with the requirements of the
degree of Bachelor of Science (Hons) in the Faculty of Science.

March 20, 2016



Copyright © 2016 Paul Ionele

ABSTRACT

This is a short example of an abstract.

ACKNOWLEDGEMENTS

Thank some people that you like here.

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of the University of Ontario Institute of Technology. The work is original except where indicated by special reference in the text and no part of the dissertation has been submitted for any other degree. Any views expressed in the dissertation are those of the author and in no way represent those of the University of Ontario Institute of Technology. The thesis has not been presented to any other University for examination either in Canada or elsewhere.

Paul Ionele
March 20, 2016

Contents

Abstract	ii
Acknowledgements	iii
Author's Declaration	iv
Table of Contents	v
List of Figures	vi
List of Tables	vii
1 Introduction	1
1.1 Diffusion	1
1.2 Monte Carlo Simulations	1
1.3 Master Equation Simulations	2
1.4 Simple Cells and Tissues	2
1.4.1 Tissues	3
2 Models and Simulations	4
2.1 1-Dimensional Systems	5
2.1.1 Homogenous System	5
2.1.2 Heterogeneous System	5
2.2 2-Dimensional System	5
3 Results and Analysis	7
4 Future Work and Conclusion	8
References	9
A Appendix	10

List of Figures

1.1	Composite cell showing various important and common internal cell structures. Most of the volume within the cell is occupied by the cytosol fluid. Figure courtesy of Essential Cell Biology, Alberts, 3rd. .	2
2.1	2D lattice unit cell with cellular and extracellular regions. Dashed lines indicate semi-permeable barrier.	6

List of Tables

1. INTRODUCTION

What was/is the purpose of this thesis? Why did we develop these computational simulations? Really, we need a goal!

In this thesis, a lattice Monte Carlo approach was used to simulate diffusion. Also used a finite difference method to simulate diffusion. Both problems were boundary-value problems.

We also performed an analysis: MSD, mean position, etc.?? Analysis is kind of empty!

Overall overview to thesis? What kind of an overview? Or just a general background to some main concepts needed or used in this thesis? How does this compare with the abstract?

1.1 Diffusion

Introduction. Theory. Supporting Figures.

Refer to page 91 in (Patton, Thibodeau , 2013). Figure 4.1 may also be used? Or is it too ‘simple’?

1.2 Monte Carlo Simulations

Introduction. Theory. Supporting figures.

1.3 Master Equation Simulations

Introduction. Theory. Supporting figures.

1.4 Simple Cells and Tissues

Nearly all human cells are microscopic in size; their diameters range from $7.5\text{ }\mu\text{m}$ to approximately $150\text{ }\mu\text{m}$ and a cell exhibits a particular size or shape that reflects the specific task it's designated to perform. There are many different types of cells including nerve cells, muscle cells, and gland cells, but despite their anatomical and functional differences, the cells of the human body have many similarities. It is a fact that no cell contains all cellular components found in all the cell types, so often a composite cell (Figure 1.1) is used to exhibit the most important characteristics.




Figure 1.1: Composite cell showing various important and common internal cell structures. Most of the volume within the cell is occupied by the cytosol fluid. Figure courtesy of Essential Cell Biology, Alberts, 3rd.

Each cell is enclosed by a plasma membrane that separates the cell contents from the surrounding environment. The inside of the cell is mostly composed of a gel-like substance called cytoplasm that is a dense arrangement of proteins, organelles, and

other molecules, suspended in a watery fluid called cytosol. The dense crowding of molecules and organelles results in frequent physical interactions which promotes high metabolic efficiency (Patton, Thibodeau , 2013). All of the fluid inside the cell may be referred to as intracellular fluid or simply, cellular fluid (CF).

The cellular fluid is separated from the extracellular fluid by the cell/plasma membrane. This membrane is a phospholipid bilayer with various embedded macromolecular structures. Each phospholipid molecule is amphipathic, having both a hydrophobic and hydrophilic region. A collection of phospholipid molecules will naturally arrange themselves into a bilayer that does not allow water, polar molecules, or ions to pass through easily. However, water and other molecules or ions need to traverse this membrane and the water transporting task is accomplished by aquaporin gated channel proteins. Aquaporins facilitate the passive diffusion of water through the plasma membrane, between the intracellular and extracellular regions. (Patton, Thibodeau , 2013).

1.4.1 Tissues

In a multicellular organism, there are several levels of biological organization. A cell is the lowest level of organization that is considered living; tissues are the next higher level of organization and are composed of cells similar in structure and function. This ensemble of cells resides in an extracellular matrix (ECM); a medium containing water, fibrous and adhesive proteins, glycoproteins, and other molecules. The ECM varies in composition between different tissues, but providing structural support and facilitating cell-to-cell communication are common functions of the ECM. **Normally, the cytoplasm is more viscous than the extracellular matrix (is this true?).** (Campbell, Reece , 2008).

2. MODELS AND SIMULATIONS

A simulation is intended to imitate in many cases, a real-world process or system, that may be too difficult or costly to analyze directly. Before any such simulation can begin, a model of the system studied must be constructed. Models capture the characteristics and behaviours of the system they represent and in general, a model should be as simple as possible (since resources are limited) while still explaining experimental observations and making predictions with a given degree of accuracy. The simulation is the implementation of the model and can be executed on a computer to produce data for testing, analysis, and visual presentation.

In our simplified model of a biological tissue, the relatively ordered and periodic nature of cells in most simple tissues is captured as a series of repeating *unit cells*. These unit cells are the building blocks of the heterogeneous 1D and 2D models. Each unit cell is characterized by a cellular domain, separated from an extracellular domain by a semi-permeable membrane. Both domains are isotropic everywhere except at the boundaries. Regarding the boundaries, there exists two kinds in our models. The first kind is a totally-reflecting boundary; it forms the absolute boundary of the model system and represents an insurmountable physical barrier. The second kind is a semi-permeable non-active/passive boundary and represents the selectively permeable nature of the plasma membrane. In a real biological plasma membrane, the integral membrane proteins may facilitate active or passive transport. More specifically, in the simple cell model, the semi-permeable membranes behave in a passive transport manner and this is implemented as a transition probability, a concept explained in

Section 1.1.

So what is being simulated? Currently, there is no net force on the particles (no directed motion) and the particles are non-interacting.

For all 1D and 2D models, both Monte Carlo and master equation approaches were used to simulate the diffusion of particles on a lattice.

Step size is fixed and equal in all directions

Particle motion occurs along a line in the 1D simulation and two orthogonal directions in the 2D simulation.

2.1 1-Dimensional Systems

In 1D, homogenous and heterogeneous simple cell systems were simulated using Monte Carlo (MC) and master equation approaches.

2.1.1 Homogenous System

include figure of homogenous model

2.1.2 Heterogeneous System

include figure of heterogeneous model

2.2 2-Dimensional System

include figure of heterogeneous model




Figure 2.1: 2D lattice unit cell with cellular and extracellular regions. Dashed lines indicate semi-permeable barrier.

3. RESULTS AND ANALYSIS

4. FUTURE WORK AND CONCLUSION

May just put this into the results section near the end as conclusions and future work.

REFERENCES

Navarro, J.F., Frenk, C.S., & White, S.D.M. 1995, MNRAS, 275, 720

Campbell, N.A., Reece, J.B., et al. 2008. Biology. 8th ed. Pearson Benjamin Cummings.

Patton K.T., Thibodeau G.A.. 2013, 8th ed. Elsevier.

A. APPENDIX

This is just an example of an appendix.