

Math Review (Not in the book)

ECON 5001
Prof. Healy

Updated 2026-08-12 at 23:26:01

Why a Math Review?

- Official prereqs: ECON 4001.0x and MATH 1151 (Calc 1)
- Reality: huge range of experience in the room
- Also, some non-calculus concepts
- Noting too hard! (IMO)

A **set** is simply a collection of things.

- Example: $A = \{\text{rock}, \text{paper}, \text{scissors}\}$
 - Shorthand names: $A = \{r, p, c\}$
 - Use curly brackets (“braces”)
 - Comma-separated list
 - Order doesn’t matter
 - No repetition
-
- “Singleton” set:
 - Empty set:

Set Operations

- Subset: $A \subseteq B$ and $A \subset B$
- Union: $A \cup B$
- Intersection: $A \cap B$
- Set difference (“set minus”): $A \setminus B$

Set Operations

- Universal set: set containing “all things” (e.g., possible die rolls) (only things we’re currently talking about)
- Complement: $A^c = U \setminus A$
- Element of: $\in$ (and not an element of: $\notin$)
- “Disjoint” (mutually exclusive) sets: $A \cap B = \emptyset$

Math Language: Quantifiers

- “For all” (or “for every”): $\forall$
- “There exists ___ such that”: $\exists$ ___ s.t.

Examples: Suppose $A = \{1, 3\}$, $B = \{-1, 2, 4\}$

Math Language: Quantifiers

- “For all” (or “for every”): $\forall$
- “There exists ___ such that”: $\exists$ ___ s.t.

Examples: Suppose $A = \{1, 3\}$, $B = \{-1, 2, 4\}$

- Is this true: $\forall x \in A \exists y \in B$ s.t. $y > x$
When $x = 1$ set $y = 2$. When $x = 3$ set $y = 4$ ✓

Math Language: Quantifiers

- “For all” (or “for every”): $\forall$
- “There exists ___ such that”: $\exists$ ___ s.t.

Examples: Suppose $A = \{1, 3\}$, $B = \{-1, 2, 4\}$

- Is this true: $\forall x \in A \exists y \in B$ s.t. $y > x$
When $x = 1$ set $y = 2$. When $x = 3$ set $y = 4$ ✓
- Is this true: $\exists x \in B$ s.t. $\forall y \in A$ $x > y$
Set $x = 4$. When $y = 1$ then $x > y$, and when $y = 3$ then $x > y$ ✓

Math Language: Quantifiers

- “For all” (or “for every”): $\forall$
- “There exists ___ such that”: $\exists$ ___ s.t.

Examples: Suppose $A = \{1, 3\}$, $B = \{-1, 2, 4\}$

- Is this true: $\forall x \in A \exists y \in B$ s.t. $y > x$
When $x = 1$ set $y = 2$. When $x = 3$ set $y = 4$ ✓
- Is this true: $\exists x \in B$ s.t. $\forall y \in A$ $x > y$
Set $x = 4$. When $y = 1$ then $x > y$, and when $y = 3$ then $x > y$ ✓
- Is this true: $\forall x \in B \exists y \in A$ s.t. $y > x$
?

Cartesian Product

The set $S = \{(r, r), (r, p), (r, c), (p, r), (p, p), (p, c), (c, r), (c, p), (c, c)\}$ contains **all possible combinations** of $A = \{r, p, c\}$ and $B = \{r, p, c\}$

This is called the **Cartesian product** (or just **product**) of A and B

Notation: $A \times B$.

Set B	r			
	p			
	c			
		r	p	c
		Set A		

Cartesian Product

The set $S = \{(r, r), (r, p), (r, c), (p, r), (p, p), (p, c), (c, r), (c, p), (c, c)\}$ contains **all possible combinations** of $A = \{r, p, c\}$ and $B = \{r, p, c\}$

This is called the **Cartesian product** (or just **product**) of A and B

Notation: $A \times B$.

Set B	r	(r, r)	(p, r)	(c, r)
	p	(r, p)	(p, p)	(c, p)
	c	(r, c)	(p, c)	(c, c)
		r	p	c
		Set A		


Cartesian Product

You've seen Cartesian products before:

Set B	r	(r, r)	(p, r)	(c, r)
	p	(r, p)	(p, p)	(c, p)
	c	(r, c)	(p, c)	(c, c)
		r	p	c
		Set A		


Cartesian Product

You've seen Cartesian products before:




Cartesian Product

You've seen Cartesian products before:



Cartesian Product

You've seen Cartesian products before:



Indexed Sets

Sets may be numbered (“indexed”). Example:

Player 1's strategies: $S_1 = \{a, b\}$

Player 2's strategies: $S_2 = \{c, d\}$

Player 3's strategies: $S_3 = \{x, y\}$

We can define $S = S_1 \times S_2 \times S_3$

We can do this for any number n sets: $S = S_1 \times S_2 \times \cdots \times S_n$

Shorthand notation: $S = \times_{i=1}^n S_i$

Lists

Suppose $S_1 = \{r, p, c\}$ and $S_2 = \{r, p, c\}$.

So $S_1 \times S_2 = \{(r, r), (r, p), (r, c), (p, r), (p, p), (p, c), (c, r), (c, p), (c, c)\}$.

Elements of this set are **lists**

Other names: **ordered pairs** (if 2), ***n*-tuples** (if *n*), **vectors**, **profiles**

List (r, c) means P1 played Rock and P2 played Scissors

Sets vs. Lists:

Sets	Lists
Ex: $\{a, b, c, d\}$	Ex: (a, d, d, a)
Braces $\{\dots\}$	Parentheses $(\dots)$
Comma-separated	Comma-separated
Order doesn't matter	Order matters!!!
Cannot repeat	Can repeat

Confusing: $(0, 10)$ can represent a list or an interval.

Cartesian Products & Lists

Rock-Paper-Scissors:

$$S_1 = \{r, p, c\}$$

$$S_2 = \{r, p, c\}$$

$$S = S_1 \times S_2 \text{ (Cartesian product)}$$

$$S = \{(r, r), (r, p), (r, c), (p, r), (p, p), (p, c), (c, r), (c, p), (c, c)\}$$

Each element of S is a **list**

List says (1) what P_1 chose in S_1 , (2) what P_2 chose in S_2

In game theory, lists of strategies are **strategy profiles**. e.g. (r, c) .

Strategy Profiles With n Players

Game theory notation:

- Suppose a game has n players
- What does this mean: “Each player i chooses some $s_i \in S_i$ ”
- Any particular player number: $i \in \{1, 2, \dots, n\}$
- Set of possible strategies for player i : S_i
- A particular strategy for player i : s_i

Player 1 chooses $s_1 \in S_1$

Player 2 chooses $s_2 \in S_2$

$\vdots$

Player n chooses $s_n \in S_n$

Strategy profile $s = (s_1, s_2, \dots, s_n) \in S = S_1 \times S_2 \times \dots \times S_n$

“Not i ” Notation: s_{-i}

- Suppose there are n players
- Each player i plays strategy s_i
- The profile is $s = (s_1, s_2, s_3, \dots, s_n)$
- Suppose we want to focus on player 7
 - “I am player 7”, so set $i = 7$
- In this case, $i = 7$. So $s_i = s_7$
- The strategies of others are $s_{-7} = (s_1, s_2, \dots, s_6, s_8, \dots, s_n)$
- For any player i , call this s_{-i}
 - $s_{-7} = (s_1, s_2, \dots, s_6, s_8, \dots, s_n)$
 - $s_{-3} = (s_1, s_2, s_4, \dots, s_n)$
 - $s_{-1} = (s_2, s_3, \dots, s_n)$
- The entire profile can be written as $s = (s_i, s_{-i})$
 - s_i is what player i plays
 - s_{-i} is what all the other players play

Functions

Function you're probably used to: $y = x^2$ or $f(x) = x^2$

What is this function doing? Mapping one set into another!

Domain vs. Range (or “Codomain”)

More General Functions

Any domain and any range. Ex: classes of animals.

Domain

Axolotl

Platypus

Komodo dragon

Coelacanth

Kiwi

Pangolin

Hammerhead shark

Hoatzin

Range

Fish

Amphibians

Reptiles

Birds


Mammals

More General Functions

Any domain and any range. Ex: classes of animals.

Domain

Range



A Game Theory Example

Rock-Paper-Scissors:

9 possible strategy pairs: $S = \left\{ \begin{array}{ccc} (r,r), & (r,p), & (r,c), \\ (p,r), & (p,p), & (p,c), \\ (c,r), & (c,p), & (c,c) \end{array} \right\}$

First letter: P1's choice. Second letter: P2's choice

Outcomes: $O = \{P1Wins, Tie, P2Wins\}$

“Outcome function”

A Game Theory Example

We can represent this in table format.

Convention: Player 1 is rows, Player 2 is columns

Strategy pairs:

		Set S_2		
		r	p	c
Set S_1	r	(r, r)	(r, p)	(r, c)
	p	(p, r)	(p, p)	(p, c)
	c	(c, r)	(c, p)	(c, c)

→

Outcome function:

		r	p	c
	r			
	p			
	c			

A Game Theory Example

We can represent this in table format.

Convention: Player 1 is rows, Player 2 is columns

Strategy pairs:

		Set S_2		
		r	p	c
Set S_1	r	(r, r)	(r, p)	(r, c)
	p	(p, r)	(p, p)	(p, c)
	c	(c, r)	(c, p)	(c, c)

→


Outcome function:

	r	p	c
r	Tie	P2	P1
p	P1	Tie	P2
c	P2	P1	Tie

A Bit of Calculus


Set of all “real” numbers: $\mathbb{R} = (-\infty, \infty)$ (interval)

“Real” function: domain and range are both $\mathbb{R}$




A Bit of Calculus

What is the slope at a point?




A Bit of Calculus

What is the derivative?



A Bit of Calculus

How to find the max of a function using calculus?



A Bit of Calculus

A worked example.

- Function: $f(x) = 4 + 2x - 6x^2$
- Derivative: $f'(x) =$
- Set equal to zero:
- Solve:

