

Untangling Knots Through Curve Repulsion

Joo-Hyun Paul Kim

March 2, 2023



What the curious folks ponder about

1 Introduction

2 Tangent-Point Energy

3 Gradient Flow

4 Curve Repulsion

Introduction

A Cool Knot




Figure: Imagine your earphones getting tangled like this...

Aim

- Finding a “homotopy” from a knot to an unknot.




Figure: Unknots of test knots.[3]

Aim

- Finding a “homotopy” from a knot to an unknot.




Figure: Unknots of test knots.[3]

- “Avoiding self-intersection”

General Strategy

- ① Define curve energy; penalizing the closeness of points on a curve.
 - Extreme-closeness of points on curve is a natural characteristic of a tangled curve.

General Strategy

- ① Define curve energy; penalizing the closeness of points on a curve.
 - Extreme-closeness of points on curve is a natural characteristic of a tangled curve.
- ② Attempt to decrease the curve energy by continuously deforming the curve.
 - We evolve the curve according to the gradient flow equation.
 - There is a freedom in choosing the “gradient” here.

General Strategy

- ① Define curve energy; penalizing the closeness of points on a curve.
 - Extreme-closeness of points on curve is a natural characteristic of a tangled curve.
- ② Attempt to decrease the curve energy by continuously deforming the curve.
 - We evolve the curve according to the gradient flow equation.
 - There is a freedom in choosing the “gradient” here.
- ③ We expect the stationary state to be the “unknot”
 - Or at least a simpler state...

Tangent-Point Energy

Defining Curve Energy

Given an (arc-length parameterised) curve $\gamma : M \rightarrow \mathbb{R}^3$, we wish to assign energy of the form:

$$\mathcal{E}(\gamma) := \iint_{M^2} k(\gamma_x, \gamma_y) \, d\gamma_x \, d\gamma_y \quad (1)$$

such that

- \mathcal{E} is very high when two non-neighbouring points are very close.

Defining Curve Energy

Given an (arc-length parameterised) curve $\gamma : M \rightarrow \mathbb{R}^3$, we wish to assign energy of the form:

$$\mathcal{E}(\gamma) := \iint_{M^2} k(\gamma_x, \gamma_y) \, d\gamma_x \, d\gamma_y \quad (1)$$



such that

- \mathcal{E} is very high when two non-neighbouring points are very close.

A naïve choice is $k(\gamma_x, \gamma_y) := \frac{1}{\|\gamma_x - \gamma_y\|}$



Pitfall of the “Simple Energy”

$$\mathcal{E}(\gamma) := \iint_{M^2} \frac{1}{||\gamma_x - \gamma_y||} d\gamma_x d\gamma_y$$





Pitfall of the “Simple Energy”

$$\mathcal{E}(\gamma) := \iint_{M^2} \frac{1}{||\gamma_x - \gamma_y||} d\gamma_x d\gamma_y$$



Pitfall of the “Simple Energy”

$$\mathcal{E}(\gamma) := \iint_{M^2} \frac{1}{\|\gamma_x - \gamma_y\|} d\gamma_x d\gamma_y$$



This energy is not well-defined for a lot of curves!

Buck-Orloff Tangent-Point Energy

- From the simple energy, need a way to suppress the infinite contribution of the “singularity”.

Buck-Orloff Tangent-Point Energy

- From the simple energy, need a way to suppress the infinite contribution of the “singularity”.

Definition (Buck-Orloff Tangent-Point Energy)

For a smooth curve γ , define

$$\mathcal{E}(\gamma) := \iint_{M^2} k_4^2(\gamma_x, \gamma_y, \mathbf{T}_x) d\gamma_x d\gamma_y$$

where \mathbf{T}_x is the unit tangent vector at γ_x , with the kernel defined as

$$k_4^2(\mathbf{p}, \mathbf{q}, \mathbf{T}) := \frac{\|\mathbf{T} \wedge (\mathbf{p} - \mathbf{q})\|^2}{\|\mathbf{p} - \mathbf{q}\|^4}$$



as **Buck-Orloff Tangent-Point Energy**.[1]

Intuition

What is the intuition behind the kernel $k_4^2(\mathbf{p}, \mathbf{q}, \mathbf{T}) := \frac{\|\mathbf{T} \wedge (\mathbf{p} - \mathbf{q})\|^2}{\|\mathbf{p} - \mathbf{q}\|^4}$?



Intuition

What is the intuition behind the kernel $k_4^2(\mathbf{p}, \mathbf{q}, \mathbf{T}) := \frac{\|\mathbf{T} \wedge (\mathbf{p} - \mathbf{q})\|^2}{\|\mathbf{p} - \mathbf{q}\|^4}$?



Intuition

What is the intuition behind the kernel $k_4^2(\mathbf{p}, \mathbf{q}, \mathbf{T}) := \frac{\|\mathbf{T} \wedge (\mathbf{p} - \mathbf{q})\|^2}{\|\mathbf{p} - \mathbf{q}\|^4}$?



Remark

Note that closer does not necessarily mean the kernel is larger.

Intuition




Figure: When two points are very close, the kernel converges to the curvature of the curve.

Example: Buck-Orloff Tangent-Point Energy of a Circle

Example


Suppose we wish to compute the Buck-Orloff tangent-point energy of a unit circle.

Example: Buck-Orloff Tangent-Point Energy of a Circle

Example

Suppose we wish to compute the Buck-Orloff tangent-point energy of a unit circle. Parameterise unit circle as:

$$\gamma(t) = (\cos t, \sin t, 0)$$




Example: Buck-Orloff Tangent-Point Energy of a Circle

Example

Suppose we wish to compute the Buck-Orloff tangent-point energy of a unit circle. Parameterise unit circle as:

$$\gamma(t) = (\cos t, \sin t, 0)$$



Then write:

$$\begin{cases} \gamma_x(\theta) = (\cos \theta, \sin \theta, 0) \\ \gamma_y(\phi) = (\cos \phi, \sin \phi, 0) \\ \mathbf{T}_x(\theta) = (-\sin \theta, \cos \theta, 0) \end{cases}$$

Example (Cont.)

$$\begin{cases} \gamma_x(\theta) = (\cos \theta, \sin \theta, 0) \\ \gamma_y(\phi) = (\cos \phi, \sin \phi, 0) \\ \mathbf{T}_x(\theta) = (-\sin \theta, \cos \theta, 0) \end{cases}$$

Substituting to Buck-Orloff Tangent-Point energy formula:

$$\mathcal{E}(\gamma) := \iint_{M^2} \frac{\|\mathbf{T}_x \wedge (\gamma_x - \gamma_y)\|^2}{\|\gamma_x - \gamma_y\|^4} d\gamma_x d\gamma_y$$

Example (Cont.)

$$\begin{cases} \gamma_x(\theta) = (\cos \theta, \sin \theta, 0) \\ \gamma_y(\phi) = (\cos \phi, \sin \phi, 0) \\ \mathbf{T}_x(\theta) = (-\sin \theta, \cos \theta, 0) \end{cases}$$

Substituting to Buck-Orloff Tangent-Point energy formula:

$$\mathcal{E}(\gamma) := \iint_{M^2} \frac{\|\mathbf{T}_x \wedge (\gamma_x - \gamma_y)\|^2}{\|\gamma_x - \gamma_y\|^4} d\gamma_x d\gamma_y$$

Using a few identities:

$$\mathcal{E}(\gamma) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{\|\mathbf{T}_x\|^2 \|\gamma_x - \gamma_y\|^2 - (\mathbf{T}_x \cdot (\gamma_x - \gamma_y))^2}{\|\gamma_x - \gamma_y\|^4} d\theta d\phi$$

Example (Cont.)

$$\mathcal{E}(\gamma) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{||\mathbf{T}_x||^2 ||\gamma_x - \gamma_y||^2 - (\mathbf{T}_x \cdot (\gamma_x - \gamma_y))^2}{||\gamma_x - \gamma_y||^4} d\theta d\phi \quad (2)$$

$$= \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{\sin^2 \frac{\theta-\phi}{2}}{(-1 + \cos(\theta - \phi))^2} d\theta d\phi \quad (3)$$

$$= \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{\sin^2 \frac{\theta-\phi}{2}}{\sin^4 \frac{\theta-\phi}{2}} d\theta d\phi \quad (4)$$

$$= \pi^2 \quad (5)$$

Example (Cont.)

$$\mathcal{E}(\gamma) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{||\mathbf{T}_x||^2 ||\gamma_x - \gamma_y||^2 - (\mathbf{T}_x \cdot (\gamma_x - \gamma_y))^2}{||\gamma_x - \gamma_y||^4} d\theta d\phi \quad (2)$$

$$= \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{\sin^2 \frac{\theta-\phi}{2}}{(-1 + \cos(\theta - \phi))^2} d\theta d\phi \quad (3)$$

$$= \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \frac{\sin^2 \frac{\theta-\phi}{2}}{\sin^4 \frac{\theta-\phi}{2}} d\theta d\phi \quad (4)$$

$$= \pi^2 \quad (5)$$

Remark

Note that (4) suggests the order at “singularity” is inverse-square.

General Tangent-Point Energy

A more general form of tangent-point energy comes from Yu, Schumacher, and Crane [3]:

Definition (Generalised Tangent-Point Energy)

$$\mathcal{E}_\beta^\alpha(\gamma) := \iint_{M^2} \frac{\|\mathbf{T}_x \wedge (\gamma_x - \gamma_y)\|^\alpha}{\|\gamma_x - \gamma_y\|^\beta} d\gamma_x d\gamma_y$$

where $\alpha > 1$ and $\beta \in [\alpha + 2, 2\alpha + 1]$

Remark

When $\alpha = 2$ and $\beta = 4$, we are back to Buck-Orloff.

Gradient Flow

Motivating Gradient Flow

For minimising a functional $\mathcal{E}(f)$, we wish to “move” f in the direction that decreases \mathcal{E} .

Motivating Gradient Flow

For minimising a functional $\mathcal{E}(f)$, we wish to “move” f in the direction that decreases \mathcal{E} .

Definition (Gradient Flow Equation)

For functional $\mathcal{E}(f)$, the **gradient flow equation** is: $\frac{\partial f}{\partial t} = - \text{grad}_X \mathcal{E}(f)$

Motivating Gradient Flow

For minimising a functional $\mathcal{E}(f)$, we wish to “move” f in the direction that decreases \mathcal{E} .

Definition (Gradient Flow Equation)

For functional $\mathcal{E}(f)$, the **gradient flow equation** is: $\frac{\partial f}{\partial t} = - \text{grad}_X \mathcal{E}(f)$

Remark

cf) For minimising a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, one may consider

$$\frac{\partial \mathbf{x}}{\partial t} = -\nabla f(\mathbf{x})$$




Figure: Steepest Descent [2]

Gradient in Space of Function?

(Gradient of Function)

$\nabla f(\mathbf{x})$ is such that for all $\mathbf{y} \in \mathbb{R}^n$

$$\nabla f(\mathbf{x}) \cdot \mathbf{y} = \frac{\partial}{\partial \epsilon} f(\mathbf{x} + \epsilon \mathbf{y}) \Big|_{\epsilon=0}$$

Gradient in Space of Function?

(Gradient of Function)

$\nabla f(\mathbf{x})$ is such that for all $\mathbf{y} \in \mathbb{R}^n$

$$\nabla f(\mathbf{x}) \cdot \mathbf{y} = \frac{\partial}{\partial \epsilon} f(\mathbf{x} + \epsilon \mathbf{y}) \Big|_{\epsilon=0}$$

(Gradient of Functional)

$\text{grad}_X E(f)$ is such that for all $g \in X$,

$$\langle \langle \text{grad}_X E, g \rangle \rangle_X = \frac{\partial}{\partial \epsilon} E(f + \epsilon g) \Big|_{\epsilon=0}$$

where X is the inner product space of functions (eg. L^2, H^1)

Gradient in Space of Function?

(Gradient of Function)

$\nabla f(\mathbf{x})$ is such that for all $\mathbf{y} \in \mathbb{R}^n$

$$\nabla f(\mathbf{x}) \cdot \mathbf{y} = \frac{\partial}{\partial \epsilon} f(\mathbf{x} + \epsilon \mathbf{y}) \Big|_{\epsilon=0}$$

(Gradient of Functional)

$\text{grad}_X E(f)$ is such that for all $g \in X$,

$$\langle \langle \text{grad}_X E, g \rangle \rangle_X = \frac{\partial}{\partial \epsilon} E(f + \epsilon g) \Big|_{\epsilon=0}$$

where X is the inner product space of functions (eg. L^2 , H^1)

Remark

For $X = L^2$, $(\text{grad}_X E)(f)$ is the first order variation from f with respect to ϵ .

Example: Dirichlet Energy in 1D

Definition (1D Dirichlet Energy)

For a differentiable function $f : \mathcal{I} \rightarrow \mathbb{R}$, define **1D Dirichlet energy**

$$E_D(f) := \int_{\mathcal{I}} |\nabla f(x)|^2 \, dx = \int_{\mathcal{I}} |f'(x)|^2 \, dx$$

Example: Dirichlet Energy in 1D

Definition (1D Dirichlet Energy)

For a differentiable function $f : \mathcal{I} \rightarrow \mathbb{R}$, define **1D Dirichlet energy**

$$E_D(f) := \int_{\mathcal{I}} |\nabla f(x)|^2 \, dx = \int_{\mathcal{I}} |f'(x)|^2 \, dx$$

Remark

Dirichlet energy is high for function f that varies a lot, and minimised by any constant function.

Example: L^2 Gradient Flow of Dirichlet Energy

Because Dirichlet energy is simple enough, we may explicitly write down $\text{grad}_X E$ for a few inner product function spaces.

Example: L^2 Gradient Flow of Dirichlet Energy

Because Dirichlet energy is simple enough, we may explicitly write down $\text{grad}_X E$ for a few inner product function spaces.

Lemma (L^2 Gradient of Dirichlet Energy)

$$\text{grad}_{L^2} E_D(f) = -\Delta f$$

(for suitable boundary condition if finite interval)

Example: L^2 Gradient Flow of Dirichlet Energy

Because Dirichlet energy is simple enough, we may explicitly write down $\text{grad}_X E$ for a few inner product function spaces.

Lemma (L^2 Gradient of Dirichlet Energy)

$$\text{grad}_{L^2} E_D(f) = -\Delta f$$

(for suitable boundary condition if finite interval)

Then the gradient flow equation for Dirichlet energy becomes:

$$\frac{\partial f}{\partial t} = \Delta f \tag{6}$$




Figure: Solution to L^2 Gradient Flow Equation for Dirichlet Energy (AKA Heat Equation)

Dirichlet Gradient Flow in Other Function Space

Definition (Famous Inner Product Function Spaces)

- $L^p = \left\{ f \left| \left(\int |f|^p dx \right)^{1/p} < \infty \right. \right\}$
 - $\langle \langle f, g \rangle \rangle_{L^2} = \int fg dx$
- $p = 2$ Sobolev Space $H^k = \left\{ f \left| \sum_{i=0}^k \int |\mathcal{D}^{(i)} f|^2 dx < \infty \right. \right\}$
 - $\langle \langle f, g \rangle \rangle_{H^k} = \sum_{i=0}^k \langle \langle \mathcal{D}^{(i)} f, \mathcal{D}^{(i)} g \rangle \rangle_{L^2}$
 - For our case, sufficient to take $\langle \langle f, g \rangle \rangle_{H^k} = \langle \langle \mathcal{D}^{(k)} f, \mathcal{D}^{(k)} g \rangle \rangle_{L^2}$

Dirichlet Gradient Flow in Other Function Space

Definition (Famous Inner Product Function Spaces)

- $L^p = \left\{ f \left| \left(\int |f|^p dx \right)^{1/p} < \infty \right. \right\}$
 - $\langle \langle f, g \rangle \rangle_{L^2} = \int fg dx$
- $p = 2$ Sobolev Space $H^k = \left\{ f \left| \sum_{i=0}^k \int |\mathcal{D}^{(i)} f|^2 dx < \infty \right. \right\}$
 - $\langle \langle f, g \rangle \rangle_{H^k} = \sum_{i=0}^k \langle \langle \mathcal{D}^{(i)} f, \mathcal{D}^{(i)} g \rangle \rangle_{L^2}$
 - For our case, sufficient to take $\langle \langle f, g \rangle \rangle_{H^k} = \langle \langle \mathcal{D}^{(k)} f, \mathcal{D}^{(k)} g \rangle \rangle_{L^2}$
- $L^2 = H^0: \frac{\partial f}{\partial t} = -\Delta f$
- $H^1: \frac{\partial f}{\partial t} = -f$
- $H^{-1}: \frac{\partial f}{\partial t} = -\Delta^2 f$
 - $\langle \langle f, g \rangle \rangle_{H^{-1}} = \langle \langle \Delta^{-1} f, \Delta^{-1} g \rangle \rangle_{H^1}$
- $H^2: \frac{\partial f}{\partial t} = \Delta^{-1} f$

Curve Repulsion

Gradient Flow on Tangent-Point Energy

We now take the tangent-point energy and gradient flow together:

$$\frac{\partial \gamma}{\partial t} = -\text{grad}_{L^2} \mathcal{E} \quad (7)$$

with $\gamma|_{t=0}$ is the tangled curve.

Gradient Flow on Tangent-Point Energy

We now take the tangent-point energy and gradient flow together:

$$\frac{\partial \gamma}{\partial t} = -\text{grad}_{L^2} \mathcal{E} \quad (7)$$

with $\gamma|_{t=0}$ is the tangled curve.

For numerical solution, one discretises the curve.




Figure: Discretised Curve

Gradient Flow on Tangent-Point Energy

We now take the tangent-point energy and gradient flow together:

$$\frac{\partial \gamma}{\partial t} = -\text{grad}_{L^2} \mathcal{E} \quad (7)$$

with $\gamma|_{t=0}$ is the tangled curve.

For numerical solution, one discretises the curve. Enumerate each point as $\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{J-1}$, and write Γ for the resulting polygonal curve.




Figure: Discretised Curve

Subtlety




Figure: Tangent-point energy is not well-defined for polygonal curves.

Energy Discretisation

Note the form of tangent-point energy:

$$\mathcal{E}_\beta^\alpha(\gamma) := \iint_{M^2} k_\beta^\alpha(\gamma_x, \gamma_y, \mathbf{T}_x) d\gamma_x d\gamma_y \quad (8)$$

Energy Discretisation

Note the form of tangent-point energy:

$$\mathcal{E}_\beta^\alpha(\gamma) := \iint_{M^2} k_\beta^\alpha(\gamma_x, \gamma_y, \mathbf{T}_x) d\gamma_x d\gamma_y \quad (8)$$

Definition (Discretised Energy)

Given a closed curve with discretised points $\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{J-1}$, define **discretised energy** E as:

$$E(\Gamma) := \sum_{\substack{i,j \in \{0,1,\dots,J-1\} \\ \text{dist}_{\text{geodesic}}(\Gamma(i), \Gamma(j)) > 1}} k_{i,j} \|\mathbf{x}_{i+1} - \mathbf{x}_i\| \|\mathbf{x}_{j+1} - \mathbf{x}_j\|$$

where the discretised kernel $k_{i,j}$ is given by:

$$\frac{1}{4} (k_\beta^\alpha(x_i, x_j, T_i) + k_\beta^\alpha(x_i, x_{j+1}, T_i) + k_\beta^\alpha(x_{i+1}, x_j, T_i) + k_\beta^\alpha(x_{i+1}, x_{j+1}, T_i))$$

Bibliography

- [1] Gregory Buck and Jeremey Orloff. “A simple energy function for knots”. In: *Topology and its Applications* 61.3 (Feb. 1995), pp. 205–214. doi: [10.1016/0166-8641\(94\)00024-w](https://doi.org/10.1016/0166-8641(94)00024-w).
- [2] “Chapter 8: Primal and Dual Projected Subgradient Methods”. In: *First-Order Methods in Optimization*, pp. 195–245. doi: [10.1137/1.9781611974997.ch8](https://doi.org/10.1137/1.9781611974997.ch8). eprint: <https://epubs.siam.org/doi/pdf/10.1137/1.9781611974997.ch8>. URL: <https://epubs.siam.org/doi/abs/10.1137/1.9781611974997.ch8>.
- [3] Chris Yu, Henrik Schumacher, and Keenan Crane. “Repulsive Curves”. In: *ACM Transactions on Graphics* 40.2 (Apr. 2021), pp. 1–21. doi: [10.1145/3439429](https://doi.org/10.1145/3439429).