Tackling 
$$u_t = \Delta \left( u^2 \right)$$

Paul Kim

December 31, 2022

## 1 The Equation

The equation we are concerned with is:

$$u_t = \Delta \left( u^2 \right) \tag{1}$$

## 2 Explicit Scheme

For convenience we consider explicit Euler schemes.

## 2.1 Scheme Construction After Simplification

Note that (1) can be written as:

$$u_t = \Delta \left( u^2 \right) \tag{2}$$

$$=2u\Delta u + 2||\nabla u||^2\tag{3}$$

In 1-dimensional case, we simplify it as:

$$\frac{\partial u}{\partial t} = 2u \frac{\partial^2 u}{\partial x^2} + 2\left(\frac{\partial u}{\partial x}\right)^2 \tag{4}$$

Explicit Euler scheme can now be written

$$\frac{U_j^{m+1} - U_j^m}{\Delta t} = 2U_j^m \left( \frac{U_{j+1}^m - 2U_j^m + U_{j-1}^m}{\Delta x^2} \right) + 2\left( \frac{U_{j+1}^m - U_{j-1}^m}{2\Delta x} \right)^2 \tag{5}$$

$$=2U_{j}^{m}\left(\frac{U_{j+1}^{m}-2U_{j}^{m}+U_{j-1}^{m}}{\Delta x^{2}}\right)+\frac{\left(U_{j+1}^{m}-U_{j-1}^{m}\right)^{2}}{2\Delta x^{2}}\tag{6}$$

which can be rewritten as:

$$U_j^{m+1} = 2\mu U_j^m \left( U_{j+1}^m - 2U_j^m + U_{j-1}^m \right) + \frac{1}{2}\mu \left( U_{j+1}^m - U_{j-1}^m \right)^2 \tag{7}$$