

The **discretized energy** E can be expressed as:

$$E = \sum_{i=1}^J \sum_{\substack{j=1 \\ |j-i|>1}}^J k_{i,j} \|x_{i+1} - x_i\| \|x_{j+1} - x_j\| \quad (1)$$

$$k_{i,j} = \frac{1}{4} (k_{\beta}^{\alpha}(x_i, x_j, T_i) + k_{\beta}^{\alpha}(x_i, x_{j+1}, T_i) + k_{\beta}^{\alpha}(x_{i+1}, x_j, T_i) + k_{\beta}^{\alpha}(x_{i+1}, x_{j+1}, T_i)) \quad (2)$$

where we index cyclically.

For derivative with respect to x_k (note that it is a vector), the only terms indexed by (i, j) that involves x_k are enumerated by the following indices:

- $(k, 1), \dots, (k, k-2), (k, k+2), \dots, (k, J)$
- $(k-1, 1), \dots, (k-1, k-3), (k-1, k+1), \dots, (k-1, J)$
- $(1, k), \dots, (k-2, k), (k+2, k), \dots, (J, k)$
- $(1, k-1), \dots, (k-3, k-1), (k+1, k-1), \dots, (J, k-1)$

We now attempt to construct derivative (in a “modular fashion”). Write

$$k_{\beta}^{\alpha}(x_p, x_q, T_r) = k_{\beta}^{\alpha}\left(x_p, x_q, \frac{x_{r+1} - x_r}{\|x_r - x_{r+1}\|}\right) \quad (3)$$

$$= \frac{\sqrt{\|x_{r+1} - x_r\|^2 \|x_p - x_q\|^2 - ((x_{r+1} - x_r) \cdot (x_p - x_q))^2}^{\alpha}}{\|x_p - x_q\|^{\beta} \|x_r - x_{r+1}\|^{\alpha}} \quad (4)$$

$$= \frac{\xi_{p,q,r}^{\alpha/2}}{\eta_{p,q,r}} \quad (5)$$

where

$$\xi_{p,q,r} = \|x_{r+1} - x_r\|^2 \|x_p - x_q\|^2 - ((x_{r+1} - x_r) \cdot (x_p - x_q))^2 \quad (6)$$

$$\eta_{p,q,r} = \|x_p - x_q\|^{\beta} \|x_r - x_{r+1}\|^{\alpha} \quad (7)$$

Derivative with respect to x_k can now be written as:

$$\frac{\partial}{\partial x_k} \left(\frac{\xi^{\alpha/2}}{\eta} \right) = \frac{1}{\eta^2} \left(\frac{\alpha}{2} \xi^{\alpha/2-1} \frac{\partial \xi}{\partial x_k} \eta - \xi^{\alpha/2} \frac{\partial \eta}{\partial x_k} \right) \quad (8)$$

where p, q, r are omitted.