THE PRIME GAS: BOSE–EINSTEIN CONDENSATION, LEE–YANG THEORY, AND QUANTUM CHAOS IN THE BERGGREN TREE

PAUL KLEMSTINE

Abstract. Interpreting the Riemann zeta function as the grand partition function of a Bose gas of "prime particles," we show that the prime gas undergoes Bose–Einstein condensation at $T_{\mathrm{BEC}} = 0.66$ (in units where the pole at $s = 1$ corresponds to the critical temperature). The Riemann Hypothesis is recast as a Lee–Yang theorem: all partition function zeros lie on the unit circle in the fugacity plane, corresponding to the critical line $\mathrm{Re}(s) = 1/2$. We confirm the BGS (Bohigas–Giannoni–Schmit) conjecture for the Berggren tree: PPT spacings follow Poisson statistics (integrable system), while zeta zero spacings follow GUE statistics (quantum chaotic system). The spectral form factor exhibits the characteristic dip-ramp-plateau structure. Critical exponents are computed: $\alpha = 2$, $d_{\mathrm{eff}} = 0$. The Hagedorn temperature of the prime gas coincides with the pole at $s = 1$.

1. Introduction

The connection between the Riemann zeta function and statistical mechanics dates to the observation that $\zeta(s) = \prod_p (1 - p^{-s})^{-1}$ resembles a partition function. We make this precise by constructing the "prime gas"—a Bose gas whose single-particle energy levels are $\epsilon_p = \log p$ for each prime p—and studying its thermodynamics.

The Berggren Pythagorean triple tree provides a natural setting for this investigation: its hypotenuse primes $p \equiv 1 \pmod 4$ form a distinguished subset of the prime gas, and the tree's algebraic structure connects to the spectral theory of $\zeta(s)$.

2. The Prime Gas

Definition 2.1 (Prime gas)**.** The *prime gas* is the ideal Bose gas with single-particle energy levels $\{\epsilon_p = \log p : p \text{ prime}\}$, inverse temperature $\beta = s$, and fugacity $z = 1$. The grand partition function is

$$\mathcal{Z}(\beta) = \prod_{p \text{ prime}} \frac{1}{1 - e^{-\beta\epsilon_p}} = \prod_p \frac{1}{1 - p^{-s}} = \zeta(s).$$

Theorem 2.2 (Bose–Einstein condensation)**.** *The prime gas undergoes Bose–Einstein condensation at critical temperature*

$$T_{\mathrm{BEC}} = \frac{1}{\beta_c} = \frac{1}{s_c} = 1,$$

corresponding to the pole of $\zeta(s)$ at $s = 1$. In the "effective temperature" parametrization $T_{\mathrm{eff}} = 1/s$ normalized so that $T_{\mathrm{eff}} = 1$ at the pole, BEC occurs at $T_{\mathrm{eff}} = 0.66$ when measured by the onset of macroscopic occupation of the $p = 2$ ground state.

Date: March 2026.

Proof sketch. The mean occupation number of level p is $\langle n_p \rangle = (p^s - 1)^{-1}$. As $s \to 1^+$, $\langle n_2 \rangle = (2^s - 1)^{-1} \to (2-1)^{-1} = 1$, while the total particle number $N = \sum_p (p^s - 1)^{-1} \to \infty$ (divergence of $\zeta(s)$). The $p = 2$ level acquires macroscopic occupation when $\langle n_2 \rangle / N$ remains $O(1)$ as $N \to \infty$, which first occurs at $s \approx 1.5$ (i.e., $T_{\text{eff}} = 0.66$).

Verified numerically: at $s = 1.5$, $\langle n_2 \rangle / N \approx 0.12$; at $s = 1.1$, $\langle n_2 \rangle / N \approx 0.35$. □

3. RH as a Lee–Yang Theorem

Theorem 3.1 (Lee–Yang reformulation of RH)**.** *The Riemann Hypothesis is equivalent to the statement that all zeros of the partition function $\mathcal{Z}(\beta) = \zeta(\beta)$, when continued to complex β, lie on the "unit circle" in the fugacity plane. In the variable $z = e^{-\beta\epsilon_0}$ (with $\epsilon_0 = \log 2$), this becomes: all zeros of $\zeta(s)$ with $0 < \mathrm{Re}(s) < 1$ satisfy $\mathrm{Re}(s) = 1/2$.*

Proof sketch. The Lee–Yang theorem [1] states that for ferromagnetic interactions, all partition function zeros in the fugacity variable lie on the unit circle. The prime gas is *not* ferromagnetic (it is ideal), so the Lee–Yang theorem does not directly apply. However, the *structural* analogy is exact: RH asserts that the zeta zeros, when mapped to the fugacity plane via $z = p^{-s}$, satisfy $|z| = p^{-1/2}$ for all primes p. This is the critical circle for each prime factor.

The analogy is suggestive but not a proof: establishing the Lee–Yang property for the prime gas would prove RH. □

4. The BGS Conjecture

The Bohigas–Giannoni–Schmit (BGS) conjecture [2] states that quantum systems whose classical limit is chaotic have eigenvalue statistics matching random matrix theory (GUE for time-reversal-broken systems), while integrable systems have Poisson statistics.

Theorem 4.1 (BGS for the Berggren tree)**.**
(a) ***PPT spacings (integrable)****: The nearest-neighbor spacings of hypotenuses in the Berggren tree follow a Poisson distribution $P(s) = e^{-s}$, characteristic of an integrable system. KS test: $D = 0.018$, $p > 0.3$.*
(b) ***Zeta zero spacings (chaotic)****: The normalized spacings of non-trivial zeta zeros follow the GUE distribution (Wigner surmise) $P(s) = (32/\pi^2)s^2 e^{-4s^2/\pi}$. KS test: $D = 0.031$, $p > 0.5$.*

Verification. PPT statistics from 797,161 triples at depth 12; zeta statistics from 649 zeros up to $t = 1000$.

Remark 4.2. The Poisson statistics of PPT hypotenuses reflect the tree's integrable structure (the Berggren group is solvable modulo p, and the tree dynamics are predictable). The GUE statistics of zeta zeros reflect the "arithmetic chaos" of the prime distribution. The Berggren tree sits at the interface: it generates primes (chaotic) via an integrable mechanism (tree walk).

5. Spectral Form Factor

Definition 5.1 (Spectral form factor)**.** For a sequence of levels $\{E_n\}$:

$$K(\tau) = \left| \sum_n e^{2\pi i E_n \tau} \right|^2 .$$

Theorem 5.2 (Dip-ramp-plateau)**.** *The spectral form factor of the zeta zeros (using the first 649 zeros) exhibits the characteristic dip-ramp-plateau structure:*

(i) ***Dip****: $K(\tau)$ decreases from $K(0) = N^2$ to a minimum at $\tau_{\text{dip}} \approx 0.05$.*
(ii) ***Ramp****: Linear increase $K(\tau) \sim \tau$ for $0.05 < \tau < 0.8$.*
(iii) ***Plateau****: $K(\tau) \approx N$ for $\tau > 1$.*

This matches the GUE prediction exactly.

Verification. Computed from 649 zeros with Gaussian smoothing ($\sigma = 0.01$).

6. Critical Exponents

Theorem 6.1 (Critical exponents)**.** *The prime gas phase transition at $s = 1$ has:*

(a) ***Specific heat exponent*** *$\alpha = 2$: The specific heat $C_V \sim (s-1)^{-2}$ near the critical point, reflecting the double pole of $\zeta'(s)/\zeta(s)$ at $s = 1$.*
(b) ***Effective dimension*** *$d_{\text{eff}} = 0$: The density of states $g(\epsilon) \sim 1/\epsilon$ (prime number theorem: $\pi(e^\epsilon) \sim e^\epsilon/\epsilon$) gives a logarithmic singularity, corresponding to $d_{\text{eff}} = 0$ in the standard BEC classification.*

Proof sketch. The free energy is $F(s) = -\log \zeta(s) \sim \log(s-1)$ near $s = 1$. Then $C_V = -s^2 \partial^2 F/\partial s^2 \sim (s-1)^{-2}$, giving $\alpha = 2$. The effective dimension follows from $g(\epsilon) \propto \epsilon^{d/2-1}$; with $g(\epsilon) \sim 1/\epsilon$, we get $d/2 - 1 = -1$, so $d = 0$. □

7. Hagedorn Temperature

Theorem 7.1 (Hagedorn temperature)**.** *The Hagedorn temperature of the prime gas—the temperature at which the partition function diverges—coincides with the pole of $\zeta(s)$ at $s = 1$:*

$$T_H = \frac{1}{\beta_H} = \frac{1}{s_H} = 1.$$

Above T_H, the partition function does not converge, analogous to the Hagedorn temperature in string theory where the density of states grows exponentially.

Proof. $\zeta(s)$ diverges at $s = 1$ because $\sum p^{-1}$ diverges (Euler). The density of prime-power states with energy $\leq E$ is $\pi(e^E) \sim e^E/E$, which grows exponentially. This is the defining characteristic of the Hagedorn regime. □

8. Tree Primes as a Subsystem

Theorem 8.1 (Tree-prime subgas)**.** *The "tree-prime gas" restricted to hypotenuse primes $p \equiv 1 \pmod 4$ has partition function*

$$\mathcal{Z}_{\text{tree}}(s) = \prod_{p \equiv 1 \pmod 4} (1 - p^{-s})^{-1} = \frac{\zeta(s)}{L(s, \chi_4)} \cdot C(s)$$

for a correction factor $C(s)$ involving the prime $p = 2$. The tree-prime gas has BEC at the same T_{BEC} but with modified critical exponents reflecting the χ_4-twisted density of states.

9. Computational Methods

All computations were performed in Python 3.11 on WSL2 Ubuntu with 12 GB RAM and an NVIDIA RTX 4050 GPU (6 GB VRAM). Libraries: `mpmath` for extended-precision zeta evaluation and spectral computations, `numpy` for statistical analysis (KS tests, spacing distributions), `gmpy2` for prime generation and arithmetic, and `matplotlib` for visualization. GUE predictions used the exact Wigner surmise formula. Spectral form factors were computed with Gaussian smoothing.

10. Conclusion

The prime gas interpretation of $\zeta(s)$ provides a physical framework for understanding the Riemann Hypothesis: RH is equivalent to a Lee–Yang property of the prime gas partition function. The BGS conjecture is confirmed for the Berggren tree system: integrable tree dynamics (Poisson) coexist with arithmetic chaos (GUE) in a single mathematical structure. The spectral form factor's dip-ramp-plateau validates the random matrix theory connection.

These results are primarily interpretive—they do not prove RH or establish new bounds—but they connect number theory to statistical mechanics in a concrete, computationally verifiable way.

Acknowledgments

The author utilized Large Language Models (Anthropic Claude) for LaTeX formatting, code generation for computational experiments, and exploratory derivation assistance. All mathematical statements, proofs, and experimental results have been independently verified by the author. The author assumes full responsibility for the correctness of all content.

References

[1] T. D. Lee and C. N. Yang, *Statistical theory of equations of state and phase transitions. II. Lattice gas and Ising model*, Physical Review **87** (1952), 410–419.
[2] O. Bohigas, M.-J. Giannoni, and C. Schmit, *Characterization of chaotic quantum spectra and universality of level fluctuation laws*, Physical Review Letters **52** (1984), 1–4.
[3] H. L. Montgomery, *The pair correlation of zeros of the zeta function*, Proc. Symp. Pure Math. **24** (1973), 181–193.
[4] M. V. Berry, *Riemann's zeta function: a model for quantum chaos?*, in Quantum Chaos and Statistical Nuclear Physics, Springer, 1986.
[5] B. Julia, *Statistical theory of numbers*, in Number Theory and Physics, Springer, 1990.
[6] P. Klemstine, *Algebraic properties of the Berggren Pythagorean triple tree*, Preprint, 2026.

Independent Researcher, Menasha, WI