FROM PYTHAGOREAN TRIPLES TO THE MONSTER: A TOWER OF EXCEPTIONAL STRUCTURES VIA THE THETA GROUP

PAUL KLEMSTINE

Abstract. We construct an explicit chain of group-theoretic connections from the elementary $(3, 4, 5)$ Pythagorean triple to the Monster group $\mathbb{M}$, the largest sporadic simple group of order $\approx 8 \times 10^{53}$. The chain proceeds through eight stages:

$$(3, 4, 5) \to \Gamma_\theta \to \mathrm{SL}(2, \mathbb{F}_p) \to E_6 \to E_8 \to \text{Klein} \to M_{11} \to M_{24} \to \text{Leech} \to \mathbb{M}.$$

The key observation is that the Berggren matrices generating the tree of all primitive Pythagorean triples (PPTs) generate the theta group $\Gamma_\theta = \langle S, T^2 \rangle$, an index-3 subgroup of $\mathrm{SL}(2, \mathbb{Z})$. Reduction modulo primes $p = 3, 5, 7, 11, 23$ connects PPTs to the binary tetrahedral group (E_6 under McKay), the binary icosahedral group (E_8), the Klein quartic group, the Mathieu group M_{11}, and the Mathieu group M_{24} acting on $\mathbb{P}^1(\mathbb{F}_{23})$. The 24 points of this projective line give the rank of the Leech lattice, whose automorphism group Co_0 embeds in the Monster via Griess's construction. As a parallel thread, the modular lambda function $\lambda(\tau) = (\theta_2/\theta_3)^4$, a Hauptmodul for Γ_θ, determines the j-invariant $j(\tau) = 256(1 - \lambda + \lambda^2)^3/(\lambda(1 - \lambda))^2$, whose Fourier coefficients are the representation dimensions of $\mathbb{M}$ (monstrous moonshine). Every link in the chain is either proven here or cited to a precise theorem in the literature, and all finite-group claims are verified computationally.

Contents

1. Introduction 1
The tower 1
Significance 2
What is proven vs. cited 2
2. The Theta Group Γ_θ 2
2.1. The Berggren matrices 2
2.2. Berggren generates the theta group 2
2.3. The theta group as modular group 3
3. The $\mathrm{SL}(2, \mathbb{F}_p)$ Theorem 3
4. ADE Classification 4
5. Klein Quartic and Exceptional Isomorphisms 4
6. Sporadic Groups: Mathieu Groups 5
6.1. M_{11} from $p = 11$ 5
6.2. M_{24} from $p = 23$ 5
7. The Leech Lattice 6
8. Monstrous Moonshine 6
8.1. From Γ_θ to the j-invariant 6
8.2. The moonshine module 7

Date: March 2026.

8.3. The complete chain 7
9. Physical Interpretation 7
9.1. Bosonic string theory 7
9.2. E_8 heterotic string 8
9.3. Gravitational partition function 8
10. Computational Verification 8
10.1. Group generation modulo p 8
10.2. McKay graph verification 8
10.3. Modular function computation 8
10.4. Golay code verification 8
10.5. Platform 8
11. Discussion 9
11.1. Uniqueness of the Berggren tree 9
11.2. The role of primes 9
11.3. Open problems 9
Acknowledgments 9
References 9

1. Introduction

A *primitive Pythagorean triple* (PPT) is a triple (a, b, c) of positive integers satisfying $a^2 + b^2 = c^2$ with $\gcd(a, b, c) = 1$. In 1934, Berggren [2] proved that the set of all PPTs carries the structure of an infinite ternary tree rooted at $(3, 4, 5)$, generated by three 3×3 integer matrices. This elementary combinatorial object has surprising connections to some of the deepest structures in mathematics.

The tower. We establish a chain of explicit connections:

(1)
$$\underbrace{(3,4,5)}_{\text{root PPT}} \xrightarrow{\S 2} \underbrace{\Gamma_\theta}_{\text{theta group}} \xrightarrow{\S 3} \underbrace{\mathrm{SL}(2,\mathbb{F}_p)}_{\text{mod } p} \xrightarrow{\S 4} \underbrace{E_6,\ E_8}_{\text{ADE}} \xrightarrow{\S 5} \underbrace{\text{Klein}}_{\text{quartic}} \xrightarrow{\S 6} \underbrace{M_{11},\ M_{24}}_{\text{Mathieu}} \xrightarrow{\S 7} \underbrace{\Lambda_{24}}_{\text{Leech}} \xrightarrow{\S 8} \underbrace{\mathbb{M}}_{\text{Monster}}$$

The first three arrows are proven directly. The ADE and Klein links follow from the McKay correspondence and classical isomorphisms. The sporadic and Leech connections use the action of $\mathrm{PSL}(2, \mathbb{F}_{23})$ on $\mathbb{P}^1(\mathbb{F}_{23})$. The final arrow is monstrous moonshine, which we access through the modular lambda function—itself a Hauptmodul for Γ_θ.

Significance. The tower is remarkable for two reasons. First, it connects one of the oldest objects in mathematics (Pythagorean triples, known to the Babylonians circa 1800 BCE) to one of the most modern (the Monster, classified in 1981). Second, the chain passes through essentially every exceptional structure in mathematics: the E-type Dynkin diagrams, the Klein quartic (the unique genus-3 Hurwitz surface), the Mathieu groups (the first sporadic groups discovered), the Leech lattice (the densest sphere packing in 24 dimensions), and the Monster (which controls monstrous moonshine and the j-function).

What is proven vs. cited. Links 1–3 are proved in full in Sections 2–3. Link 4 (ADE) follows from the McKay correspondence (McKay [13], Steinberg [18]). Link 5 (Klein) uses the exceptional isomorphism $\mathrm{PSL}(2, \mathbb{F}_7) \cong \mathrm{GL}(3, \mathbb{F}_2)$. Links 6–7 (Mathieu, Leech) follow from

the classification of sharply k-transitive groups and Conway's work on the Leech lattice [6]. Link 8 (Monster) is the Borcherds–Conway–Norton moonshine theorem [3, 5]. We make no claim to have discovered these individual results; our contribution is the *unified chain* from PPTs to $\mathbb{M}$ and the observation that Γ_θ is the organizing thread.

2. The Theta Group Γ_θ

2.1. **The Berggren matrices.** Every PPT is obtained from $(3, 4, 5)$ by successive application of

$$B_1 = \begin{pmatrix} 1 & -2 & 2 \\ 2 & -1 & 2 \\ 2 & -2 & 3 \end{pmatrix}, \quad B_2 = \begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 3 \end{pmatrix}, \quad B_3 = \begin{pmatrix} -1 & 2 & 2 \\ -2 & 1 & 2 \\ -2 & 2 & 3 \end{pmatrix}. \tag{2}$$

These matrices preserve the quadratic form $Q(x, y, z) = x^2 + y^2 - z^2$, so the Berggren monoid embeds in $\mathrm{SO}(Q; \mathbb{Z}) \cong \mathrm{SO}(2, 1; \mathbb{Z})$.

Via the parametrization $(a, b, c) = (m^2 - n^2, 2mn, m^2 + n^2)$, the 3×3 matrices act on (m, n) via 2×2 matrices:

$$M_1 = \begin{pmatrix} 2 & -1 \\ 1 & 0 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}, \quad M_3 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}. \tag{3}$$

We have $\det M_1 = \det M_3 = 1$ and $\det M_2 = -1$, so $M_1, M_3 \in \mathrm{SL}(2, \mathbb{Z})$ while $M_2 \in \mathrm{GL}(2, \mathbb{Z}) \setminus \mathrm{SL}(2, \mathbb{Z})$.

2.2. **Berggren generates the theta group.**

Theorem 2.1 (Berggren = Theta Group)**.** *The group $\langle M_1, M_3 \rangle \subset \mathrm{SL}(2, \mathbb{Z})$ equals the theta group $\Gamma_\theta = \langle S, T^2 \rangle$, an index-3 subgroup of $\mathrm{SL}(2, \mathbb{Z})$, where $S = \left(\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}\right)$ and $T = \left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right)$.*

Proof. Direct computation:

$$M_3^{-1} M_1 = \begin{pmatrix} 1 & -2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = S.$$

Since $M_3 = T^2$, we have $S \in \langle M_1, M_3 \rangle$ and $T^2 \in \langle M_1, M_3 \rangle$, giving $\Gamma_\theta \subset \langle M_1, M_3 \rangle$. Conversely, $M_1 = M_3 S = T^2 S$ and $M_3 = T^2$, so $\langle M_1, M_3 \rangle \subset \Gamma_\theta$.

The theta group is classically known to have index 3 in $\mathrm{SL}(2, \mathbb{Z})$ with coset representatives $\{I, T, T^{-1}\}$. □

Remark 2.2 (Uniqueness of the PPT tree). The Berggren tree is the *unique* tree of all PPTs compatible with the arithmetic Γ_θ-action. The only other known PPT tree (due to Price [16]) uses a different set of generators that do *not* generate Γ_θ; the Berggren tree is distinguished by its modular-group origin.

2.3. **The theta group as modular group.** The theta group Γ_θ is the automorphism group of the Jacobi theta function

$$\theta(\tau) = \sum_{n \in \mathbb{Z}} q^{n^2}, \qquad q = e^{2\pi i \tau}, \quad \tau \in \mathbb{H}. \tag{4}$$

It is a genus-0 congruence subgroup with two cusps (at 0 and ∞) and one elliptic point of order 2. The modular lambda function

$$\lambda(\tau) = \left(\frac{\theta_2(\tau)}{\theta_3(\tau)} \right)^4 \tag{5}$$

is a Hauptmodul for Γ_θ: it generates the field of meromorphic functions on $\mathbb{H}/\Gamma_\theta$ and provides a bijection $\mathbb{H}/\Gamma_\theta \xrightarrow{\sim} \mathbb{P}^1(\mathbb{C}) \setminus \{0, 1, \infty\}$.

3. The $\mathrm{SL}(2, \mathbb{F}_p)$ Theorem

Theorem 3.1 (Surjectivity)**.** *For every odd prime p, the reduction map $\Gamma_\theta \to \mathrm{SL}(2, \mathbb{F}_p)$ is surjective.*

Proof. Since p is odd, 2 is invertible in $\mathbb{F}_p$. Thus $T = (T^2)^{2^{-1} \bmod p} \in \langle T^2 \rangle \bmod p$. Since S and T generate $\mathrm{SL}(2, \mathbb{Z})$, their reductions generate $\mathrm{SL}(2, \mathbb{F}_p)$. □

Remark 3.2. At $p = 2$, the map fails: $T^2 \equiv I \pmod 2$, so $\langle S, T^2 \rangle \bmod 2 = \langle S \rangle$ has order 2, while $|\mathrm{SL}(2, \mathbb{F}_2)| = 6$. This is consistent with $[\mathrm{SL}(2, \mathbb{Z}) : \Gamma_\theta] = 3$.

Table 1. Computational verification: $|\langle M_1, M_3 \rangle \bmod p| = |\mathrm{SL}(2, \mathbb{F}_p)| = p(p^2 - 1)$ for odd primes $p \leq 31$.

p	$\lvert\langle M_1, M_3\rangle \bmod p\rvert$	$\lvert\mathrm{SL}(2,\mathbb{F}_p)\rvert$	Status
3	24	24	✓
5	120	120	✓
7	336	336	✓
11	1 320	1 320	✓
13	2 184	2 184	✓
17	4 896	4 896	✓
19	6 840	6 840	✓
23	12 144	12 144	✓
29	24 360	24 360	✓
31	29 760	29 760	✓

Corollary 3.3 (Expander family)**.** *The Cayley graphs* $\mathrm{Cay}(\mathrm{SL}(2, \mathbb{F}_p), \{M_1^{\pm 1}, M_3^{\pm 1}\})$ *form an expander family by the Bourgain–Gamburd theorem* [4].

4. ADE Classification

The McKay correspondence [13] associates to each finite subgroup $G \subset \mathrm{SL}(2, \mathbb{C})$ a simply-laced Dynkin diagram via the tensor product of the natural 2-dimensional representation with each irreducible representation of G. The classification is:

$$\begin{array}{lcl} \mathbb{Z}/n\mathbb{Z} & \longleftrightarrow & A_{n-1} \\ \text{binary dihedral } \widetilde{D}_n & \longleftrightarrow & D_{n+2} \\ \text{binary tetrahedral } \widetilde{T} & \longleftrightarrow & E_6 \\ \text{binary octahedral } \widetilde{O} & \longleftrightarrow & E_7 \\ \text{binary icosahedral } \widetilde{I} & \longleftrightarrow & E_8 \end{array}$$

Theorem 4.1 (ADE Tower from PPTs)**.** *The surjection $\Gamma_\theta \twoheadrightarrow \mathrm{SL}(2, \mathbb{F}_p)$ specializes to:*

(i) $p = 3$, E_6: $\mathrm{SL}(2, \mathbb{F}_3)$ is the binary tetrahedral group $\widetilde{T}$ of order 24, corresponding to E_6 under McKay. Its McKay graph has 7 vertices with adjacency matrix equal to the extended E_6 Dynkin diagram.

(ii) $p = 5$, E_8: $\mathrm{SL}(2, \mathbb{F}_5)$ *is the binary icosahedral group* $\widetilde{I}$ *of order* 120*, corresponding to* E_8 *under McKay. Its McKay graph has* 9 *vertices matching the extended* E_8 *Dynkin diagram.*

Proof. That $\mathrm{SL}(2, \mathbb{F}_3) \cong \widetilde{T}$ follows from $|\mathrm{SL}(2, \mathbb{F}_3)| = 3(9-1) = 24 = |\widetilde{T}|$ combined with the unique group of order 24 that is a central extension $\mathbb{Z}/2\mathbb{Z} \to \widetilde{T} \to A_4$.

That $\mathrm{SL}(2, \mathbb{F}_5) \cong \widetilde{I}$ follows from $|\mathrm{SL}(2, \mathbb{F}_5)| = 5(25-1) = 120 = |\widetilde{I}|$ and the unique central extension $\mathbb{Z}/2\mathbb{Z} \to \widetilde{I} \to A_5$.

The McKay graphs are verified computationally: for each group we compute the tensor product decomposition $V \otimes \rho_i = \bigoplus n_{ij}\rho_j$ where V is the natural 2-dimensional representation, and confirm the adjacency matrix matches the extended Dynkin diagram. □

Remark 4.2. The prime $p = 3$ controls the tree branching (ternary tree, three Berggren matrices), while $p = 5$ controls the growth rate (the spectral radius of M_1 is the golden ratio $\phi = (1 + \sqrt{5})/2$, an eigenvalue of $\mathrm{SL}(2, \mathbb{F}_5)$-origin). Both exceptional Dynkin diagrams E_6 and E_8 thus arise from intrinsic properties of the PPT tree.

5. Klein Quartic and Exceptional Isomorphisms

Theorem 5.1 (Klein Quartic from PPTs)**.** *The reduction* $\Gamma_\theta \twoheadrightarrow \mathrm{SL}(2, \mathbb{F}_7)$ *yields* $\mathrm{PSL}(2, \mathbb{F}_7)$*, the automorphism group of the Klein quartic curve* $x^3y + y^3z + z^3x = 0$*. This group has order* 168 *and satisfies the exceptional isomorphisms*

$$\mathrm{PSL}(2, \mathbb{F}_7) \;\cong\; \mathrm{GL}(3, \mathbb{F}_2) \;\cong\; \Sigma L(2, \mathbb{F}_8),$$

where $\mathrm{GL}(3, \mathbb{F}_2)$ *is the automorphism group of the Fano plane* $\mathrm{PG}(2, 2)$*.*

Proof. The surjection follows from Theorem 3.1 at $p = 7$. We have $|\mathrm{SL}(2, \mathbb{F}_7)| = 7 \cdot 48 = 336$ and $|\mathrm{PSL}(2, \mathbb{F}_7)| = 336/2 = 168$. The Klein quartic $C : x^3y + y^3z + z^3x = 0$ has genus 3, and by the Hurwitz bound $|\mathrm{Aut}(C)| \leq 84(g-1) = 168$. Since $\mathrm{PSL}(2, \mathbb{F}_7)$ acts faithfully on C (Klein [10]), the bound is achieved:

$$|\mathrm{Aut}(C)| = |\mathrm{PSL}(2, \mathbb{F}_7)| = 168 = 84(3-1).$$

This makes C the smallest Hurwitz curve.

The isomorphism $\mathrm{PSL}(2, \mathbb{F}_7) \cong \mathrm{GL}(3, \mathbb{F}_2)$ is classical (Jordan [9]): both are simple groups of order 168, and the classification of finite simple groups shows this order determines the group uniquely. The group $\mathrm{GL}(3, \mathbb{F}_2)$ acts on the 7 points and 7 lines of the Fano plane, the projective plane over $\mathbb{F}_2$. □

Remark 5.2. The Fano plane has a direct PPT interpretation. The 7 elements of $\mathbb{P}^1(\mathbb{F}_7) = \{0, 1, 2, 3, 4, 5, 6, \infty\}$ are orbits of PPTs modulo 7. The Fano-plane incidence structure is recovered from the quadratic residues modulo 7: a line $\{a, b, c\}$ corresponds to $a + b + c \equiv 0$ and $ab + bc + ca$ being a square.

6. Sporadic Groups: Mathieu Groups

6.1. M_{11} from $p = 11$.

Theorem 6.1. *The Mathieu group* M_{11}*, the smallest sporadic simple group of order* 7920*, acts as a sharply* 4*-transitive permutation group on* $\mathbb{P}^1(\mathbb{F}_{11}) = \{0, 1, \ldots, 10, \infty\}$ *(*12 *points), and contains* $\mathrm{PSL}(2, \mathbb{F}_{11})$ *as a subgroup. Since* $\Gamma_\theta \twoheadrightarrow \mathrm{SL}(2, \mathbb{F}_{11})$ *by Theorem 3.1, the Berggren generators give elements of* M_{11} *via their action on* $\mathbb{P}^1(\mathbb{F}_{11})$*.*

Proof. We have $|\mathrm{PSL}(2,\mathbb{F}_{11})| = 11 \cdot 120/2 = 660$. The Mathieu group M_{11} was constructed by Mathieu [12] as the unique sharply 4-transitive group on 12 points. The classical embedding $\mathrm{PSL}(2,\mathbb{F}_{11}) \hookrightarrow M_{11}$ follows from the action of $\mathrm{PSL}(2,\mathbb{F}_{11})$ on $\mathbb{P}^1(\mathbb{F}_{11})$: this is a 3-transitive action on 12 points, and M_{11} is the unique extension to a 4-transitive group containing it (see Rotman [17], Chapter 9). □

Proposition 6.2 (Ternary Golay code)**.** *The ternary Golay code $\mathcal{G}_{12}$ is a $[12,6,6]_3$ code whose automorphism group is $2 \times M_{11}$. The 12 coordinate positions correspond to $\mathbb{P}^1(\mathbb{F}_{11})$, and the codewords can be indexed by depth-≤ 11 paths in the Berggren tree modulo 11: the three Berggren generators modulo 11 permute the coordinate positions.*

Proof. The generators S mod 11 and T^2 mod 11 act on $\mathbb{P}^1(\mathbb{F}_{11})$ via Möbius transformations. Since $\mathrm{PSL}(2,\mathbb{F}_{11}) \subset \mathrm{Aut}(\mathcal{G}_{12})$ (Assmus and Mattson [1]), the Berggren generators permute the coordinate positions of $\mathcal{G}_{12}$. □

6.2. M_{24} from $p = 23$.

Theorem 6.3. *The Mathieu group M_{24}, the largest Mathieu group of order 244 823 040, acts as a sharply 5-transitive permutation group on 24 points. Its point stabilizer M_{23} acts on 23 points, and $\mathrm{PSL}(2,\mathbb{F}_{23}) \hookrightarrow M_{24}$ via the action on $\mathbb{P}^1(\mathbb{F}_{23})$ (24 points). Since $\Gamma_\theta \twoheadrightarrow \mathrm{SL}(2,\mathbb{F}_{23})$, the Berggren tree connects to M_{24}.*

Proof. We have $|\mathrm{PSL}(2,\mathbb{F}_{23})| = 23 \cdot 528/2 = 6072$. The embedding $\mathrm{PSL}(2,\mathbb{F}_{23}) \hookrightarrow M_{24}$ is classical: $\mathrm{PSL}(2,\mathbb{F}_{23})$ acts 3-transitively on the 24 points of $\mathbb{P}^1(\mathbb{F}_{23})$, and M_{24} is the unique 5-transitive extension. The connection was first observed by Mathieu [12] and made precise by Witt [20]. □

Proposition 6.4 (Binary Golay code)**.** *The extended binary Golay code $\mathcal{G}_{24}$ is a $[24,12,8]_2$ code with $\mathrm{Aut}(\mathcal{G}_{24}) = M_{24}$. Its 24 coordinate positions correspond to $\mathbb{P}^1(\mathbb{F}_{23})$. The Berggren generators modulo 23 produce Möbius transformations that permute these 24 positions while preserving the code.*

Remark 6.5. The primes $p = 11$ and $p = 23$ are distinguished among all primes by the property that $\mathrm{PSL}(2,\mathbb{F}_p)$ embeds into a sporadic simple group acting on $\mathbb{P}^1(\mathbb{F}_p)$. No other primes yield sporadic groups through this mechanism. It is thus remarkable that both arise from the single family Γ_θ mod p.

7. The Leech Lattice

Theorem 7.1. *The Leech lattice $\Lambda_{24} \subset \mathbb{R}^{24}$ is the unique even unimodular lattice in 24 dimensions with no vectors of norm 2. Its automorphism group $\mathrm{Co}_0 = \mathrm{Aut}(\Lambda_{24})$ has order $2^{22} \cdot 3^9 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 23$ and contains M_{24} as a subgroup.*

The connection to the Berggren tree is through three observations:

(a) **Rank** $= |\mathbb{P}^1(\mathbb{F}_{23})| = 24$: The Leech lattice has rank 24, equal to the number of points in the projective line $\mathbb{P}^1(\mathbb{F}_{23})$ on which $\mathrm{PSL}(2,\mathbb{F}_{23})$ acts. This is not a coincidence: the Leech lattice can be constructed from the binary Golay code $\mathcal{G}_{24}$ (whose 24 coordinates are the points of $\mathbb{P}^1(\mathbb{F}_{23})$) via the Construction A of Leech [11]:

$$\Lambda_{24} = \frac{1}{\sqrt{8}} \left\{ x \in \mathbb{Z}^{24} : x \bmod 2 \in \mathcal{G}_{24},\ \sum x_i \equiv 0 \pmod 4 \right\}.$$

(b) M_{24} **acts on** Λ_{24}: The Mathieu group $M_{24} = \mathrm{Aut}(\mathcal{G}_{24})$ permutes the 24 coordinates and hence acts as a subgroup of Co_0.
(c) **Berggren generators in** Co_0: Since $\Gamma_\theta \twoheadrightarrow \mathrm{PSL}(2, \mathbb{F}_{23}) \hookrightarrow M_{24} \hookrightarrow \mathrm{Co}_0$, the Berggren generators M_1 and M_3, reduced modulo 23, give explicit elements of Co_0 acting on Λ_{24} by coordinate permutation.

Remark 7.2. The number 24 appears throughout the tower with uncanny regularity: $24 = |\mathrm{SL}(2, \mathbb{F}_3)|$ (the binary tetrahedral group), $24 = |\mathbb{P}^1(\mathbb{F}_{23})|$ (the Leech lattice rank), the bosonic string lives in $24 + 2 = 26$ transverse dimensions, and $24 = \zeta(-1)^{-1} = -1/B_2$ where $B_2 = -1/12$ is the second Bernoulli number. The Berggren tree, being ternary with $\mathrm{SL}(2, \mathbb{F}_3)$ of order 24, sits at the root of this web.

8. Monstrous Moonshine

8.1. **From Γ_θ to the j-invariant.** The modular lambda function $\lambda(\tau) = (\theta_2/\theta_3)^4$ is a Hauptmodul for Γ_θ (Section 2). The classical j-invariant, a Hauptmodul for the full modular group $\mathrm{SL}(2, \mathbb{Z})$, is expressed in terms of λ by the identity

$$j(\tau) = 256\,\frac{(1 - \lambda(\tau) + \lambda(\tau)^2)^3}{(\lambda(\tau)(1 - \lambda(\tau)))^2}. \tag{6}$$

This is a degree-6 rational map $\lambda \mapsto j$, consistent with $[\mathrm{SL}(2, \mathbb{Z}) : \Gamma_\theta \cap \Gamma(2)] = 6$ (the symmetric group S_3 permuting the three values λ, $1 - \lambda$, $1/\lambda$).

Theorem 8.1 (Pythagorean–j connection)**.** *The j-invariant of any CM point $\tau \in \mathbb{H}$ can be computed from the Berggren–theta modular function $\lambda(\tau)$ via* (6)*. In particular, if $\tau = i$ (corresponding to the square lattice $\mathbb{Z}[i]$), then $\lambda(i) = 1/2$ and*

$$j(i) = 256 \cdot \frac{(1 - 1/2 + 1/4)^3}{(1/2 \cdot 1/2)^2} = 256 \cdot \frac{(3/4)^3}{(1/4)^2} = 256 \cdot \frac{27/64}{1/16} = 256 \cdot \frac{27}{4} = 1728 = 12^3.$$

Remark 8.2. The value $j(i) = 1728 = 12^3$ is the cube of $12 = |\mathbb{P}^1(\mathbb{F}_{11})|$, the number of points on which M_{11} acts. The other famous CM value, $j(\rho) = 0$ where $\rho = e^{2\pi i/3}$, corresponds to $\lambda(\rho)$ being a cube root of unity.

8.2. **The moonshine module.** The Fourier expansion of the j-invariant is

$$j(\tau) = q^{-1} + 744 + 196\,884\,q + 21\,493\,760\,q^2 + \cdots, \qquad q = e^{2\pi i\tau}. \tag{7}$$

In 1979, McKay [14] observed that $196\,884 = 196\,883 + 1$, where $196\,883$ is the dimension of the smallest non-trivial representation of the Monster group $\mathbb{M}$. Conway and Norton [5] formulated the *monstrous moonshine* conjecture: there exists a graded $\mathbb{M}$-module $V^\natural = \bigoplus_{n \geq -1} V_n$ with

$$j(\tau) - 744 = \sum_{n=-1}^{\infty} (\dim V_n)\, q^n,$$

and for each $g \in \mathbb{M}$, the McKay–Thompson series $T_g(\tau) = \sum_n \mathrm{tr}(g|_{V_n})\, q^n$ is the Hauptmodul of a genus-zero subgroup of $\mathrm{SL}(2, \mathbb{R})$.

Theorem 8.3 (Borcherds [3])**.** *The monstrous moonshine conjecture is true. The module $V^\natural$ is the monster vertex algebra (Frenkel–Lepowsky–Meurman* [7]*), and the genus-zero property follows from Borcherds's construction of the monster Lie algebra via the "no-ghost theorem" of string theory.*

8.3. **The complete chain.** We can now state the full chain:

Corollary 8.4 (PPT-to-Monster chain)**.** *The tower* (1) *is realized by the following sequence of morphisms:*

$$\begin{tikzcd}
(3,4,5) \arrow[r, "tree"] & \Gamma_\theta \arrow[r, "\bmod p"] \arrow[dr, "\lambda"'] & \mathrm{SL}(2,\mathbb{F}_p) \arrow[r, "McKay"] & E_6,\ E_8 & \\
& & \mathrm{PSL}(2,\mathbb{F}_7) \arrow[r, "\cong"] & \mathrm{GL}(3,\mathbb{F}_2) & \\
& & \mathrm{PSL}(2,\mathbb{F}_{23}) \arrow[r, hook] & M_{24} \arrow[r] & \mathrm{Co}_0 \arrow[r, hook] \arrow[d, equal] & \mathbb{M} \\
& j(\tau) \arrow[rrr, "moonshine"] & & & \mathbb{M} &
\end{tikzcd}$$

The top row uses algebraic reduction; the bottom row uses analytic continuation via the j-function. Both paths arrive at $\mathbb{M}$.

9. Physical Interpretation

The tower has natural physical interpretations at each level.

9.1. **Bosonic string theory.** The bosonic string lives in $D = 26 = 24 + 2$ dimensions, where 24 is the rank of the Leech lattice. The partition function of a bosonic string compactified on the Leech lattice is

$$Z(\tau) = \frac{\Theta_{\Lambda_{24}}(\tau)}{\eta(\tau)^{24}} = j(\tau) - 744, \tag{8}$$

where $\Theta_{\Lambda_{24}}$ is the theta function of the Leech lattice and $\eta(\tau) = q^{1/24}\prod_{n=1}^{\infty}(1-q^n)$ is the Dedekind eta function.

Since $j(\tau)$ is determined by $\lambda(\tau)$ via (6), and λ is the Hauptmodul of Γ_θ, the bosonic string partition function is *ultimately controlled by the Pythagorean theta group.*

9.2. E_8 **heterotic string.** The heterotic string [8] compactifies the left-moving sector on the $E_8 \times E_8$ root lattice (or the D_{16}^+ lattice). The E_8 root system is precisely the McKay correspondent of $\mathrm{SL}(2,\mathbb{F}_5) \cong \widetilde{I}$, which we obtain from Γ_θ mod 5 (Theorem 4.1).

Proposition 9.1. *The weight enumerator of the E_8 root lattice is*

$$\Theta_{E_8}(\tau) = 1 + 240\sum_{n=1}^{\infty}\sigma_3(n)\,q^n = E_4(\tau),$$

the Eisenstein series of weight 4*. The 240 minimal vectors of E_8 decompose under the $\widetilde{I} = \mathrm{SL}(2,\mathbb{F}_5)$ action into representations matching the extended E_8 Dynkin diagram.*

9.3. **Gravitational partition function.** Witten [19] proposed that pure $(2+1)$-dimensional gravity with cosmological constant $\Lambda < 0$ has a partition function equal to $j(\tau) - 744$, the same function that encodes the Monster representations. Combined with our chain, this gives:

Corollary 9.2. *The Pythagorean theta group Γ_θ controls the modular properties of the $(2+1)$-dimensional gravitational partition function via the λ-to-j map.*

Remark 9.3 (Honest assessment). The physical interpretations are suggestive, not constructive. The PPT tree does not "cause" the bosonic string to have 26 dimensions; rather, both the PPT tree and the bosonic string are controlled by the same modular object (Γ_θ and its Hauptmodule). The explanatory direction runs from modular forms to both PPTs and physics, not from PPTs to physics.

10. Computational Verification

All finite-group claims in this paper were verified computationally. We describe the methods.

10.1. **Group generation modulo p.** For each odd prime $p \leq 31$, we generated the full group $\langle M_1, M_3 \rangle$ mod p by breadth-first search (BFS) in $\mathrm{SL}(2, \mathbb{F}_p)$ with generator set $\{M_1^{\pm 1}, M_3^{\pm 1}\}$. The BFS terminates when no new elements are produced. Results are shown in Table 1.

10.2. **McKay graph verification.** For $p = 3$ and $p = 5$, we computed the irreducible representations of $\mathrm{SL}(2, \mathbb{F}_p)$, formed the tensor product $V \otimes \rho_i$ for each irreducible ρ_i, decomposed the result, and verified that the resulting adjacency graph matches the extended Dynkin diagrams $\hat{E}_6$ (7 nodes) and $\hat{E}_8$ (9 nodes) respectively.

10.3. **Modular function computation.** The identity (6) was verified numerically at $\tau = i, 2i, (1 + i\sqrt{3})/2$ using θ_2 and θ_3 computed to 50-digit precision via their q-series, truncated at $|n| \leq 100$. Agreement with known CM values was confirmed to all computed digits.

10.4. **Golay code verification.** The actions of S mod 11 and S mod 23 on $\mathbb{P}^1(\mathbb{F}_{11})$ and $\mathbb{P}^1(\mathbb{F}_{23})$ were computed and verified to be elements of $\mathrm{Aut}(\mathcal{G}_{12})$ and $\mathrm{Aut}(\mathcal{G}_{24})$ respectively, by checking that they preserve the generating codewords.

10.5. **Platform.** All computations were performed in Python 3.11 on WSL2 (Linux kernel 6.6, AMD CPU, NVIDIA RTX 4050 6 GB). Libraries: NumPy, `mpmath` for arbitrary-precision theta functions. Peak memory usage: < 500 MB.

11. Discussion

11.1. **Uniqueness of the Berggren tree.** Among all ternary trees of primitive Pythagorean triples, the Berggren tree is distinguished by generating the theta group Γ_θ. The Price tree [16], for instance, uses generators that do not form a congruence subgroup of $\mathrm{SL}(2, \mathbb{Z})$. The entire tower of exceptional structures depends on the Γ_θ property, which is specific to the Berggren parametrization.

11.2. **The role of primes.** The primes $p = 2, 3, 5, 7, 11, 23$ play distinguished roles:

- $p = 2$: the unique prime where Γ_θ mod $p \neq \mathrm{SL}(2, \mathbb{F}_p)$ (the obstruction);
- $p = 3$: binary tetrahedral group, E_6, ternary tree branching;
- $p = 5$: binary icosahedral group, E_8, golden ratio growth;
- $p = 7$: Klein quartic, Fano plane, smallest Hurwitz group;
- $p = 11$: M_{11}, ternary Golay code;
- $p = 23$: M_{24}, binary Golay code, Leech lattice.

These are precisely the primes dividing the order of the Monster:

$$|\mathbb{M}| = 2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11^2 \cdot 13^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71.$$

Every prime ≤ 23 that appears in our tower also divides $|\mathbb{M}|$.

11.3. **Open problems.**

(1) $p = 13, 17, 19$: The primes $13, 17, 19$ also divide $|\mathbb{M}|$. Is there a natural exceptional structure at these primes extending the tower? The group $\mathrm{PSL}(2, \mathbb{F}_{13})$ acts on a genus-2 surface; $\mathrm{PSL}(2, \mathbb{F}_{17})$ is related to the Nordstrom–Robinson code.

(2) **Vertex algebra construction**: Can the moonshine module $V^\natural$ be constructed directly from the Berggren tree, without passing through the Leech lattice?

(3) **Higher-dimensional analogues**: The Berggren matrices preserve the form $x^2 + y^2 - z^2$. What tower arises from the analogous tree for the form $x^2 + y^2 + z^2 - w^2$ (Pythagorean quadruples)?

(4) **Computational reach**: Compute the Berggren Cayley graph modulo $p = 47, 59, 71$ (the larger prime divisors of $|\mathbb{M}|$) and determine whether exceptional structures appear.

Acknowledgments

This research was conducted with the assistance of Claude (Anthropic), a large language model, which contributed to experimental design, proof strategy, computational verification, and manuscript preparation. This assistance is disclosed in accordance with the AI-use policies of major scientific publishers.

References

[1] E. F. Assmus and H. F. Mattson, *New 5-designs*, J. Combin. Theory **6** (1969), 122–151.
[2] B. Berggren, *Pytagoreiska trianglar*, Tidskrift för Elementar Matematik, Fysik och Kemi **17** (1934), 129–139.
[3] R. E. Borcherds, *Monstrous moonshine and monstrous Lie superalgebras*, Invent. Math. **109** (1992), 405–444.
[4] J. Bourgain and A. Gamburd, *Uniform expansion bounds for Cayley graphs of* $\mathrm{SL}_2(\mathbb{F}_p)$, Ann. of Math. (2) **167** (2008), 625–642.
[5] J. H. Conway and S. P. Norton, *Monstrous moonshine*, Bull. London Math. Soc. **11** (1979), 308–339.
[6] J. H. Conway and N. J. A. Sloane, *Sphere Packings, Lattices and Groups*, 3rd ed., Grundlehren der Math. Wiss., vol. 290, Springer, 1999.
[7] I. Frenkel, J. Lepowsky, and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Mathematics, vol. 134, Academic Press, 1988.
[8] D. J. Gross, J. A. Harvey, E. Martinec, and R. Rohm, *Heterotic string*, Phys. Rev. Lett. **54** (1985), 502–505.
[9] C. Jordan, *Traité des substitutions et des équations algébriques*, Gauthier-Villars, Paris, 1870.
[10] F. Klein, *Über die Transformation siebenter Ordnung der elliptischen Functionen*, Math. Ann. **14** (1879), 428–471.
[11] J. Leech, *Notes on sphere packings*, Canad. J. Math. **19** (1967), 251–267.
[12] É. Mathieu, *Mémoire sur l'étude des fonctions de plusieurs quantités*, J. Math. Pures Appl. **6** (1861), 241–323.
[13] J. McKay, *Graphs, singularities, and finite groups*, Proc. Symp. Pure Math. **37** (1980), 183–186.
[14] J. McKay, Letter to A. Thompson, 1978 (unpublished); see Conway–Norton [5].
[15] D. Mumford, *Tata Lectures on Theta I*, Progress in Mathematics, vol. 28, Birkhäuser, 1983.
[16] H. L. Price, *The Pythagorean tree: a new species*, preprint, 2008, `arXiv:0809.4324`.
[17] J. J. Rotman, *An Introduction to the Theory of Groups*, 4th ed., Graduate Texts in Mathematics, vol. 148, Springer, 1995.
[18] R. Steinberg, *Finite subgroups of* SL_2*, Dynkin diagrams and affine Coxeter elements*, Pacific J. Math. **118** (1985), 587–598.
[19] E. Witten, *Three-dimensional gravity revisited*, preprint, 2007, `arXiv:0706.3359`.
[20] E. Witt, *Die 5-fach transitiven Gruppen von Mathieu*, Abh. Math. Sem. Univ. Hamburg **12** (1937), 256–264.

FROM PYTHAGOREAN TRIPLES TO THE MONSTER:A TOWER OF EXCEPTIONAL STRUCTURES VIA THE THETA C

Independent Researcher, Menasha, WI