

The Murty–Simon bound at order 28

A computer-assisted candidate proof

Paul Lenz*

9 September 2026

Fan-free reviewer edition 2; independent mathematical review open

Abstract

We present a computer-assisted candidate proof that a simple diameter-two edge-critical graph on 28 vertices has at most 196 edges, with equality precisely for $K_{14,14}$. A complement-based selection of quasi-edges gives residual degree, demand, and source-supplement constraints. Two separate finite calculations handle 196 edges and 197 edges; the latter uses a newly generated domain, not a transfer of equality-level certificates. Exact integer linear-combination certificates and complete branching proofs exclude the retained necessary-condition systems. Fan’s published strict estimate is retained for historical attribution, but edition 2 replaces it logically by a direct Fan-free exclusion of every edge count above 197. This edition consolidates the direct LocalIncidence-v7/direct197-v8 route and identifies its complete computational dependencies. An internal red-team audit found no blocking defect in that route. Independent mathematical review, external reproduction, and formal verification remain open. No novelty or general-order resolution is claimed.

Reading this edition. Sections 1–5 establish the structural and finite-domain framework; Sections 6–8 explain the stronger numerical systems and exact certificates; Section 9 assembles all degree cases; Section 10 records reproducibility and review limits. The accompanying *Computational verification and reviewer guide* specifies immutable evidence, executable entry points, the hardened checking release, and outstanding review questions. The full finite systems are specified by the pinned source constructors and certificates, not by the displayed summary counts alone. Original historical notes and archives are preserved unchanged.

1 Statement, conventions and external input

A graph is simple and finite. Distance between disconnected vertices is infinite. A graph G is diameter-two edge-critical when $\text{diam}(G) = 2$ and $\text{diam}(G - e) > 2$ for every edge e . Write $n = |V(G)|$, $m = e(G)$ and $b = \Delta(G)$. The Murty–Simon conjecture asserts $m \leq \lfloor n^2/4 \rfloor$, with equality only for the balanced complete bipartite graph [2, 1].

Theorem 1.1 (Order-28 candidate). *Every simple diameter-two edge-critical graph on 28 vertices has at most 196 edges. Equality holds if and only if the graph is $K_{14,14}$.*

The proof below uses the verified computational propositions in Section 8. The word *candidate* records the review status of the mathematical derivations and graph-to-system implication; it does not turn numerical solver messages into proofs.

*Project direction: Paul Lenz. Mathematical development, software, manuscript drafting and internal checking: ChatGPT (“Geeps”). This attribution describes the work performed; it does not assert independent human verification.

Historically, reviewer edition 1 used Fan’s strict estimate, as reported by Wang [2, p. 2], to deduce $m \leq 197$. Fan remains cited for attribution, but his theorem is *not* a logical dependency of edition 2. The replacement is the project note [project/research/fan-free-fixed-orders/2026-09-09-v](#) the existing 197-edge calculation remains unchanged, while every $m \geq 198$ is excluded directly. Degree sum handles $\Delta \leq 14$; fresh exact arithmetic closes $\Delta = 15$ for $m = 198, \dots, 210$; the pointwise charging bound at $\Delta = 16$ and residual h-index bounds at $\Delta = 17, \dots, 26$ only strengthen as m increases; and $\Delta = 27$ is the star case. The historical Fan-based reviewer-v1 manuscript remains preserved unchanged.

A bipartite diameter-two graph is complete bipartite: an absent cross-part edge would have odd distance at least three. Its edge count is at most 196, with equality only for $K_{14,14}$. A universal vertex forces an edge-critical graph to be a star, since an edge between other vertices would be redundant. That case has only 27 edges. These exceptions are handled explicitly.

2 Complement selection and residual identities

Put $H = \overline{G}$. An adjacent pair totally dominates a graph when every vertex has a neighbour in that pair. The complement of any graph has diameter greater than two exactly when the graph has such an adjacent pair: the endpoints are nonadjacent in the complement and have no common neighbour there. This direct observation, also recorded in the complement literature [2, 3], suffices for the quasi-edge argument; no bound on $\text{diam}(H)$ is assumed.

Lemma 2.1 (Quasi-edge with a prescribed exceptional endpoint). *Suppose uw is a nonedge of H and u, w both miss a third vertex. There is an existing edge ui , or symmetrically wi , whose endpoints totally dominate every vertex of H except the opposite endpoint. Write $ui \rightarrow w$ in the former case. The exception of that existing edge is unique.*

Proof. Deleting uw from G makes some distance exceed two. Hence $H + uw$ has an adjacent total dominating pair, while H has none. The pair cannot be $\{u, w\}$ because it still misses the third vertex. It must use an endpoint of the newly added edge; otherwise its domination is unchanged. Its edge already existed. The only newly dominated vertex is the opposite endpoint of uw , which is therefore its unique exception in H . $\square$

Choose v of minimum H -degree and define

$$a = n - 1 - b, \quad A = N_H(v), \quad B = V(H) \setminus N_H[v], \quad C = H[A], \quad F = \overline{C}.$$

Thus $|A| = a$, $|B| = b$, and $d_i = d_F(i)$ for $i \in A$. Every missing unordered B -pair misses v . Choose exactly one edge $ui \rightarrow w$ for that pair. Its auxiliary i is in A because it must dominate v . Call these cross-edges *selected*; all other edges of $H[A, B]$ are *residual*. The missing pair is oriented $u \rightarrow w$; u is its source and w its supplement.

Distinct missing B -pairs select distinct cross-edges: the B -endpoint and unique exception recover the pair. At a fixed source, labels and supplements are distinct. Each unordered B -pair supports at most one selected arc, not one in each orientation.

The notation used throughout is

Quantity	Meaning
ρ_u, R_i, r	Residual degrees at B , at A , and their common total
q_u, p_u, σ_u	Selected outdegree, indegree, and $q_u + p_u$
x_i, Q	Actual selected degree at label i ; total selected incidences
t	Surplus $m - b(n - b)$
ℓ	$b - a - 1 = 2b - n$
s_i, S	Minimum demand $\max(0, d_i - R_i)$; its sum

Selected cross-edges and existing $H[B]$ -edges together number $\binom{b}{2}$. Counting all edges of H gives

$$\begin{aligned} e(C) + r &= \binom{a}{2} - t, & e(F) &= r + t, \\ \sum_i d_i &= 2(r + t), & \sum_i R_i &= \sum_u \rho_u = r. \end{aligned} \tag{1}$$

Minimum H -degree gives $d_H(i) = a - d_i + R_i + x_i \geq a$ and $d_H(u) = b - 1 + \rho_u - p_u \geq a$. Therefore

$$\begin{aligned} x_i &\geq s_i, & S &\geq r + 2t, & q_u + \rho_u &\leq a, \\ p_u &\leq \rho_u + \ell, & q_u + p_u &\leq b - 1, & Q &\leq r + \ell b. \end{aligned} \tag{2}$$

Actual selected degree is not minimum demand: x_i may exceed s_i , including when $s_i = 0$. All finite tests retain that possibility.

3 Structural injections

Lemma 3.1 (Selected-edge constraints). *For every selected $ui \rightarrow w$,*

$$\begin{aligned} d_i &\leq \rho_u + R_i, & d_i &\leq \rho_u + \rho_w, \\ d_i &\leq \rho_u + q_u - 1, & \rho_w + q_w &\geq q_u - 1, \\ R_i + x_i &\geq q_u + p_u. \end{aligned} \tag{3}$$

In particular $s_i \leq \rho_u$ at every selected incidence.

Proof. Every F -neighbour j of i must neighbour u . At most ρ_u of the edges uj are residual. A remaining uj is selected and has a distinct supplement $w_j \neq w$. Since ui must dominate w_j and uw_j is absent, iw_j exists. It is residual: i, w_j both miss the A -vertex j , while a selected edge's exception is in B . Distinct supplements give $d_i - \rho_u \leq R_i$.

For the pair bound, each of those selected uj must dominate w , forcing ju . This edge is residual because j, w both miss i . Distinct j give $d_i - \rho_u \leq \rho_w$. The source neighbours i and all its F -neighbours, giving the third inequality. Every other selected label at u neighbours w , giving the fourth.

Finally, i neighbours u and every missing- B -pair neighbour of u except w . These are $q_u + p_u$ distinct vertices, so $d_H(i, B) = R_i + x_i \geq q_u + p_u$. $\square$

Lemma 3.2 (Residual activity). *If $t > 0$, then every $u \in B$ has $\rho_u \geq 1$.*

Proof. Suppose $\rho_u = 0$ and set $U = N_H(u) \cap A$, $T = A \setminus U$. Every ui with $i \in U$ is selected. No F -edge joins U and T , since such an edge would give a selected quasi-edge an A -exception as well as its B -exception.

Let w_i be the distinct supplements at u . An F -edge ij within U forces ju and iw_j , by domination of the other selected edge's supplement. Both are residual. Their A -endpoints and supplement labels distinguish all $2e_F(U)$ such edges.

An F -edge within T misses u . Its quasi-edge auxiliary must dominate u . It cannot be u or v ; if in A it lies in U and is adjacent in C to both exceptional-pair endpoints, impossible. It therefore lies in B and gives a residual cross-edge with A -endpoint in T . The A -endpoint and unique exception identify the original missing pair, so these $e_F(T)$ edges are distinct. The two families are disjoint. Thus $r \geq 2e_F(U) + e_F(T) \geq e(F) = r + t$, a contradiction. Empty sets cause no exception. $\square$

Lemma 3.3 (No isolated C -vertex in the degree-15 scopes). *For $(a, b, t) = (12, 15, 1)$ or $(12, 15, 2)$, $\delta(C) \geq 1$. Consequently $d_i \leq 10$ and $r \leq 59$ or 58 , respectively.*

Proof. If x is isolated in C , put $X = A \setminus \{x\}$. Every missing pair inside X misses x ; its quasi-edge auxiliary lies in B . This gives a family P of distinct residual edges, with $|P| = \binom{a-1}{2} - e(C)$. Every B -endpoint u used by P must neighbour x to dominate it. The edge xu is residual: x, u both miss the original A -exception. It lies outside P . Each unused B -vertex supplies another residual edge by Lemma 3.2. Distinct B -endpoints distinguish these b extra edges. Hence

$$b \leq r - |P| = \binom{a}{2} - t - \binom{a-1}{2} = a - 1 - t,$$

requiring $15 \leq 10$ or $15 \leq 9$. Finally $e(C) \geq \lceil a/2 \rceil = 6$ and use (1). $\square$

Lemma 3.4 (Source-local restrictions). *Fix $u \in B$ and partition A into selected neighbours S_u , residual neighbours T_u , and missing neighbours M_u . Then $F(S_u, M_u)$ is empty, and*

$$d_j \leq a - q_u - 1 \quad (j \in M_u), \quad (4)$$

$$R_i \geq |N_F(i) \cap S_u| \quad (i \in S_u), \quad (5)$$

$$\rho_u + 2e_F(S_u) + e_F(M_u) - e_C(T_u, M_u) \leq r. \quad (6)$$

Proof. Every selected edge at u must dominate each $j \in M_u$ through its label. Thus j has at least q_u neighbours in C , proving (4) and the absence of $F(S_u, M_u)$. For (5), a selected w_i must dominate the distinct supplements w_j of selected uj for $j \in N_F(i) \cap S_u$. The forced iw_j are residual and distinct.

For (6), internal F -edges in S_u force $2e_F(S_u)$ residual edges with A -endpoints in S_u , as above. A missing pair in M_u misses u . Its quasi-edge auxiliary lies in T_u or B : it cannot be in S_u , because every S_u – M_u pair is a C -edge. At most $e_C(T_u, M_u)$ missing pairs use auxiliaries in T_u , since those quasi-edges are distinct. Every remaining pair gives a distinct residual cross-edge with an A -exception and A -endpoint in M_u . Finally the ρ_u residual edges at u have endpoints in T_u . The three endpoint classes are disjoint, proving the inequality (in fact with the M_u contribution replaced by its positive part). $\square$

The last injection uses full edge-criticality for missing pairs in A . It is applied here to the original graph, not to a weakened object retaining only selected B -quasi-edges.

Using $|T_u| = \rho_u$, $|S_u| = q_u$, $|M_u| = a - q_u - \rho_u$ and (1), (6) gives

$$\sum_{i \in S_u} d_i \leq \rho_u(2a - \rho_u - 3 + q_u) - 2t, \quad (7)$$

$$\sum_{i \in S_u} R_i \leq r - \rho_u. \quad (8)$$

Indeed $t + \rho_u + e_F(S_u) \leq e_F(T_u) + e_F(S_u, T_u) + \rho_u|M_u|$; multiply by two and add $e_F(S_u, T_u)$, then bound $2e_F(T_u) \leq \rho_u(\rho_u - 1)$ and $e_F(S_u, T_u) \leq q_u\rho_u$. The source's residual edges have no endpoints in S_u , proving (8).

4 Complete demand and residual domains

A positive demand needs a selected source, so Lemma 3.1 and $q_u + \rho_u \leq a$ give $0 \leq s_i \leq a - 1$. Put

$$g_a(s) = \frac{s(s-1)}{a-s}, \quad f_a(s) = s - g_a(s) = \frac{s(a+1-2s)}{a-s}.$$

Lemma 4.1 (Charging). *At positive surplus,*

$$r - b \geq \sum_i g_a(s_i), \quad \sum_i f_a(s_i) \geq b + 2t. \quad (9)$$

Proof. Choose exactly s_i actual selected incidences at label i . At a source with residual degree $\rho < a$, charge $(\rho - 1)/(a - \rho)$ per chosen incidence. At most $a - \rho$ incidences are present, so their total charge is at most $\rho - 1$. A source with $\rho = a$ has no selections. The charged function is increasing on $[1, a)$, and every selected demand has $s_i \leq \rho$, so the total charge at label i is at least $g_a(s_i)$. Sum and apply $S \geq r + 2t$. $\square$

Every possible graph therefore gives a nondecreasing length- a tuple in $\{0, \dots, a - 1\}$ obeying (9). For each tuple the residual total lies in

$$r_{\min} = b + \left\lceil \sum_i g_a(s_i) \right\rceil, \quad r_{\max} = \min\{S - 2t, \binom{a}{2} - t\}. \quad (10)$$

An empty interval is impossible. Sorting demands and residual rows separately only relabels A and B . It does *not* sort the subsequently paired d_i, R_i columns independently.

The checker determines the full demand-domain size independently, verifies every listed tuple's membership and uniqueness, and checks equality of cardinalities. Residual rows are every nondecreasing list of b integers in $[1, a]$ with total in (10). A multiplicity recursion and the coefficients of

$$\prod_{j=1}^a (1 - XY^j)^{-1}$$

independently enumerate and count these multisets. Non-graphical degree sequences are retained. Thus the domain is a superset of all actual graphs.

4.1 Demand support and exact source cuts

These initial tests are inherited from the preserved demand-support-v3 notes [4]. Their formulas are included to make the early numerical reductions auditable. For $2 \leq h \leq a - 1$, put $I = \{i : s_i \geq h\}$, $k = |I|$, $D = \sum_{i \in I} s_i$, and $Z = \{u : \rho_u \geq h\}$ of size z . Activity gives

$$z \leq z_{\max} = \min\left(b, \left\lfloor \frac{r_{\max} - b}{h - 1} \right\rfloor\right).$$

If $I \neq \emptyset$, its largest demand cannot exceed $z_{\max}$. Outside Z at most $a - k$ labels can be selected, so $d_H(u, A) \leq a - k + h - 1$. Write $K = a - k + h$. A source in Z with $q_u \geq K + 1$ must therefore use only supplements in Z . If p sources are this large, their pair budget is

$$P(z, p) = \binom{z}{2} - \binom{z - p}{2}.$$

Necessarily either $p = 0$, or $a - h \geq K + 1$ and $p(K + 1) \leq P(z, p)$. The I -incidences obey

$$D \leq \max_{z, p} \{(z - p) \min(k, K, a - h) + \min(p \min(k, a - h), P(z, p))\}, \quad (11)$$

where $0 \leq z \leq z_{\max}$ and only the stated p values are admitted. Non-large sources contribute the first term; large sources are limited by both labels and distinct unordered pairs. A strict violation is an exact rejection.

For descending demands $s_{(i)}$, set $D_k = \sum_{i \leq k} s_{(i)}$ and $A_{kj} = \min(a - j, |\{i \leq k : s_{(i)} \leq j\}|)$. If n_j sources have residual degree j , then

$$D_k \leq \sum_j A_{kj} n_j, \quad \sum_j n_j = b, \quad \sum_j (j - 1) n_j = r - b.$$

Integer weights $y_k \geq 0$, $\mu \in \mathbb{Z}$, $\Lambda > 0$ satisfying $\mu + \sum_k y_k A_{kj} \leq \Lambda(j-1)$ for each j imply

$$b\mu + \sum_k y_k D_k \leq \Lambda(r_{\max} - b). \quad (12)$$

Every saved dual rejection checks these integer inequalities and a strict reverse of (12); no optimisation status is trusted.

For a residual row, start with

$$c_u = \min(a - \rho_u, b - 1, |\{i : s_i \leq \rho_u\}|).$$

Refine c_u to the largest $q \leq c_u$ having at least q other vertices w with $\rho_w + c_w \geq q - 1$, using old caps simultaneously. Every actual q_u remains possible by Lemma 3.1. Decreasing integer caps terminate. Every prefix then satisfies

$$D_k \leq \sum_u \min(c_u, |\{i \leq k : s_{(i)} \leq \rho_u\}|). \quad (13)$$

The checker reconstructs the iterations and any violated prefix. Zero-demand labels are included.

5 Columns, one-source projections and activation

An individual column option is (d_i, R_i, z_i) with

$$s_i = \max(0, d_i - R_i), \quad z_i = R_i + s_i - d_i \geq 0, \quad \sum_i R_i = r, \quad \sum_i z_i = S - r - 2t.$$

In the degree-15 cases $0 \leq d_i \leq 10$, $0 \leq R_i \leq b$, and $R_i + s_i \leq b$. For positive demand $z_i = 0$; zero-demand labels may have positive slack and selected degree. A multichoice generating product retains exactly the options participating in a collection with the two fixed sums. Each option remains paired with its label and demand.

A safely overestimated eligible-source set E_i has $|E_i| \geq s_i$. If equality holds and $s_i > 0$, all its sources are forced to select i . A supplement cannot be one of those forced sources, because it must miss i .

For each source u and trial q , the one-source projection assigns every label an option and one of the statuses selected, residual, or missing. Counts are exactly q , ρ_u , and $a - q - \rho_u$; both global column sums are fixed. Selected status requires the selected-edge inequalities, a distinct allowed supplement, and $R_i + \max(1, s_i) \leq b$. Residual status requires $R_i \geq 1$. Missing status requires (4) and $R_i + s_i \leq b - 1$. Forced selections prohibit the other statuses. Label-supplement matchings have distinct endpoints and contain all mandatory labels.

A dynamic program tests these finite choices. A value of q is discarded only when no witness exists; an option is removed only when it has no witness for at least one source at any remaining q . Each real graph continues to witness every projection, so iterative pruning is sound. Different sources may still use different witnesses: survival is not a graph-realisation claim. The separately written checker reconstructs all remaining options, caps and forced sets [6].

For fixed columns the distinct-pair injection also gives, for every integer j ,

$$\sum_{i: d_i \geq j} s_i \leq |\{\{u, w\} \subset B : \rho_u + \rho_w \geq j\}|. \quad (14)$$

Projected zero-slack and pair cuts are checked before or during refinement. Their precise finite domains are specified by the pinned checking source, including the identities above, rather than an assumption of independent column sorting.

Lemma 5.1 (Activation budget). *Let $l_u \leq q_u$ and $P_w \geq p_w$ be proved bounds. For any subset $\mathcal{T}$ of actual selected arcs,*

$$\sum_{(u,w) \in \mathcal{T}} \frac{[l_u - 1 - \rho_w - l_w]_+}{P_w} \leq Q - \sum_w l_w. \quad (15)$$

A term with $P_w = 0$ cannot occur.

Proof. For each selected $u \rightarrow w$, Lemma 3.1 gives $q_w - l_w \geq [l_u - 1 - \rho_w - l_w]_+$. At most P_w terms enter w , and each is at most $q_w - l_w$. Division by P_w prevents charging the same forced increase once for every incoming arc. Sum over w . $\square$

Choose exactly s_i actual incidences at each label. Their possible placement is relaxed to a capacitated network through label, source–label, ordered-pair and supplement nodes. Capacities enforce label demands, distinct source–label and ordered-pair incidences, and incoming caps. Some opposite-orientation and outgoing-total constraints are intentionally omitted; that enlarges the feasible set and cannot cause a false rejection. Costs are the terms of (15), scaled to integers; the available budget is a proved upper bound on $Q - \sum l$.

A cut below total demand rejects the flow. Alternatively integer vertex potentials π give the exact cost lower bound

$$S(\pi_{\text{sink}} - \pi_{\text{source}}) - \sum_{e=(u,w)} C_e[\pi_w - \pi_u - c_e]_+. \quad (16)$$

For a feasible flow f_e , use $c_e f_e \geq (\pi_w - \pi_u) f_e - C_e[\pi_w - \pi_u - c_e]_+$ and sum; conservation cancels interior potentials. A value exceeding the available budget is an integer contradiction. The verifier rebuilds the network, capacities, costs and inequality; it does not trust the discovery flow result.

6 Shared fractional systems and safe symmetry

Later tests force different sources to share the same column marginals, F -adjacencies and pair usage. These are necessary fractional systems, not full graph constructions. The complete named-variable constraints are specified in the preserved v6 and LocalIncidence-v7 source and proof notes [7, 8]; this section states their graph-to-system interpretation and principal identities.

Group labels only when demands, full option domains and forced-source masks agree. Group sources only when residual degree, safe cap and forced-label patterns agree. Denote class sizes by N_g and M_k . Average labelled indicators over permutations within these classes. This averages relabelled copies; it does not assert that the unknown graph has any corresponding automorphism. Every linear, invariant necessary inequality remains valid. Average coordinates may be fractional for an actual integral graph.

Let y_{go} be the probability of option o at a label of class g . Let S_{kgo}, T_{kgo} be joint selected and residual event probabilities with that option, and use S_{kg}, T_{kg} for their sums over o . Write $d_g = \sum_o d(o)y_{go}$, $R_g = \sum_o R(o)y_{go}$. Basic equations include

$$\begin{aligned} \sum_o y_{go} &= 1, & \sum_g N_g R_g &= r, & \sum_g N_g d_g &= 2(r + t), \\ \sum_g N_g T_{kg} &= \rho_k, & q_k &= \sum_g N_g S_{kg} \leq c_k, & \sum_k M_k T_{kg} &= R_g, & x_g &= \sum_k M_k S_{kg} \geq s_g. \end{aligned} \quad (17)$$

All event coordinates lie in $[0, 1]$, $S_{kgo} + T_{kgo} \leq y_{go}$, and safe impossible statuses are omitted. Forced selections have $S_{kg} = 1$.

The common edge probability F_{gh} is for two *distinct* labels, with

$$\sum_h (N_h - \mathbf{1}_{\{g=h\}}) F_{gh} = d_g.$$

For a source and a distinct label pair, selected domination gives $F_{gh} + S_{kg} \leq S_{kh} + T_{kh} + 1$ and its reverse-label version. A joint auxiliary V_{kgh} can be chosen to equal the indicator of an F -edge with both labels selected. The relaxation requires

$$V_{kgh} \geq F_{gh} + S_{kg} + S_{kh} - 2, \quad \sum_h (N_h - \mathbf{1}_{\{g=h\}}) V_{kgh} \leq R_g.$$

These are averaged consequences of Lemma 3.4; they do not impose nonlinear independence.

6.1 Selected triple and source-degree types

Let U_{klg} represent $ui \rightarrow w$ with $u \neq w$ in classes k, l and label class g . Then

$$\begin{aligned} \sum_l (M_l - \mathbf{1}_{\{k=l\}}) U_{klg} &= S_{kg}, & \sum_g N_g (U_{klg} + U_{lkg}) &\leq 1, \\ U_{klg} + S_{lg} + T_{lg} &\leq 1. \end{aligned} \tag{18}$$

When $k = l$ both orientations still count; there is no loop. Domination of every other selected label supplies $U_{klg} + S_{kh} - S_{lh} - T_{lh} \leq 1$ and $U_{klg} + F_{gh} + S_{kh} - T_{lh} \leq 2$ for distinct labels. The base model also retains safe linear implications of (3) and (4), using bounds valid in each unselected case.

To stop incompatible source degrees sharing optimistic capacities, use $w_{kq} = \Pr(q_u = q)$ over actual integer values $f_k \leq q \leq c_k$, and A_{kgoq}^X for the joint event of option o , source degree q and status $X \in \{S, T, M\}$. Their marginals recover the same y_{go} and status variables at every source class. Their weighted label counts are qw_{kq} , $\rho_k w_{kq}$, and $(a - q - \rho_k)w_{kq}$.

Typed oriented-pair events P_{klqv} require $q \geq 1$ and $v + \rho_l \geq q - 1$. Their sums recover (18), and their outgoing total is qw_{kq} ; incoming total is at most $(\rho_k + \ell)w_{kq}$. These are joint events, not products of marginals.

6.2 Source-local pair events

The first LocalIncidence-v7 model additionally records $J_{kqghXYe}$ for a source degree, two distinct labels' statuses and their F -adjacency $e \in \{0, 1\}$. Let $c_S = q$, $c_T = \rho_k$, $c_M = a - q - \rho_k$. Marginal identities give

$$\begin{aligned} \sum_h (N_h - \mathbf{1}_{\{g=h\}}) \sum_e J_{kqghXYe} &= (c_Y - \mathbf{1}_{\{X=Y\}}) \sum_o A_{kgoq}^X, \\ \sum_h (N_h - \mathbf{1}_{\{g=h\}}) \sum_Y J_{kqghXY1} &= \sum_o d(o) A_{kgoq}^X, \\ \sum_h (N_h - \mathbf{1}_{\{g=h\}}) J_{kqghSS1} &\leq \sum_o R(o) A_{kgoq}^S. \end{aligned} \tag{19}$$

Events with an F -edge across S, M are omitted. Averaging (6) adds the local residual inequality; unordered pair multiplicities are $N_g N_h$ or $\binom{N_g}{2}$, with both orders for different statuses in one class. Summing over the source degree and statuses gives the same common F_{gh} .

7 Actual endpoint consistency

A fixed column total remains fixed conditional on $q_u = q$. Thus

$$\begin{aligned} \sum_{g,o,X} N_g R(o) A_{kgoq}^X &= r w_{kq}, & \sum_{g,o,X} N_g d(o) A_{kgoq}^X &= 2(r+t) w_{kq}, \\ \sum_{g,o} N_g R(o) A_{kgoq}^S &\leq (r - \rho_k) w_{kq}, & \sum_{g,o} N_g d(o) A_{kgoq}^S &\leq [\rho_k(2a - \rho_k - 3 + q) - 2t] w_{kq}. \end{aligned} \quad (20)$$

These use (7)–(8), not an independence assumption.

Finally retain actual selected label degree x and source indegree p . Set

$$V_{gox} = \Pr(o_i = o, x_i = x), \quad W_{kqp} = \Pr(q_u = q, p_u = p),$$

where $\max(s_g, f_g) \leq x \leq b - R(o)$ and $0 \leq p \leq \min(\rho_k + \ell, b - 1 - q)$; f_g counts forced sources for a label. Require $\sum_x V_{gox} = y_{go}$, $\sum_p W_{kqp} = w_{kq}$ and

$$\sum_p p W_{kqp} = \sum_{l,v} (M_l - \mathbf{1}_{\{l=k\}}) P_{lvq}.$$

The same selected incidence has a joint coordinate

$$Z_{kgoqpx} = \Pr(ui \text{ selected}, o_i = o, q_u = q, p_u = p, x_i = x).$$

It is admitted only for permitted selected status, $x \geq 1$, and

$$R(o) + x \geq q + p. \quad (21)$$

Both endpoints' incidence counts agree:

$$\begin{aligned} \sum_{p,x} Z_{kgoqpx} &= A_{kgoq}^S, & \sum_{o,x} Z_{kgoqpx} &\leq W_{kqp} \quad \text{for each } g, \\ \sum_{g,o,x} N_g Z_{kgoqpx} &= q W_{kqp}, & \sum_{q,p} Z_{kgoqpx} &\leq V_{gox} \quad \text{for each } k, \\ \sum_{k,q,p} M_k Z_{kgoqpx} &= x V_{gox}. \end{aligned} \quad (22)$$

Every equation counts actual incidences; every inequality expresses simplicity. Equation (21) is the actual-type version of Lemma 3.1, stronger than applying it only to averages.

The final production model uses the source-degree model, (20) and (21)–(22), but omits the larger J extension. The three LocalIncidence stages therefore need not be nested. Each rejection has a certificate for its own necessary system, and all stage dispositions are checked. The final seven 197-edge rows are rejected by this endpoint model without integer branching.

8 Exact certificates and complete computation

Proposition 8.1 (Certificate rule). *Let a necessary system be $Ay \leq b$, $Ey = f$, $y \geq 0$. Nonnegative integer inequality multipliers λ and signed integer equality multipliers μ with*

$$\lambda A + \mu E \geq 0 \quad \text{coefficientwise}, \quad \lambda b + \mu f < 0$$

prove infeasibility.

Proof. Their linear combination requires a nonnegative expression in y to be at most a strictly negative number. $\square$

The checking implementation reconstructs named-variable constraints and accumulates exact integers. It verifies multiplier types and signs, constraint identities, the final coefficients, and the strict negative right side. Numerical optimisation proposes multipliers only. Unknown or duplicate certificate records, missing inputs, unsupported scope, timeouts, and unclosed trees are not accepted as proofs.

The equality calculation also uses 91 complete binary proof trees. A split is permitted only on an integral count for every lifted graph: $N_g y_{go}$; $N_g N_h F_{gh}$ or $\binom{N_g}{2} F_{gg}$; $M_k N_g S_{kg}$ or $M_k N_g T_{kg}$; or $M_k w_{kq}$. Children $L \leq h$ and $L \geq h + 1$ cover every integer L . The verifier reconstructs the class-size factors and checks both children at every node. It never branches on a raw average as though it were Boolean. All 572 leaves have exact certificates.

Proposition 8.2 (Recorded finite exclusions). *The complete specified necessary-condition domains at $(n, \Delta, m) = (28, 15, 196)$ and $(28, 15, 197)$ are excluded by the pinned v3–v8 direct-route evidence. The separately specified $(28, 16, 196)$ domain is likewise excluded.*

The exact executed partitions are shown below. The full checking route regenerates the initial domains and the intermediate data, reconstructs all retained column domains, and checks disjoint, complete disposition lists. It compares five equality-stage handoffs byte-for-byte (decompressing where specified). Cardinalities alone are not the coverage proof.

Degree-15 equality stage ($m = 196, t = 1$)	Input	Retained
Charging demand tuples	18,645	1,976
Residual rows of retained intervals	17,669,896	24,411
Projected slack and pair tests	24,411	13,196
Joint columns and one-source constraints	13,196	7,725
Activation cuts and costs	7,725	6,918
Shared-adjacency model	6,918	4,617
Source-degree-type model	4,617	479
Complete branching proofs	479	388
Source-local incidence model	388	366
Conditional global ledgers	366	347
Actual endpoint-degree model	347	0

Degree-15 upper-bound stage ($m = 197, t = 2$)	Input	Retained
Fresh charging demand tuples	12,012	1,229
Residual rows of retained intervals	8,216,928	5,154
Projected slack and pair tests	5,154	2,959
Joint columns and one-source constraints	2,959	1,584
Shared-adjacency model	1,584	797
Source-degree-type model	797	7
Actual endpoint-degree model	7	0

For $\Delta = 16, m = 196, a = 11, t = 4$. The initial domain has 39 demand tuples: 35 are rejected by support/source counts, three by integer duals, and one by all 604 residual rows. A useful hand explanation appears in the v3 proof [4, Section 8]; no graph construction is inferred from its intermediate tuples.

The independent checking route is separate from coefficient discovery, but both implementations were developed by the same assistant. The additional audit kernel checked 8,983 final certificate leaves and 833,923 cited constraint terms [10]. That kernel reuses the preserved model constructors: it is a further arithmetic and branching check, not a third independent derivation of the graph constraints.

9 All degree cases and the equality boundary

9.1 Excluding 197 edges

For $\Delta \leq 14$, the degree sum gives $2m \leq 28 \cdot 14 = 392$. Degree 15 is excluded by Proposition 8.2.

For $\Delta = 16$, $a = 11$ and $t = 5$. Equation (9) requires a sum of eleven scores to be at least $b + 2t = 26$. For every integer $0 \leq s \leq 10$,

$$\frac{s(12 - 2s)}{11 - s} \leq \frac{16}{7}, \quad \frac{16}{7} - \frac{s(12 - 2s)}{11 - s} = \frac{2(s - 4)(7s - 22)}{7(11 - s)} \geq 0.$$

The two numerator factors have the same sign at every permitted integer (including the zero at $s = 4$). Thus the sum is at most $176/7 < 26$, impossible.

For larger degrees use the residual h -index: the largest h for which at least h residual degrees are at least h . A demand s_i requires s_i distinct sources of residual degree at least s_i , so $s_i \leq h$. Activity gives $r \geq b + h(h - 1)$. Hence

$$b + 2t \leq (a + 1)h - h^2 \leq \left\lfloor \frac{(n - b)^2}{4} \right\rfloor. \quad (23)$$

At $m = 197$ it fails for every $b = 17, \dots, 26$:

b	17	18	19	20	21	22	23	24	25	26
$b + 2t$	37	52	71	94	121	152	187	226	269	316
Permitted maximum	30	25	20	16	12	9	6	4	2	1

Degree 27 is the star case. Thus 197 edges are impossible at every maximum degree. Equation (??) now gives $m \leq 196$.

9.2 Characterising 196 edges

If $m = 196$ and $\Delta \leq 14$, every vertex has degree 14. For nonadjacent vertices the two 14-element neighbourhoods lie among the other 26 vertices, and so have at least two common vertices. An edge in a triangle could be deleted without increasing the diameter: its endpoints retain their two-edge path, and any other nonadjacent pair loses at most one of its two common neighbours. This contradicts edge-criticality. The graph is triangle-free.

Choose a vertex. Its 14 neighbours form an independent set, each of whose vertices has degree 14 and therefore neighbours every vertex outside that set. The resulting 196 cross-edges exhaust the graph. It is $K_{14,14}$.

At degree 15 use the equality calculation in Proposition 8.2; at degree 16 use the separate v3 exclusion. At degrees 17–26 the left sides of the preceding table decrease by two when m decreases by one, and remain greater than the right sides. Degree 27 is a star. Conversely $K_{14,14}$ has 196 edges and diameter two; deleting a cross-edge makes its endpoints' distance three. This proves Theorem 1.1, through the stated structural lemmas and computational propositions.

10 Reproducibility, audit limits and reviewer questions

The six immutable archives are v3 demand-support, v4 label-tail, v5 column-propagation, v6 shared-adjacency, LocalIncidence-v7, and direct197-v8. Their 400 original payload files and six manifests are hash-pinned in the companion and release inventory. The separate degree-load-v7 weak-core route is *not* a dependency. No n=25 or n=27 frozen theorem is needed as a black box in the direct order-28 assembly; inherited residual arguments are restated here.

The committed internal audit reran all six components, compared the handoffs, and checked the final certificates with an additional kernel [10]. It found a non-blocking input-validation defect in the standalone v8 C++ helper: an unsupported interval could give a successful empty report. The full proof driver rejects that input first. The new reviewer release makes a guarded derivative of the helper mandatory, verifies the original source hash before substitution, and leaves original archives unchanged. The companion reports the new release’s own tests separately from the historical audit.

Actual-graph regressions and abstract systems test local constraints and implementations, including fractional averages and zero-demand selections. They contain no positive-surplus critical graph. That fact limits what those tests demonstrate; universal necessity must follow from the arguments, not from the sample. The additional kernel’s reuse of model constructors is another explicit trust boundary.

Reviewers should examine: uniqueness in the residual injections; the completeness of the sorted domains and paired column choices; preservation of actual witnesses under cap and option pruning; both-endpoint event normalisations and distinct-endpoint class factors; licensing of integer branch expressions; and the exact statement and applicability of the external Fan bound. Computational reproduction and mathematical review should be reported separately. No author external to this project has yet certified this manuscript, and no formal proof assistant has checked it.

References

- [1] G. Fan, *On diameter 2-critical graphs*, Discrete Mathematics 67 (1987), 235–240. DOI: 10.1016/0012-365X(87)90174-9. Original theorem input; its full proof is not re-audited here.
- [2] T. Wang, *On Murty-Simon Conjecture*, arXiv:1205.4397 (2012), especially p. 2. <https://arxiv.org/abs/1205.4397>. The strict Fan estimate used here is the one stated in this manuscript.
- [3] T. W. Haynes, M. A. Henning, L. C. van der Merwe and A. Yeo, *A maximum degree theorem for diameter-2-critical graphs*, Central European Journal of Mathematics 12 (2014), 1882–1889. DOI: 10.2478/s11533-014-0449-3. Background complement and quasi-edge framework; the observation required here is proved in Section 2.
- [4] P. Lenz project / ChatGPT, *General-order demand–support cuts*, checkpoint 2026-09-07-demand-support-v3, original PROOF.md, source and certificates. Repository `paullenz/MurtySimon25`; exact archive in the companion inventory.
- [5] Same project, *Label-tail v4*, checkpoint 2026-09-07-label-tail-v4, original proof and complete equality-domain replay.
- [6] Same project, *Column propagation and supplement activation v5*, checkpoint 2026-09-07-column-propagation-v5, original proof and source/checking programs.
- [7] Same project, *Shared columns, degree-type transport and threshold activation v6*, checkpoint 2026-09-07-shared-adjacency-v6, especially original PROOF.md Sections 3–6 and named-constraint checking source.
- [8] Same project, *Local incidence and endpoint-degree consistency v7*, checkpoint 2026-09-07-local-incidence-v7, especially original PROOF.md Sections 3–6. This is distinct from degree-load-v7.
- [9] Same project, *Direct exclusion of the 197-edge case at order 28 v8*, checkpoint 2026-09-07-direct-197-v8, original proof, exact inputs, source, and certificates.
- [10] Same project, *Red-team audit of the direct order-28 candidate*, 2026-09-07-redteam-v1. Publication commit `6176ce6428c105b791d9129682cbacd55137edd1`; receipt commit `cd3c22411911280624a8223035e6de2343abbb7c`. Same-assistant internal review; independent review open.