When Acting on Knowledge Makes Free Evidence Harmful: Good's Theorem in Normality Models

Abstract. Knowledge-based decision theory maximizes expected utility with probabilities conditioned on what one knows. I test it against Good's theorem inside Goodman and Salow's normality theory of inductive knowledge. In finite statist, comparable models with K-normality, a free question has non-negative value in every decision problem if and only if no state compatible with one's knowledge is one where the answer would reopen a possibility one now rules out. Given suitable questions, every question is safe if and only if knowledge is gapped, and in graded models no factive state that contains rational belief is safe unless it is the whole evidence. In Goodman and Salow's model of repeated weighing, a free second weighing has negative value at ten-to-one odds. I then examine three expected-utility developments of Salow's Conditional Knowledge Dominance. Conditioning on the options' differing fails wherever the unconditional rule fails; conditioning on the evidence's changing one's choice also fails. Conditioning answer by answer is safe, but it makes the agent pay to check what she knows, which Conditional Knowledge Dominance forbids.

1. Introduction

If you know p, any evidence against p is misleading, so it seems you may avoid it. This is the dogmatism puzzle. Salow (2025) offers the most economical recent dissolution. If Preservation holds (discovering something consistent with your beliefs never requires giving one up), you cannot believe both p and that you might meet evidence defeating p, so the puzzle never arises. If Preservation fails, the puzzle needs Weak Knowledge Dominance, which Salow replaces with a principle that looks equivalent but is not:

> Conditional Knowledge Dominance: If you know that B's consequences would be better than A's if they're not equally good, you should choose B over A. (Salow 2025, 33)

In Salow's Suitcase, you do not know that the suitcase weighs under 18 kg on the hypothesis that it weighs over 17.5 kg, since discovering the latter would defeat that knowledge. So the principle does not tell you to avoid checking. Salow grants that his response "by itself doesn't predict that openness to further evidence is always required", but holds that "a natural development of it will" (2025, 35). The development he has in mind solves "a general problem" about how all-out attitudes guide choices under uncertainty, and "Any solution to that problem will generalize to this one" (2025, 34).

The most developed proposal of that kind is knowledge-based decision theory (KDT): maximize expected utility using probabilities conditioned on what you know. Goldschmidt (2024) gives representation-theorem foundations for knowledge-based decision theories, including Hawthorne and Stanley's (2008), and Hong (2024) explores the view. I ask whether KDT, and expected-utility versions of Salow's conditional principle, keep Good's (1967) theorem that free evidence never has negative expected value, once knowledge is modeled by the most developed formal theory of inductive knowledge, the normality framework of Goodman and Salow (2023, 2025). My thesis:

> In finite statist, comparable normality models, conditioning expected utility on knowledge preserves Good's theorem exactly where knowledge is gapped; of three expected-utility versions of Salow's conditional repair, two fail, and the third succeeds only by abandoning the knowledge-based verdict on costly inquiry.

Theorem 1 identifies when a free question is safe under KDT. Theorem 2 extends its necessity result to factive states containing rational belief. Bjorn's repeated weighing supplies a

continuous example with ordinary payoff ratios (§3). Section 5 then distinguishes three conditional evaluations of inquiry. Its negative results concern specified expected-utility rules, not a refutation of Salow's qualitative principle or every knowledge-based decision theory.

Salow (2025, 33) already notes that a 'for all you know' dominance principle can support dogmatism even with Preservation. Goodman and Salow (2025, henceforth BRN) establish the normality conditions for Preservation and its failure in Bjorn. Geanakoplos (1989; 2002 revision) gives sufficient conditions for valuable information beyond partitional learning; Dorst (2020) and Das (2023) develop related epistemological questions. Hong (2026, §§1–3) challenges normality knowledge through lottery choices, sweetened prospects and information. The general decision-theoretic objection is therefore not new.

Dorst et al.'s (2021) Deference Done Better supplies an even closer general comparison: Total Trust is equivalent to Value, the requirement that every recommended expert-dependent strategy perform at least as well as acting on one's own probabilities (Theorem 2.2). Here the current probabilities are $\pi = \Pr(\cdot \mid K_E(x))$ and the expert's state-indexed probabilities are $P_s = \Pr(\cdot \mid K_{E\cap\Pi(s)}(s))$. The fixed tie order generates a recommended strategy. Universal safety for that strategy implies their Weak Value condition; their Appendix B establishes its equivalence to Value and Total Trust. Theorem 1 thus characterizes, within these normality models, when the post-answer rule merits Total Trust. Its increment is the structural boundary in terms of relative defeat and gappedness, together with the graded-model consequence and the tests of three conditional inquiry rules.

2. Framework

I follow Goodman and Salow (2023) and BRN. A normality structure has a set S of states, a family $\mathcal{E}$ of possible bodies of evidence, and two relations on situations $\langle s, E\rangle$ (with $s \in E \in \mathcal{E}$): $\succeq$ (at least as normal as), a preorder, and $\gg$ (sufficiently more normal than). $\gg$ implies $\succeq$, is irreflexive, and chains: if $w_1 \succeq w_2 \gg w_3 \succeq w_4$ then $w_1 \gg w_4$. Under statism both relations are evidence-independent, so I treat them as relations on states. Comparability says any two states with the same evidence are $\succeq$-related. Collapse says every strict difference in normality is sufficient. For the theorems S is finite.

Rational belief, given evidence E, rules out exactly the evidential possibilities that something else in E is sufficiently more normal than:

$$B(E) = \{s \in E : \text{ no } u \in E \text{ has } u \gg s\}.$$

Knowledge goes beyond evidence by the same mechanism, but its extent depends on where one actually is. I work mainly with K-normality (Goodman and Salow 2023, 104):

$$K_E(x) = B(E) \cup \{v \in E : v \succeq x\}.$$

One knows what is true throughout $K_E(x)$, which I call one's ken. Evidence is transparent, and Goodman and Salow note that "given TRANSPARENCY, K-NORMALITY entails KK" (2023, 107). Their alternatives, I- and KM-normality, validate their principle Margins and allow failures of KK (§6.1). Discovering p at $\langle x, E\rangle$ yields $\langle x, E\cap p\rangle$. A question is a partition Π of E whose cells are discoverable ($E \cap c \in \mathcal{E}$ for each cell c); $\Pi(s)$ is the cell containing s.

Knowledge-based expected utility. Fix a prior Pr with full support. A decision problem is a finite set A of acts, each a function from states to utilities. At $\langle x, E\rangle$, KDT chooses an act maximizing $\Pr(u_a \mid K_E(x))$, ties broken by a fixed order.

The value of a free question. Let $\alpha_\Pi(s)$ be the act KDT chooses at $\langle s, E \cap \Pi(s)\rangle$. The KDT value of answering Π before choosing is

$$V_x(\Pi, A) = \sum_s \Pr(s \mid K_E(x))\, u(\alpha_\Pi(s), s) \;-\; \max_{a \in A} \Pr(u_a \mid K_E(x)).$$

This is Good's comparison of acting now with looking and then acting, with $\Pr(\cdot \mid K_E(x))$ in place of the unconditioned probability. Call Π safe at x if $V_x(\Pi, A) \geq 0$ for every decision problem A.

Relative defeat at x for Π: some $s \in K_E(x)$ has $K_{E\cap\Pi(s)}(s) \not\subseteq K_E(x)$. For all one knows, answering would leave open something one now rules out.

Gapped: every member of $K_E(x)$ is sufficiently more normal than every member of $E \setminus K_E(x)$.

States settle what an unconsulted source would say. This is already so in Goodman and Salow's model, where what Bjorn knows after one weighing depends on the second scale's reading and he has inductive knowledge "about the scales' future readings" (2023, 130 n. 40). The conditioning set is therefore fixed at the time of choice and is the same for both options. If states also settled whether one looks, the conditioning set would depend on the act being evaluated. That is a different route to failures of Good's theorem, and I set it aside.

3. Bjorn's second weighing

Goodman and Salow's (2023, §6) model of Bjorn is statist and comparable, and it violates collapse. Its states are triples $\langle w, y_1, y_2\rangle$: Bjorn's weight and the readings of two scales. With $\delta = (w - y_1)^2 + (w - y_2)^2$, one state is at least as normal as another iff its δ is no larger, and sufficiently more normal iff its δ is smaller by more than c^2. Goodman and Salow derive these relations from independent Gaussian errors (2023, 129). I take the standard deviation as the unit and, following BRN (§5, n. 18), set $c \approx 1.96$.

After the first reading r, Bjorn believes his weight lies in $[r - c, r + c]$. Under K-normality, in any situation with $\delta \leq c^2$ this belief is knowledge, and his ken is $\{\delta \leq c^2\}$. After a second reading r' he believes his weight lies within $c/\sqrt{2}$ of the average of the two readings. As Goodman and Salow show for the case where the first reading is accurate, he "loses knowledge" whenever $|r - r'| > (2 - \sqrt{2})c$ (2023, 127). Since the readings may differ by up to $\sqrt{2}c$ for all he believes, he believes that the second weighing might defeat his knowledge: a failure of Preservation, the case BRN use against it. With independent unit Gaussian errors, losing the old weight-interval belief has probability about 0.417. BRN's estimate 0.367 concerns the narrower event of losing it while the discovery remains consistent with the old beliefs: $(2-\sqrt{2})c < |r-r'| \leq \sqrt{2}c$ (§5, n. 18). Over the full state space, including the readings, Preservation fails at every nonzero difference up to $\sqrt{2}c$, with probability about 0.95. These are different events.

Ten-to-one bet. After the first weighing Bjorn is offered a bet: win 1 if his weight is within c of r, lose 10 otherwise. He knows the bet wins, so KDT takes it. A free second weighing is then offered. After it he would decline the bet whenever the readings differ by more than about $1.12c$, because his new ken gives the losing weights enough probability. Conditional on his present knowledge, this has probability about 0.05, and in each such state he forgoes a payoff he now knows he would get. So KDT values the free weighing at about -0.05; at twenty-to-one, -0.12; at a hundred-to-one, -0.27. A fallibilist who uses the unconditioned probabilities also takes the ten-to-one bet (expected value $0.95 - 10(0.05) = 0.45$), and by

Good's theorem the second weighing has positive value for her. The magnitudes depend on c. On Goodman and Salow's probabilistic account of "sufficiently more normal" (2023, 121), applied to these errors, $c^2 = -2\ln(1-t)$ for probability threshold t; so $c \approx 1.96$ corresponds to $t \approx 0.85$. At $t = 0.95$ ($c \approx 2.45$) the ten-to-one value is about -0.01 and the hundred-to-one value about -0.15.

Three claims about other situations should be kept apart. (i) At every situation after the first weighing, some state compatible with Bjorn's knowledge has a second reading that reopens a state he now rules out, and a bet on the proposition he knows, paying 1 if it holds and losing a finite amount otherwise, makes the second weighing harmful (appendix, A.5). That proposition concerns the readings as well as his weight. (ii) A reopened weight exists only if $\max(c^2, \delta(x)) < (1+\sqrt{2})^2 c^2$; beyond that, which has probability about 10^{-5}, no bet of this form on his weight alone does the job (A.5). (iii) Theorem 2 and its corollary are finite results using the questions defined in §4; the second weighing is not one of them.

4. Where Good's theorem holds

Write M for any assignment of a set $M_F(s) \subseteq F$ to situations, used in place of K in the definitions of §2.

Lemma 1 (defeat is costly). If some $s_0 \in M_E(x)$ has $M_{E\cap\Pi(s_0)}(s_0) \not\subseteq M_E(x)$, then some two-act decision problem gives Π negative value.

Proof. Take a bet β paying 1 on $M_E(x)$ and $-L$ elsewhere, and declining, which pays 0. β has conditional expectation 1, so it is chosen. At s_0 the post-answer state contains a state outside $M_E(x)$, of positive probability, so for large L the agent declines and gets 0 where β paid 1. Nowhere in $M_E(x)$ does she get more than 1. So $V \leq -\Pr(s_0 \mid M_E(x)) < 0$. □

Lemma 2 (laminar sufficiency). Suppose the post-answer states $\{M'(s) : s \in M_E(x)\}$ all lie within $M_E(x)$ and are reflexive ($s \in M'(s)$), transitive ($t \in M'(s)$ implies $M'(t) \subseteq M'(s)$) and nested (any two are disjoint or one contains the other). Then $V \geq 0$ for every decision problem.

This applies Geanakoplos's value-of-information result (1989; 2002 revision, Theorem 1); a short proof is in the appendix.

Lemma 3. Under statism and comparability, for each F the kens $\{K_F(s) : s \in F\}$ are reflexive, transitive and nested.

Theorem 1. Assume statism, comparability and K-normality. (a) A question is safe at x if and only if there is no relative defeat at x for it. (b) If $K_E(x)$ is gapped, every question is safe at x. If it is not, then for some $v \in E \setminus K_E(x)$ the question whether p_v, where $p_v = \{u \in E : \text{not } u \gg v\}$, is unsafe at x if it is available.

Part (a) combines Lemmas 1–3. Part (b) needs only one question at a time, and it rests on a lemma that uses no K-normality.

Lemma 4 (non-gaps are defeasible). Assume statism. Let $m \in M_E(x)$ and $v \in E \setminus M_E(x)$, with not $m \gg v$. Then $m, v \in p_v$ and $v \in B(E \cap p_v)$.

Proof. $m \in p_v$ by hypothesis and $v \in p_v$ by irreflexivity. Nothing in p_v is sufficiently more normal than v. □

Learning p_v is learning that things are not far more normal than v: that one's situation is somewhat abnormal. Goodman and Salow's own discoveries of this kind come by testimony (2023, §8). Probabilistic models predict widespread failures of statism, and theirs obey it only for the bodies of evidence they consider (2023, 138 n. 47); so the statism assumption must cover the extended family.

Say that M leaves open what rational belief leaves open if always $B(F) \subseteq M_F(s)$. Knowledge on each of Goodman and Salow's three completions does, and so do justified true belief and iterated knowledge.

Theorem 2 (only gapped states can be conditioned on). Assume statism, with belief defined from normality as above, and suppose M leaves open what rational belief leaves open. If $M_E(x)$ is not gapped, then for some v the question whether p_v, if available, is unsafe at x for expected utility conditioned on M. A gapped set containing x contains $C(x)$: the states reachable from x by steps, each to a state that is not sufficiently less normal than the last.

The boundary in Theorem 1 is not the boundary of Preservation. Take three states a, b, c, in decreasing order of normality, with $a \gg b$ and $a \gg c$ but not $b \gg c$. Preservation holds here for every body of evidence and every discovery. At b one knows $\{a, b\}$; learning that a is false, which is true at b, reopens c. This is the expected-utility counterpart of Salow's observation about the 'for all you know' variant, located exactly.

Call E graded if $C(x) = E$ for every $x \in E$. Bjorn's evidence after one reading is graded, since a chain of small increases in δ connects any two states.

Corollary. (i) Under statism, collapse and comparability, every K-normality ken is gapped, so KDT makes every question safe. (ii) In a finite graded model, if the questions whether p_v are available, expected utility conditioned on a factive state that leaves open what rational belief leaves open makes them all safe at x only if that state is all of E.

Part (i) is the decision-theoretic counterpart of BRN's result that statism, collapse and comparability validate Preservation. Part (ii) says that, given these questions, the only belief-containing state one can condition on in graded models without some free question becoming harmful is one's evidence. Conditioning on evidence is the response on which "decisions should be governed by underlying probabilities rather than our knowledge" (Salow 2025, 23). Part (ii) depends on the questions. In Bjorn, the state $\{|\mu| \leq 1000\}$ (coordinates as in A.5) is factive at its members, contains rational belief and is not gapped, yet every K-normality ken after the second reading lies inside it, so by Lemma 2 the second weighing is safe for it.

5. The conditional repair

5.1 Three developments

Conditional Knowledge Dominance evaluates options on the hypothesis that they differ. Salow's moral is that "you should always think that, if the evidence you would discover tells strongly against your view, there is a good chance you're wrong" (2025, 36). He gives no expected-utility version. Write $K_E^D(x)$ for what one knows on the hypothesis D. Its K-normality form restricts evidential accessibility to D, as BRN's definition of conditional belief does (App. D.3):

$$K_E^D(x) = \{v \in E \cap D : \text{no } u \in E \cap D \text{ has } u \gg v\} \cup \{v \in E \cap D : v \succeq x\}.$$

Under statism, when $x \in D$, this is the ken one would have on discovering D; in every case it contains the belief one would then have. So it respects Salow's constraint that one is "entitled to believe q on the hypothesis that p only if discovering p would not require you to stop believing q" (2025, 33). Write $u_\ell(s) = u(\alpha_\Pi(s), s)$ for the payoff of looking, and compare looking with an act a in three ways:

(Difference) CEU: look if $\Pr(u_\ell - u_a \mid K_E^D(x)) \geq 0$, where $D = \{s : u_\ell(s) \neq u_a(s)\}$.

(Verdict) VER: look if $\sum_b \Pr(D_b \mid K_E(x)) \Pr(u_b - u_a \mid K_E^{D_b}(x)) \geq 0$, where $D_b = \{s : \alpha_\Pi(s) = b\}$; one conditions on what the evidence would lead one to do.
(Answer) ANS: look if $\sum_c \Pr(c \mid K_E(x)) \Pr(u_\ell - u_a \mid K_E^c(x)) \geq 0$, summing over the cells of Π. These formulas evaluate the future KDT policy α_Π; they do not define fully recursive VER or ANS decision theories.

All three handle Bjorn's bet as Salow's diagnosis predicts. Looking and betting now differ exactly where the second reading would make Bjorn decline, and on that hypothesis his conditional ken includes losing weights. With the actual readings perfectly accurate, at ten-to-one the bet's expected value on that hypothesis is -0.63, so CEU and VER prefer to look; ANS does too, since each decline cell's conditional ken is the post-answer ken.

5.2 Difference and verdict

CEU's hypothesis carries information only because equal-payoff states are excluded from it. A perturbation removes them: if looking and acting now differ at every state, $D = E$, conditional knowledge is ordinary knowledge, and CEU coincides with KDT.

Theorem 3. Assume statism, comparability and K-normality. If there is relative defeat at x for Π, some three-act decision problem makes CEU strictly prefer acting now to answering Π, even if CEU also governs every later choice. KDT gives Π negative value too. These verdicts persist on an open neighborhood of the constructed payoffs.

Construction. Let $M = K_E(x)$ and choose $s_0 \in M$ witnessing relative defeat. Take the bet β of Lemma 1, declining, and $\beta_1 = \beta + \varepsilon\zeta$, where $\zeta(s_0) = N$ and $\zeta = -1$ elsewhere, N gives ζ positive current mean, and $0 < \varepsilon < 1$. Choose L so large that every post-answer ken meeting $E \setminus M$ strictly prefers declining; there are finitely many. Every other post-answer ken excludes s_0, since otherwise transitivity would put s_0's reopened possibilities in it; there β strictly beats β_1 and declining. So looking never selects β_1, which is uniquely best now. All acts differ pairwise at every state, and β_1 differs everywhere from the looking policy; so every comparison CEU makes, now or later, uses ordinary knowledge, CEU agrees with KDT, and no cycle can arise. The gain from acting now at s_0 outweighs the small losses elsewhere (A.6).

For a numerical illustration, take three states $a \succ b \succ c$ with only $a \gg c$, a uniform prior, and $x = a$, whose ken is $\{a, b\}$. Ask whether a; on the other answer the ken is $\{b, c\}$. Let the acts pay $(1, 1, -10)$, $(0, 0, 0)$ and $(0.9, 1.5, -10.1)$. Both rules choose the third now (expectation 1.2). After looking they choose the first at a and decline at b and c, a policy with current expectation 0.5. CEU prefers acting now by 0.7.

Theorem 3 refutes CEU, not Conditional Knowledge Dominance. In the constructed problems the qualitative principle is silent: the most normal state m^* of E lies in the ken, its post-answer ken lies inside the ken and excludes s_0, and there looking yields β, which beats β_1. So one does not know that β_1 is better whenever they differ. More generally, in these models, when looking and acting now differ at every state, Conditional Knowledge Dominance never forbids a free look (A.7). CEU extends the principle into its silence, and there it behaves as KDT does.

VER conditions on the hypothesis Salow's moral names: that the evidence would change what one does. It escapes Theorem 3 for large L, because the conditional ken on 'I would decline' then contains an excluded state, where β_1 loses heavily. It fails elsewhere. Take states a, s_2, s_1, v_2, v_1 with normality ranks $0, 0.2, 0.5, 1.15, 1.3$ (lower is more normal), $u \gg w$ iff w's rank exceeds u's by more than 1, prior $(0.67, 0.01, 0.30, 0.01, 0.01)$, $x = a$, and the question $\{\{a\}, \{s_1, v_1\}, \{s_2, v_2\}\}$. The ken is $\{a, s_1, s_2\}$; both non-trivial answers reopen

something. Use Theorem 3's acts with $s_0 = s_1$, $N = 10$, $\varepsilon = 0.001$, $L = 30.4$. Looking yields β at a and declining elsewhere. On the hypothesis 'I would decline', the conditional ken is $\{s_1, s_2, v_2\}$: $s_2 \gg v_1$ excludes v_1, but v_2 survives with too little weight to sink β_1, whose expected value there is 0.028. VER prefers acting now by 0.008. Every pair of primitive acts differs everywhere, so later choices are KDT's. VER, and CEU too, can even disvalue a question where there is no relative defeat. Take $a \succ x \succ b \succ c$ with $a \gg x, b, c$ and $x \gg c$ only, a uniform prior, the situation x (ken $\{a, x\}$), the question $\{\{x\}, \{b, c\}, \{a\}\}$, and two acts paying $(-2, 3.1, -2, 2.1)$ and $(3, 3, 3, -3)$ at (a, x, b, c). No answer reopens anything, and looking does at least as well as acting now throughout the ken, so KDT values the question at 0.05. On the hypothesis 'I would choose the first act', however, the conditional ken is $\{x, b\}$, and b lies only in an answer she knows to be false; VER and CEU prefer acting now. Conditioning on what the evidence would lead one to do imports possibilities the evidence itself would not make live.

5.3 Answer by answer

Proposition 4. Assume statism, comparability and K-normality. ANS never prefers acting now to a free question.

Proof. Fix a cell c. If $s \in K_E^c(x)$, then $K_{E \cap c}(s) \subseteq K_E^c(x)$: if $s \in B(E \cap c)$ its ken is $B(E \cap c)$ (by the chaining step in A.2), and if $s \succeq x$ then $\{v \succeq s\} \subseteq \{v \succeq x\}$. These kens are reflexive, transitive and nested (Lemma 3), so Lemma 2 with $K_E^c(x)$ in place of $M_E(x)$ makes each summand non-negative. □

The price appears when evidence is not free.

Proposition 5. Under the same assumptions, if there is relative defeat at x for Π, there are a bet β and a price $\kappa > 0$ such that ANS prefers paying κ and answering Π to taking β now, although the two options differ at every state and β pays more at every state of the ken. Conditional Knowledge Dominance therefore requires taking β.

Proof sketch (A.7). Use Lemma 1's bet. The conditional ken on the cell of s_0 contains an excluded state, so for large L that cell's summand is large, while the others are non-negative. Choose also $0 < \kappa < L$. Inside the ken, looking at a price pays at most $1 - \kappa < 1$; outside it pays $-\kappa \neq -L$. □

In Bjorn at normal situations ($c \approx 1.96$), ANS pays up to 0.019 for the second weighing at ten-to-one. At a hundred-to-one it pays up to 1.23, more than the bet pays in any state she has not ruled out, for evidence about a proposition she knows. ANS secures Good's theorem by valuing inquiry with the mixture $\sum_c \Pr(c \mid K_E(x)) \Pr(\cdot \mid K_E^c(x))$, which gives excluded possibilities positive weight; for inquiry, knowledge no longer settles the probabilities. Restricting ANS to free questions, with costly inquiry left to KDT or Conditional Knowledge Dominance, avoids Proposition 5 at another cost: in Bjorn's case the agent then looks at price zero and refuses at every positive price, however small. Thus ANS, this particular conditional repair, keeps free-inquiry safety at the cost of conflicting with Conditional Knowledge Dominance on costly inquiry.

Outside statist models conditional and post-discovery knowledge can diverge (BRN App. D.3), and each rule needs separate assessment there.

6. Objections and replies

6.1 "This is self-trust, not evidence"

The objection says that KDT disvalues questions because the agent distrusts her future self, not the evidence. On the Margins completions, statism, comparability and Margins "together entail NO DEFEAT" (Goodman and Salow 2023, 128). There, by Lemma 4, the reopened possibility lies in the agent's own present ken at m, so the relative defeat in Theorem 2 is a failure of KK at x. Such failures harm the empty question too: by Lemma 1 with the trivial partition, a KDT agent without KK at x pays to decide now rather than later on the same evidence, a cousin of the unmarked-clock case of Ahmed and Salow (2019). For the Margins views I concede the objection. Goodman and Salow report, however, that Goldstein and Hawthorne's (2022) model of multiple observations predicts failures of No Defeat "even given MARGINS" (2023, 130 n. 40), so this escape may be less clean; I have not checked the bet in that model.

For K-normality the objection fails. KK holds, so by Lemmas 2 and 3 the empty question never has negative value (at abnormal situations it can even have positive value). In Bjorn the new reading destroys knowledge despite positive introspection. The structure is the classic dogmatist's: I know the bet wins, the evidence might stop me relying on that, so I avoid it. On the KK-friendly completion, KDT avoids evidence; on the others, it avoids deferring to itself.

6.2 Stakes and encroachment

A ten-to-one payoff ratio is not an absolute measure of stakes: at a fixed ken, positive rescaling of utilities preserves choices and the sign of information value. The finite theorems use large L; Bjorn shows that ten-to-one suffices at $c \approx 1.96$ (§3).

The main reply is pragmatic encroachment. Hawthorne (2004, 181), as Salow reports, suggests that given Good's result the puzzle is handled by the proposal "developed by Fantl and McGrath (2002) and Weatherson (2005)" that encroachment occurs "whenever the difference between maximal and actual probability affects your preferences over available options" (Salow 2025, 24 n. 3). Since the theorems hold the ken fixed across problems, such views escape them. The cost can be made exact. Offered the ten-to-one bet alone, Bjorn bets whether the proposition has probability 1 or 0.95, so on this criterion he keeps his knowledge. Offered the bet and a free second scale, with probability 1 he is indifferent to looking, and with 0.95 he strictly prefers to look. So, on this criterion, the mere offer of free evidence destroys ordinary inductive knowledge. By the reasoning of Corollary (ii), in finite graded models every factive, belief-containing state short of the evidence faces some offer of a free question and a bet that removes it in the same way. Salow objects that nothing would confine encroachment to "these relatively rare counterexamples to Preservation" (2025, 36); in the measurement model the counterexamples are not rare (§3).

6.3 Statism and questions

BRN argue that statism is "the exception rather than the rule" (§7). Bjorn, however, is statist. Beyond it, the questions are whether post-answer kens stay nested and whether Lemma 4's questions still reopen excluded possibilities. On the probabilistic grounding, normality is relative to a question: Goodman and Salow note that relativizing to how much does Bjorn weigh yields a model like Goldstein and Hawthorne's, in which current knowledge is not sensitive to future observations (2023, 130 n. 40). A weight bet may make that question salient. I have not computed the bet there.

6.4 "Good's theorem is not sacrosanct"

Ahmed and Salow (2019) argue that Good's theorem "may be false", using observations that are not partitional. Bjorn's readings partition the evidential possibilities exactly. The failure arises elsewhere: knowledge is lost, so the post-answer probabilities are not the present knowledge-conditioned probabilities conditioned on the reading, and the later self's policy is not the reading-dependent policy the present self would choose. A defender may accept that foresight of such choices can make looking bad. That concedes predictable, evidence-driven knowledge loss with practical cost, which Salow's dissolution aimed to show harmless. If the agent could commit to ignoring an answer, the present best act stays available and a free signal cannot hurt; the results concern looking followed by the specified later rule, without commitment.

7. Conclusion

In finite statist, comparable models with K-normality, expected utility conditioned on knowledge makes every free question safe exactly where knowledge is gapped; any factive state that leaves open what rational belief leaves open needs the same condition when the questions whether p_v are available. On graded normality, which Goodman and Salow defend for measurement, that condition means not going beyond the evidence, and in their own model a free second weighing has negative value at ordinary odds. Of three expected-utility developments of Conditional Knowledge Dominance, conditioning on the options' differing and on the evidence changing one's choice both fail; conditioning answer by answer succeeds, but it pays to check what one knows, contrary to the principle it develops. What has not been shown is that no other way of letting all-out attitudes guide graded choice could deliver the development Salow anticipates, or what happens in non-statist models. The burden on knowledge-conditioned expected utility is precise: weaken the universal safety requirement, restrict the models or questions (collapse is one sufficient restriction), let stakes or offers change knowledge, or stop letting knowledge settle the probabilities used to evaluate inquiry.

Appendix: proofs and computations

A.1 Lemma 2. Let $\mathcal{F} = \{M'(s) : s \in M\}$, $M = M_E(x)$. By nestedness and reflexivity the inclusion-maximal members of $\mathcal{F}$ partition M. For a maximal C, let a_C be the act chosen on C. By induction on $|C|$, $\Pr(u(\alpha(s), s) \mid C) \geq \Pr(u_{a_C} \mid C)$: by transitivity every $s \in C$ has $M'(s) \subseteq C$; states with $M'(s) = C$ choose a_C; the others lie in the maximal proper subkens $D_1, \ldots, D_k$, disjoint members of $\mathcal{F}$ satisfying the hypotheses, on each of which the plan does at least as well as the best act on D_j, hence as a_C. Summing, the plan's expectation on M is at least $\sum_C \Pr(C \mid M) \max_a \Pr(u_a \mid C) \geq \max_a \Pr(u_a \mid M)$. □

A.2 Lemma 3. Reflexivity: $s \succeq s$. Transitivity: let $t \in K_F(s)$. If $t \in B(F)$, every $v \succeq t$ is in $B(F)$ (if $u \gg v$, chaining gives $u \gg t$), so $K_F(t) = B(F) \subseteq K_F(s)$. If $t \succeq s$, $\{v : v \succeq t\} \subseteq \{v : v \succeq s\}$. Nestedness: by comparability the sets $\{v : v \succeq s\}$ are linearly ordered by inclusion, and each ken is their union with $B(F)$. □

A.3 Theorem 1. (a) Kens in different cells are disjoint; by Lemma 3 the post-answer kens are reflexive, transitive and nested; without relative defeat they lie within $K_E(x)$, so Lemma 2 applies. Conversely, Lemma 1. (b) Suppose the ken is gapped, $s \in K_E(x) \cap c$, $v \in c \setminus K_E(x)$. Then $s \gg v$, so $v \notin B(E \cap c)$; and not $v \succeq s$, since $v \succeq s \gg v \succeq v$ would give $v \gg v$. So $K_{E\cap c}(s) \subseteq K_E(x)$. If the ken is not gapped, Lemma 4 gives relative defeat for the question whether p_v, because $B(E \cap p_v) \subseteq K_{E\cap p_v}(m)$. □

A.4 Theorem 2. Lemmas 4 and 1 give the first claim. If M is gapped, $m \in M$ and not $m \gg v$, then $v \in M$; so M is closed under the steps generating $C(x)$. □ Corollary (i). Let $m \in K_E(x)$, $v \notin K_E(x)$. Then not $v \succeq x$, and some $u \gg v$. If $m \succeq x$, comparability and collapse give $x \gg v$, hence $m \gg v$. If $m \in B(E)$, then $v \succeq m$ would give $u \gg m$, so m is strictly more normal than v and collapse gives $m \gg v$. □

A.5 Bjorn. After the first reading, let $\mu = (e_1+e_2)/2$ and $d = e_1-e_2$ for the scale errors, so $\delta = 2\mu^2+d^2/2$; for the numerical model, $\mu \sim N(0,1/2)$ and $d \sim N(0,2)$ independently (the swamped-prior treatment of weight). A second reading fixes d, and the post-answer ken of a state s is $|\mu| \le \max(c/\sqrt{2}, |\mu_s|)$. Let $R = \max(c^2, \delta(x))$, so the current ken is $M = \{\delta \le R\}$. Choose s with $\mu_s = 0$ and $d_s^2/2$ strictly between $R - c^2$ and R; its post-answer ken reaches $\delta = c^2 + d_s^2/2 > R$. By strictness, a small open neighborhood U of s stays inside M and has post-answer probability of M^c at least some $\eta > 0$, and U has positive probability. A bet paying 1 on M and $-L$ elsewhere, with $L > (1-\eta)/\eta$, is declined throughout U and accepted now, so looking has value at most $-\Pr(U \mid M) < 0$. For weights: $e_1 = \mu + d/2$, and by Cauchy–Schwarz $|e_1| \le \sqrt{R}$ throughout M. A post-answer ken of $s \in M$ reaches $|e_1|$ up to $\max(c/\sqrt{2}, |\mu_s|) + |d_s|/2$. If $|\mu_s| > c/\sqrt{2}$, this is at most $\sqrt{R}$; otherwise its maximum, $c/\sqrt{2} + \sqrt{R/2}$, exceeds $\sqrt{R}$ iff $R < (1+\sqrt{2})^2 c^2$. Under the Gaussian model $\Pr(\delta \ge (1+\sqrt{2})^2 c^2) \approx 1.4 \times 10^{-5}$. The normal-situation figures in §3 integrate the density over $\delta \le c^2$ for the bet on $|e_1| \le c$.

A.6 Theorem 3. Put $P = \Pr(\cdot \mid M)$, choose N with $NP(s_0) > 2$ and fix $0 < \varepsilon < 1$. For each post-answer ken F meeting $E \setminus M$, $q_F = \Pr(E \setminus M \mid F) > 0$; the expectations of β_1 and β on F are at most $1 + N - Lq_F$, so $L > (1+N)/\min_F q_F$ makes declining strictly best on every such F, for every $\varepsilon < 1$. On every other post-answer ken, $\beta_1 = 1 - \varepsilon$ and $\beta = 1$. Now β_1 is uniquely optimal since $P(\zeta) > 0$. At s_0 acting now gains $1 + \varepsilon N$; elsewhere in M looking gains at most ε. So $P(u_{\beta_1} - u_\ell) \ge P(s_0)(1+\varepsilon N) - \varepsilon > 0$. Outside M looking declines while $\beta_1 = -L - \varepsilon$; inside, looking pays 0 or 1, neither equal to β_1. So the difference event is E, and likewise for every pair of primitive acts. Finitely many strict inequalities and nonzero differences persist under small perturbations. □

A.7 CKD silence; Propositions 4–5. Silence. Let m^* be $\succeq$-maximal in E and c^* its cell. If $v \notin K_E(x)$, some $u \gg v$ and chaining gives $m^* \gg v$; so $B(E \cap c^*) \subseteq K_E(x)$, and every state there has post-answer ken $B(E \cap c^*)$ and chooses one act b^*, optimal on that set. If looking differs from a everywhere, b^* beats a at some state of $B(E \cap c^*)$, so one does not know that a is better whenever they differ. Proposition 5. Let $c_0 = \Pi(s_0)$. If $s_0 \succeq x$, $K_E^{c_0}(x) \supseteq K_{E \cap c_0}(s_0)$; if $s_0 \in B(E)$, the part of $K_{E \cap c_0}(s_0)$ above s_0 lies in $B(E)$ by chaining, so its excluded states lie in $B(E \cap c_0) \subseteq K_E^{c_0}(x)$. Either way $K_E^{c_0}(x)$ contains an excluded state, with probability $q > 0$ in it. By the proof of Proposition 4, the c_0 summand is at least $-\Pr(u_\beta \mid K_E^{c_0}(x)) \ge Lq - 1$ and the others are non-negative; so choose $0 < \kappa < \min\{L, \Pr(c_0 \mid K_E(x))(Lq-1)\}$. □

Computations. Supporting scripts enumerate statist comparable structures with up to four states, every nonempty body of evidence, situation and nontrivial partition (64,004 situation–question pairs; 3,306 with relative defeat). Structural enumeration is exhaustive within that domain; sampled priors and utilities test examples, not the universal decision-problem quantifiers proved above. The examples in §5.2 were also evaluated directly. Bjorn's decision values use numerical integration. His weight-belief-loss probability is $\operatorname{erfc}((2-\sqrt{2})c/2)$; subtracting $\operatorname{erfc}(c/\sqrt{2})$ gives loss with a consistent discovery.

References

Ahmed, Arif, and Bernhard Salow. 2019. “Don’t Look Now.” British Journal for the Philosophy of Science 70: 327–350.

Das, Nilanjan. 2023. “The Value of Biased Information.” British Journal for the Philosophy of Science 74: 25–55.

Dorst, Kevin, Benjamin A. Levinstein, Bernhard Salow, Brooke E. Husic, and Branden Fitelson. 2021. “Deference Done Better.” Philosophical Perspectives 35: 99–150. https://doi.org/10.1111/phpe.12156.

Dorst, Kevin. 2020. “Evidence: A Guide for the Uncertain.” Philosophy and Phenomenological Research 100: 586–632.

Fantl, Jeremy, and Matthew McGrath. 2002. “Evidence, Pragmatics, and Justification.” Philosophical Review 111: 67–94.

Geanakoplos, John. 1989. “Game Theory without Partitions, and Applications to Speculation and Consensus.” Cowles Foundation Discussion Paper 914, Yale University. Revised June 18, 2002; the revision was consulted for Theorem 1.

Goldschmidt, Zeev. 2024. “Foundations for Knowledge-Based Decision Theories.” Australasian Journal of Philosophy 102: 939–958.

Goldstein, Simon, and John Hawthorne. 2022. “Knowledge from Multiple Experiences.” Philosophical Studies 179: 1341–1372.

Good, I. J. 1967. “On the Principle of Total Evidence.” British Journal for the Philosophy of Science 17: 319–321.

Goodman, Jeremy, and Bernhard Salow. 2023. “Epistemology Normalized.” Philosophical Review 132: 89–145.

Goodman, Jeremy, and Bernhard Salow. 2025. “Belief Revision Normalized.” Journal of Philosophical Logic 54: 1–49.

Hawthorne, John. 2004. Knowledge and Lotteries. Oxford: Oxford University Press.

Hawthorne, John, and Jason Stanley. 2008. “Knowledge and Action.” Journal of Philosophy 105: 571–590.

Hong, Frank. 2024. “Know Your Way Out of St. Petersburg: An Exploration of ‘Knowledge-First’ Decision Theory.” Erkenntnis 89: 2473–2492.

Hong, Frank. 2026. “Normality in Action?” Philosophical Perspectives, early view. https://doi.org/10.1111/phpe.70003.

Salow, Bernhard. 2025. “Fallibility and Dogmatism.” Australasian Journal of Philosophy 103: 23–38.

Weatherson, Brian. 2005. “Can We Do Without Pragmatic Encroachment?” Philosophical Perspectives 19: 417–443.