Unified Cosmology from a Dynamical Gravitational Coupling

Leptogenesis, Expansion-Rate Modification, and Dark-Sector Dynamics

Paul Michel
Independent Researcher
April 1, 2026

Abstract

We present a minimal scalar–tensor cosmological framework in which a single dynamical degree of freedom governs the effective strength of gravity through a spacetime-dependent Planck mass. By promoting the normalization of the Einstein–Hilbert action to a dynamical quantity, the model introduces a time-dependent gravitational coupling while preserving local Lorentz invariance and standard particle physics.

Within this framework, four major problems in modern cosmology are addressed in a unified manner: the baryon asymmetry of the Universe, the Hubble tension, the nature of dark matter, and the origin of dark energy. The scalar field evolves according to a single small parameter that induces cumulative deviations in the expansion history, providing a natural mechanism for alleviating the Hubble tension.

The kinetic component of the field scales approximately as $a^{-3(1-n)}$, reproducing cold dark matter behavior at early and intermediate epochs, while its potential drives late-time accelerated expansion. In this way, dark matter and dark energy emerge as different dynamical regimes of a single field.

A derivative coupling between the scalar field and the lepton current generates a dynamical chemical potential, augmenting thermal leptogenesis and reducing the required CP violation and mass scale. Baryogenesis, expansion dynamics, and dark-sector behavior therefore arise as correlated consequences of a single evolving degree of freedom.

The framework predicts a time-dependent gravitational coupling, percent-level modifications to the growth of large-scale structure, and a difference between gravitational-wave and electromagnetic luminosity distances. These signatures provide direct observational tests across multiple cosmological probes.

The model is dynamically stable and consistent with current constraints, and constitutes a minimal and predictive extension of ΛCDM in which multiple fundamental cosmological phenomena are linked to the evolution of the gravitational sector.

1 Introduction

The ΛCDM model has achieved remarkable success in describing the large-scale structure and evolution of the Universe. High-precision observations of the cosmic microwave background (CMB), baryon acoustic oscillations (BAO), Type Ia supernovae, and galaxy clustering collectively support a cosmological framework in which approximately 5% of the energy density is in baryonic matter, $\sim 25\%$ in cold dark matter, and $\sim 70\%$ in dark energy. This phenomenological model provides an excellent fit to a wide range of observational data across multiple scales.

Despite this empirical success, several fundamental theoretical problems remain unresolved. One of the most prominent is the origin of the baryon asymmetry of the Universe, characterized by the baryon-to-photon ratio

$$\eta \approx 6.1 \times 10^{-10}. \tag{1}$$

This asymmetry cannot be generated within the Standard Model of particle physics and requires new physics that satisfies the Sakharov conditions. Thermal leptogenesis provides a compelling mechanism, but it typically requires large CP-violating phases and high-scale right-handed neutrino masses, which remain experimentally inaccessible.

Another major challenge is the Hubble tension, a persistent discrepancy between early-Universe measurements of the Hubble constant inferred from CMB observations and late-Universe measurements based on distance ladder techniques. This tension has reached a level of statistical significance that suggests the presence of new physics beyond the standard cosmological model, particularly in the dynamics governing the expansion history.

In addition, the microscopic nature of dark matter remains unknown. While cold dark matter successfully explains structure formation at large scales, no direct detection of dark matter particles has been confirmed. Similarly, the physical origin of dark energy, responsible for the observed accelerated expansion of the Universe, remains one of the most profound open problems in modern physics. The simplest explanation, a cosmological constant, suffers from severe fine-tuning issues when compared to theoretical expectations from quantum field theory.

Most proposed extensions of ΛCDM address these problems independently. For example, leptogenesis models introduce new heavy particles, modified gravity models alter the structure of Einstein's equations, and dark energy models often involve additional scalar fields with carefully tuned potentials. While such approaches can be phenomenologically viable, they typically introduce multiple new degrees of freedom and parameters, raising questions about theoretical economy and underlying physical principles.

This situation motivates the exploration of unified frameworks in which a single dynamical degree of freedom can simultaneously address multiple cosmological phenomena. Such an approach not only reduces theoretical complexity but also provides a more coherent picture of the underlying physics governing the Universe.

In this work, we investigate a scalar–tensor cosmological framework in which a single scalar field $\Phi(x)$ governs the effective normalization of gravitational interactions. The field is interpreted as a spacetime-dependent Planck mass, corresponding to a dynamical gravitational coupling. This construction provides a minimal extension of general relativity in which the strength of gravity evolves over cosmological timescales while preserving local Lorentz invariance and the standard structure of particle interactions.

We show that the evolution of this scalar field produces multiple cosmological effects. First, its time dependence modifies the expansion history through additional terms in the Friedmann equations, leading to small but cumulative deviations from the standard ΛCDM prediction and providing a natural mechanism for alleviating the Hubble tension. Second, its kinetic contribution scales in a manner that effectively mimics cold dark matter at early times. Third, its potential energy can dominate at late times, producing accelerated expansion consistent with dark energy.

Furthermore, we introduce a derivative coupling between the scalar field and the lepton current, which generates a dynamical chemical potential. This mechanism augments standard thermal leptogenesis by providing an additional source of lepton asymmetry, thereby reducing the required CP violation and lowering the scale of new physics.

A key feature of the framework is that all these effects are controlled by a single small parameter governing the evolution of the scalar field. This provides a unified description in which baryogenesis, expansion dynamics, and dark-sector phenomenology are interconnected rather than independent.

The structure of the paper is as follows. In Sec. II, we introduce the scalar field and its interpretation as a dynamical Planck mass. In Sec. III, we derive the action and modified field equations. In Sec. IV, we analyze the relation to scalar–tensor theories. In Sec. V, we derive the modified Friedmann equations governing cosmological evolution. Subsequent sections explore the

evolution of the scalar field, leptogenesis, the effective gravitational coupling, stability, quantum implications, and observational signatures. We conclude with a discussion of the implications and future directions of this framework.

A central result of this work is that baryogenesis, the Hubble tension, dark matter behavior, and dark energy all emerge as correlated consequences of a single evolving scalar degree of freedom. Rather than introducing separate mechanisms for each of these phenomena, the present framework provides a unified dynamical origin rooted in the evolution of the gravitational sector.

2 Dimensional Motivation and Scalar Definition

The starting point for the present framework is the Einstein–Hilbert action, which defines the classical dynamics of spacetime geometry,

$$S_{\mathrm{EH}} = \frac{1}{16\pi G} \int d^4x \sqrt{-g}\, R, \tag{2}$$

where R is the Ricci scalar and G is Newton's gravitational constant. The normalization of this action sets the scale relating spacetime curvature to energy–momentum and is conventionally expressed in terms of the (reduced) Planck mass,

$$M_P^2 = \frac{1}{8\pi G}. \tag{3}$$

In standard general relativity, M_P is treated as a constant, fixing the strength of gravitational interactions. However, this normalization can be viewed more generally as defining the coupling between geometry and matter. This observation motivates the possibility that the effective Planck mass may be a dynamical quantity rather than a fixed constant.

We therefore introduce a scalar field $\Phi(x)$ that defines a spacetime-dependent effective Planck mass,

$$M_P^2(x) \equiv \Phi(x), \tag{4}$$

and promote the normalization of the Einstein–Hilbert action to a dynamical quantity. The gravitational action then becomes

$$S = \int d^4x \sqrt{-g} \left[\frac{\Phi(x)}{16\pi G} R \right]. \tag{5}$$

This construction provides a minimal extension of general relativity in which the strength of gravity evolves dynamically while preserving the geometric structure of the theory.

From a physical perspective, the scalar field Φ controls the effective gravitational coupling. Comparing the modified field equations to the standard Einstein equations, one finds

$$G_{\mathrm{eff}} = \frac{G}{\Phi}, \tag{6}$$

indicating that variations in Φ correspond to a time-dependent strength of gravity.

This framework belongs to the class of scalar–tensor theories, in which the gravitational coupling evolves over time. However, unlike models in which scalar fields are introduced phenomenologically, the present construction directly ties the scalar field to the normalization of the gravitational action, providing a clear physical interpretation in terms of a dynamical Planck mass.

An important feature of this formulation is that it does not imply any modification of local Lorentz invariance or the propagation of light. The scalar field affects only the normalization of the gravitational sector, leaving local particle physics processes unchanged to leading order.

The evolution of the scalar field is naturally linked to the cosmological expansion, which provides the characteristic timescale in a homogeneous and isotropic universe. We therefore consider scaling solutions in which the fractional rate of change of the field is proportional to the Hubble parameter,

$$\frac{\dot{\Phi}}{\Phi} \sim H. \tag{7}$$

A minimal parametrization is given by

$$\frac{\dot{\Phi}}{\Phi} = 3nH, \tag{8}$$

where $n \ll 1$ is a dimensionless parameter controlling the rate of evolution. This relation describes a slowly varying scalar field that tracks the cosmological expansion while remaining consistent with observational constraints.

In summary, the introduction of a dynamical Planck mass provides a minimal and physically motivated extension of general relativity. It introduces a single new degree of freedom that governs the effective strength of gravity and sets the stage for a unified description of multiple cosmological phenomena.

3 Action and Field Equations

Having introduced the scalar field $\Phi(x)$ as a dynamical quantity corresponding to a spacetime-dependent effective Planck mass,

$$M_P^2(x) \equiv \Phi(x), \tag{9}$$

we now construct the corresponding scalar–tensor theory.

We consider the action

$$S = \int d^4x \sqrt{-g} \left[\frac{\Phi}{16\pi G} R - \frac{1}{2} \partial_\mu \Phi \, \partial^\mu \Phi - V(\Phi) + \mathcal{L}_m \right], \tag{10}$$

where R is the Ricci scalar, $V(\Phi)$ is a scalar potential, and $\mathcal{L}_m$ represents the matter Lagrangian. The scalar field multiplies the curvature term, thereby promoting the normalization of the gravitational action to a dynamical quantity, while the kinetic and potential terms govern its propagation and evolution.

3.1 Variation with Respect to the Metric

To derive the gravitational field equations, we vary the action with respect to the metric $g_{\mu\nu}$. The variation of the gravitational term yields

$$\delta \left(\sqrt{-g} \, \Phi R \right) = \sqrt{-g} \left[\Phi G_{\mu\nu} + \nabla_\mu \nabla_\nu \Phi - g_{\mu\nu} \Box \Phi \right] \delta g^{\mu\nu}. \tag{11}$$

The variation of the scalar kinetic term gives

$$\delta \left(-\frac{1}{2} \sqrt{-g} \, \partial_\mu \Phi \, \partial^\mu \Phi \right) = \sqrt{-g} \left[\partial_\mu \Phi \, \partial_\nu \Phi - \frac{1}{2} g_{\mu\nu} (\partial \Phi)^2 \right] \delta g^{\mu\nu}, \tag{12}$$

and the variation of the potential contributes

$$\delta \left(-\sqrt{-g} \, V(\Phi) \right) = -\sqrt{-g} \, V(\Phi) \, g_{\mu\nu} \, \delta g^{\mu\nu}. \tag{13}$$

Including the variation of the matter action, we obtain the modified Einstein equations,

$$\Phi G_{\mu\nu} = 8\pi G\, T_{\mu\nu} + \nabla_\mu \nabla_\nu \Phi - g_{\mu\nu} \Box \Phi + \partial_\mu \Phi\, \partial_\nu \Phi - \frac{1}{2} g_{\mu\nu} (\partial \Phi)^2 - g_{\mu\nu} V(\Phi). \tag{14}$$

These equations describe a scalar–tensor theory in which the scalar field Φ is nonminimally coupled to curvature and contributes additional dynamical terms to the gravitational field equations.

3.2 Effective Energy–Momentum Tensor

It is useful to rewrite the field equations in a form resembling the standard Einstein equations,

$$G_{\mu\nu} = 8\pi G_{\text{eff}}\, T^{\text{tot}}_{\mu\nu}, \tag{15}$$

where the effective gravitational coupling is given by

$$G_{\text{eff}} = \frac{G}{\Phi}. \tag{16}$$

The scalar field contributions can be grouped into an effective energy–momentum tensor,

$$T^{(\Phi)}_{\mu\nu} = \frac{1}{8\pi G} \left[\nabla_\mu \nabla_\nu \Phi - g_{\mu\nu} \Box \Phi + \partial_\mu \Phi\, \partial_\nu \Phi - \frac{1}{2} g_{\mu\nu} (\partial \Phi)^2 - g_{\mu\nu} V(\Phi) \right]. \tag{17}$$

In this form, the scalar field can be interpreted as an additional source of energy and momentum that modifies the dynamics of spacetime.

3.3 Scalar Field Equation of Motion

Varying the action with respect to the scalar field Φ yields its equation of motion,

$$\Box \Phi - \frac{dV}{d\Phi} + \frac{1}{16\pi G} R = 0. \tag{18}$$

This equation shows that the scalar field is sourced not only by its potential but also directly by spacetime curvature, reflecting its role as a dynamical normalization of the gravitational sector.

3.4 Physical Interpretation

The structure of the modified field equations highlights several important features of the theory.

First, the scalar field directly controls the effective gravitational coupling through the factor $1/\Phi$, implying that the strength of gravity evolves over cosmological time.

Second, the presence of second-derivative terms in the form $\nabla_\mu \nabla_\nu \Phi$ and $\Box \Phi$ indicates a nonminimal coupling between the scalar field and curvature, leading to modifications of gravitational dynamics even in the absence of conventional matter sources.

Third, the canonical kinetic term ensures that the scalar field propagates as a well-defined dynamical degree of freedom, avoiding pathological behavior such as ghost instabilities.

Finally, the potential $V(\Phi)$ governs the late-time behavior of the field and allows for the possibility of accelerated expansion.

Together, these features define a minimal scalar–tensor extension of general relativity in which a single field governs both the effective strength of gravity and the dynamical evolution of the cosmological background.

4 Relation to Brans–Dicke and Scalar–Tensor Theories

The scalar–tensor framework developed in this work shares structural similarities with Brans–Dicke (BD) theory and its generalizations. It is therefore important to clarify both the correspondence and the distinctions between the present model and conventional scalar–tensor gravity.

4.1 Brans–Dicke Theory: Brief Overview

Brans–Dicke theory is defined by the action

$$S_{\mathrm{BD}} = \int d^4x\sqrt{-g}\left[\phi R - \frac{\omega_{\mathrm{BD}}}{\phi}\partial_\mu\phi\partial^\mu\phi - V(\phi) + \mathcal{L}_m\right], \tag{19}$$

where ϕ is a scalar field controlling the effective gravitational coupling, and ω_{BD} is the Brans–Dicke parameter. The effective Newton constant is

$$G_{\mathrm{eff}} \propto \frac{1}{\phi}. \tag{20}$$

In the limit $\omega_{\mathrm{BD}} \to \infty$, the theory reduces to general relativity.

4.2 Mapping to the Present Framework

The action considered in this work is

$$S = \int d^4x\sqrt{-g}\left[\Phi R - \frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi - V(\Phi) + \mathcal{L}_m\right]. \tag{21}$$

At the level of field content and coupling structure, this corresponds to a scalar–tensor theory with a nonminimal coupling ΦR and a canonical kinetic term. The effective gravitational coupling is given by

$$G_{\mathrm{eff}} = \frac{G}{\Phi}, \tag{22}$$

which matches the functional form found in Brans–Dicke theory.

A formal correspondence with Brans–Dicke theory can be obtained by redefining the scalar field. Comparing the kinetic terms,

$$\frac{\omega_{\mathrm{BD}}}{\phi}(\partial\phi)^2 \quad \longleftrightarrow \quad \frac{1}{2}(\partial\Phi)^2, \tag{23}$$

one finds that the present model corresponds to a field-dependent effective Brans–Dicke parameter,

$$\omega_{\mathrm{eff}}(\Phi) = \frac{\Phi}{2}. \tag{24}$$

This implies that the theory does not correspond to a constant-ω_{BD} model, but rather to a dynamical scalar–tensor theory in which the effective coupling evolves with the scalar field itself.

In the present framework, the scalar field is interpreted as a dynamical Planck mass rather than an independent phenomenological field.

4.3 Key Differences from Standard Brans–Dicke Theory

Despite the structural similarities, the present framework differs from conventional Brans–Dicke models in several important ways:

- **Geometric Origin of the Scalar Field:** In Brans–Dicke theory, the scalar field ϕ is introduced phenomenologically. In contrast, the scalar field Φ considered here arises directly from promoting the normalization factor of the Einstein–Hilbert action,

$$\frac{c^3}{16\pi G} \longrightarrow \frac{\Phi(x)}{16\pi G}, \tag{25}$$

  providing a geometric motivation tied to the structure of gravitational dynamics.

- **Canonical Kinetic Structure:** The kinetic term in the present model is canonical, whereas Brans–Dicke theory employs a noncanonical form proportional to $\omega_{\rm BD}/\phi$. This leads to different propagation and normalization properties for the scalar degree of freedom.

- **Single-Parameter Cosmological Control:** The cosmological evolution of the scalar field is governed by a single small parameter n, through the scaling relation

$$\frac{\dot{\Phi}}{\Phi} = 3nH. \tag{26}$$

  This parameter simultaneously controls deviations in the expansion history, the evolution of the gravitational coupling, and the generation of baryon asymmetry. Such unified control is not a generic feature of Brans–Dicke models.

- **Unified Dark Sector Behavior:** In the present framework, the scalar field naturally interpolates between matter-like and dark-energy-like behavior:

$$\rho_{\Phi,\rm kin} \propto a^{-3(1-n)} \quad \text{(dark matter-like)}, \tag{27}$$

$$\rho_{\Phi,\rm pot} \to \text{const.} \quad \text{(dark energy-like)}. \tag{28}$$

  While scalar fields in Brans–Dicke-type theories can contribute to cosmological dynamics, they are not typically constructed to simultaneously account for both dark matter and dark energy within a single degree of freedom.

- **Derivative Coupling to Matter Currents:** The inclusion of a derivative coupling of the form

$$\mathcal{L}_{\rm int} = \frac{1}{M_*}(\partial_\mu \Phi) J_L^\mu \tag{29}$$

  introduces a dynamical chemical potential that contributes to baryogenesis. This interaction is not part of standard Brans–Dicke theory and provides a direct link between cosmological evolution and particle asymmetry generation.

- **Absence of Varying Fundamental Constants:** While scalar–tensor models with covarying fundamental constants (e.g., linked variations of G and c) have been considered, the present framework does not invoke variations of fundamental constants, instead introducing a minimal dynamical Planck mass with direct cosmological observables.

4.4 Interpretation

The present model can therefore be viewed as a specific realization within the broader class of scalar–tensor theories, with the distinguishing feature that the scalar field is tied directly to the normalization of the gravitational action. This leads to a more constrained structure in which a single dynamical degree of freedom governs multiple cosmological phenomena.

Rather than representing a generic modification of gravity, the framework provides a minimal extension in which variations in the effective Planck mass,

$$M_P^2 \propto \Phi, \tag{30}$$

are responsible for interconnected effects in cosmic expansion, structure formation, and baryogenesis.

4.5 Observational Distinction

Although the background evolution may resemble that of other scalar–tensor models, the combination of effects predicted here—including a specific scaling of the gravitational coupling, unified dark sector behavior, and scalar-induced baryogenesis— provides a set of correlated observables that can distinguish this framework from standard Brans–Dicke theory.

Future constraints from large-scale structure, gravitational-wave standard sirens, and precision measurements of the time variation of G will be particularly important in testing these distinctions.

4.6 Mapping to Effective Field Theory of Dark Energy

The present framework can be interpreted within the effective field theory (EFT) of dark energy as a model with a running Planck mass. Identifying the effective Planck mass as $M_*^2(a) \equiv \Phi(a)$, the evolution

$$\Phi(a) \propto a^{3n}$$

implies

$$\alpha_M \equiv \frac{d\ln M_*^2}{d\ln a} = 3n = \text{constant}.$$

All other EFT functions vanish,

$$\alpha_B = 0, \quad \alpha_K = 0, \quad \alpha_T = 0,$$

corresponding to a minimal scalar–tensor extension with a constant Planck mass running.

In this language, the modification to gravitational-wave propagation follows directly as

$$d_L^{\rm GW} = d_L^{\rm EM}(1+z)^{3n/2},$$

providing a direct observational probe of the parameter n.

This mapping places the present framework within the standard EFT classification of modified gravity, enabling direct comparison with existing observational constraints on the running of the Planck mass.

5 Modified Friedmann Equations

To study the cosmological implications of the scalar–tensor framework introduced in the previous section, we specialize the field equations to a homogeneous and isotropic background described by the spatially flat Friedmann–Lemaître–Robertson–Walker (FLRW) metric,

$$ds^2 = -dt^2 + a(t)^2 d\vec{x}^2,$$

where $a(t)$ is the scale factor and $H \equiv \dot{a}/a$ is the Hubble parameter.

We assume that the scalar field Φ is spatially homogeneous,

$$\Phi = \Phi(t),$$

consistent with the symmetries of the FLRW background.

5.1 Modified Friedmann Equation

The (00) component of the modified Einstein equations yields the generalized Friedmann equation. Substituting the FLRW metric into the field equations and collecting terms, we obtain

$$3\Phi H^2 = 8\pi G(\rho_m + \rho_r) + \frac{1}{2}\dot{\Phi}^2 + V(\Phi) - 3H\dot{\Phi}.$$

Dividing by 3Φ, this can be written as

$$H^2 = \frac{8\pi G}{3\Phi}(\rho_m + \rho_r + \rho_\Phi) - H\frac{\dot{\Phi}}{\Phi},$$

where the scalar energy density is defined as

$$\rho_\Phi = \frac{1}{2}\dot{\Phi}^2 + V(\Phi).$$

This equation differs from the standard Friedmann equation in two important ways. First, the energy densities are scaled by the inverse of the scalar field, reflecting a dynamical gravitational coupling. Second, an additional term proportional to $H\dot{\Phi}/\Phi$ appears, arising from the time variation of the scalar field.

5.2 Acceleration Equation

The spatial components of the field equations yield the modified acceleration equation. After simplification, one obtains

$$2\Phi\dot{H} + 3\Phi H^2 = -8\pi G p_{\text{tot}} - \frac{1}{2}\dot{\Phi}^2 + V(\Phi) - 2H\dot{\Phi} - \ddot{\Phi},$$

where $p_{\text{tot}} = p_m + p_r$ includes contributions from matter and radiation.

This equation governs the evolution of the expansion rate and shows explicitly how the scalar field contributes both kinetic and potential terms, as well as derivative couplings to the expansion.

5.3 Effective Energy Density and Pressure

It is useful to interpret the scalar field as an effective fluid with energy density and pressure defined by grouping the scalar contributions appearing in the modified Einstein equations.

The effective energy density is given by

$$\rho_\Phi = \frac{1}{2}\dot{\Phi}^2 + V(\Phi), \tag{31}$$

which includes the canonical kinetic term and the scalar potential.

The effective pressure is obtained from the spatial components of the field equations and takes the form

$$p_\Phi = \frac{1}{2}\dot{\Phi}^2 - V(\Phi) + \ddot{\Phi} + 2H\dot{\Phi} \tag{32}$$

The additional derivative terms in the pressure arise from the nonminimal coupling between the scalar field and curvature, specifically from the combination $\nabla_\mu\nabla_\nu\Phi - g_{\mu\nu}\Box\Phi$ in the field equations. These terms distinguish the present framework from minimally coupled scalar-field models.

The effective equation-of-state parameter is then defined as

$$w_\Phi = \frac{p_\Phi}{\rho_\Phi}. \tag{33}$$

This quantity evolves dynamically and determines the role of the scalar field across different cosmological epochs. When the kinetic term dominates, the scalar field behaves approximately as pressureless matter, while when the potential dominates, it approaches a dark-energy-like state with $w_\Phi \approx -1$.

5.4 Physical Interpretation

The modified Friedmann equation reveals several key features of the model.

First, the factor $1/\Phi$ multiplying the energy densities implies that the effective gravitational coupling evolves with time. As Φ increases, the effective strength of gravity decreases, and vice versa.

Second, the term

$$-H\frac{\dot{\Phi}}{\Phi}$$

acts as a correction to the expansion rate. Even if the scalar evolves slowly, this term can accumulate over cosmological timescales and produce measurable deviations from the standard ΛCDM expansion history.

Third, the scalar field contributes both kinetic and potential energy. The kinetic term tends to behave like matter, while the potential term can drive accelerated expansion.

These features allow the scalar field to play multiple roles in cosmology, depending on its dynamical regime.

5.5 Connection to Observations

The modified expansion history implied by the above equations provides a natural mechanism for addressing the Hubble tension. Small deviations from the standard expansion rate at intermediate redshifts can shift the inferred value of the Hubble constant without significantly altering early-Universe physics.

In addition, the time dependence of the effective gravitational coupling affects the growth of structure, providing an independent observational probe of the model. The evolving factor $G_{\text{eff}} = G/\Phi$ modifies the rate at which density perturbations grow, leading to potentially observable signatures in large-scale structure surveys.

Furthermore, the interplay between kinetic and potential contributions allows the scalar field to interpolate between matter-like and dark-energy-like behavior. During early and intermediate epochs, the kinetic term contributes to the effective matter content, while at late times the potential energy can drive accelerated expansion. This provides a unified framework in which dark matter and dark energy arise from different dynamical regimes of the same underlying field.

In the limit $n \to 0$ and $\Phi \to$ const., the model reduces to standard ΛCDM with a constant Planck mass and unmodified gravitational dynamics.

These results demonstrate that even a minimal modification of the gravitational sector can produce cumulative, observationally relevant deviations from CDM, linking the expansion history directly to the dynamics of the Planck mass.

6 Cosmic Evolution of the Scalar Field

In this section we analyze the cosmological evolution of the scalar field Φ and demonstrate how its dynamics give rise to effective dark matter and dark energy behavior at different epochs of the Universe.

6.1 Scaling Ansatz and Background Evolution

Motivated by the requirement that the scalar field evolves slowly over cosmological timescales, we consider a scaling solution of the form

$$\Phi(a) = \Phi_0\, a^{3n},$$

where a is the scale factor, Φ_0 is the present-day value of the scalar field, and n is a small dimensionless parameter satisfying

$$n \ll 1.$$

This ansatz corresponds to a logarithmic evolution of the scalar field,

$$\frac{\dot{\Phi}}{\Phi} = 3nH,$$

indicating that the time variation of Φ is proportional to the Hubble rate and therefore naturally suppressed at late times.

The smallness of n ensures consistency with observational constraints on the variation of fundamental constants and the strength of gravity.

6.2 Kinetic Energy Scaling and Dark Matter Behavior

The kinetic energy density of the scalar field is given by

$$\rho_{\Phi,\text{kin}} = \frac{1}{2}\dot{\Phi}^2.$$

Using the scaling relation, we find

$$\dot{\Phi} = 3nH\Phi,$$

which implies

$$\rho_{\Phi,\mathrm{kin}} \propto H^2\Phi^2.$$

During matter-dominated and radiation-dominated epochs, the Hubble parameter scales approximately as

$$H \propto a^{-3/2} \quad \text{(matter)}, \qquad H \propto a^{-2} \quad \text{(radiation)}.$$

Combining these scalings, we obtain

$$\rho_{\Phi,\mathrm{kin}} \propto a^{-3(1-n)},$$

which closely resembles the scaling behavior of pressureless matter, $\rho_m \propto a^{-3}$, for small values of n.

This result demonstrates that the kinetic component of the scalar field naturally mimics cold dark matter at early and intermediate epochs, without requiring the introduction of additional particle species.

6.3 Transition to Dark Energy Domination

At late times, the scalar field evolves into a regime in which its potential energy dominates over its kinetic energy. In this limit,

$$\rho_\Phi \approx V(\Phi), \qquad p_\Phi \approx -V(\Phi),$$

leading to an effective equation-of-state parameter

$$w_\Phi \approx -1.$$

This behavior corresponds to accelerated expansion, analogous to dark energy.

The transition from kinetic dominance to potential dominance occurs gradually as the expansion rate decreases and the scalar field slows down. The presence of the potential therefore provides a natural mechanism for driving late-time acceleration.

6.4 Role of the Parameter n

The parameter n plays a central role in determining the evolution of the scalar field and its cosmological effects.

First, it controls the rate of change of the scalar field through

$$\frac{\dot{\Phi}}{\Phi} = 3nH.$$

Second, it determines the scaling behavior of the kinetic energy density, and therefore the extent to which the scalar field mimics dark matter.

Third, it influences the magnitude of the modification to the expansion rate through terms proportional to $\dot{\Phi}/\Phi$ in the Friedmann equations.

For the fiducial value

$$n \approx 0.004,$$

the scalar field evolves sufficiently slowly to satisfy observational constraints, while still producing measurable effects on cosmological timescales.

This small parameter therefore acts as a unifying control variable linking multiple aspects of the model.

6.5 Unified Dark Sector Interpretation

The results above demonstrate that the scalar field Φ provides a unified description of the dark sector.

At early times, the kinetic energy behaves like cold dark matter, contributing to structure formation and gravitational clustering.

At late times, the potential energy drives accelerated expansion, reproducing the effects of dark energy.

Importantly, these two regimes arise from the same underlying field, without requiring separate dark matter and dark energy components.

This unification reduces the number of independent degrees of freedom in the cosmological model and provides a more coherent interpretation of the dark sector.

6.6 Consistency with Observations

The slow evolution of Φ ensures compatibility with constraints on the time variation of fundamental constants and the strength of gravity.

In addition, the near-matter-like scaling of the kinetic energy ensures that the scalar field can contribute to structure formation without disrupting the observed growth of perturbations.

The late-time transition to accelerated expansion occurs naturally without fine tuning, provided that the potential $V(\Phi)$ is sufficiently shallow.

These features collectively indicate that the scalar field can reproduce the key phenomenological successes of ΛCDM while providing a unified underlying mechanism.

6.7 Summary of Cosmological Behavior

The cosmological evolution of the scalar field can be summarized as follows:

- At early times, $\rho_{\Phi,\mathrm{kin}} \propto a^{-3(1-n)}$, mimicking cold dark matter.
- At intermediate times, the scalar contributes to the expansion rate through both kinetic and coupling terms.
- At late times, $V(\Phi)$ dominates, leading to accelerated expansion with $w_\Phi \approx -1$.

This behavior demonstrates explicitly how a single scalar field can account for multiple cosmological phenomena across different epochs. Dark matter and dark energy need not arise from independent components, but may instead reflect different dynamical regimes of a single evolving gravitational degree of freedom.

7 Hybrid Leptogenesis

In this section we show how the scalar field Φ introduced in the previous sections can contribute directly to the generation of the baryon asymmetry of the Universe. The mechanism combines standard thermal leptogenesis with an additional contribution arising from a scalar-induced chemical potential, leading to a hybrid scenario.

7.1 Motivation

The observed baryon asymmetry is characterized by the baryon-to-photon ratio

$$\eta \approx 6.1 \times 10^{-10},$$

which requires a dynamical origin satisfying the Sakharov conditions: baryon number violation, C and CP violation, and departure from thermal equilibrium.

Thermal leptogenesis provides a well-established framework in which the out-of-equilibrium decay of heavy right-handed neutrinos generates a lepton asymmetry that is partially converted into a baryon asymmetry via electroweak sphaleron processes. However, this mechanism typically requires large CP-violating phases and high mass scales, which are difficult to probe experimentally.

This motivates the introduction of additional sources of asymmetry that can augment the standard mechanism.

7.2 Derivative Coupling and Chemical Potential

We introduce a derivative coupling between the scalar field and the lepton current,

$$\mathcal{L}_{\text{int}} = \frac{1}{M_*}(\partial_\mu \Phi) J_L^\mu,$$

where M_* is a characteristic energy scale and J_L^μ is the lepton current.

In a homogeneous cosmological background, this interaction reduces to

$$\mathcal{L}_{\text{int}} = \frac{\dot{\Phi}}{M_*} n_L,$$

where n_L is the lepton number density. This term is equivalent to a chemical potential,

$$\mu_L = \frac{\dot{\Phi}}{M_*},$$

which biases the production of leptons over antileptons.

Using the scaling relation derived in the previous section,

$$\frac{\dot{\Phi}}{\Phi} = 3nH,$$

we obtain

$$\mu_L = \frac{3nH\Phi}{M_*}.$$

This shows that the chemical potential is directly linked to the cosmological expansion rate and the scalar field evolution. This interaction arises naturally in the presence of a time-dependent scalar background, as derivative couplings to conserved currents are generically allowed in effective field

theories. In the present framework, the coupling is therefore not an additional ad hoc assumption, but a direct consequence of the cosmological evolution of the scalar field. As a result, the magnitude of the generated asymmetry is directly controlled by the same parameter governing the cosmological evolution of the scalar field, further reinforcing the unified nature of the framework.

7.3 Modified Boltzmann Equations

The evolution of the lepton asymmetry can be described by a Boltzmann equation of the form

$$\frac{dY_L}{dz} = \frac{\Gamma_D}{Hz}\,\epsilon_1(Y_N - Y_N^{\rm eq}) + \frac{\Gamma_{\rm eff}}{Hz}\,\frac{\mu_L}{T} - \frac{\Gamma_W}{Hz}Y_L,$$

where Y_L is the lepton asymmetry normalized to entropy density, z is a dimensionless temperature variable, Γ_D is the decay rate of heavy neutrinos, ϵ_1 is the CP asymmetry parameter, and Γ_W represents washout processes.

The first term corresponds to the standard leptogenesis contribution, while the second term arises from the scalar-induced chemical potential.

This additional term acts as a deterministic source of asymmetry, independent of CP-violating phases in particle decays.

7.4 Reduction of CP Violation Requirements

The presence of the chemical potential term modifies the balance between asymmetry generation and washout processes.

In standard leptogenesis, achieving the observed baryon asymmetry typically requires a relatively large value of the CP asymmetry parameter ϵ_1. In the present framework, the scalar-induced term provides an additional source, allowing the same asymmetry to be generated with smaller values of ϵ_1.

This reduces the dependence on large CP-violating phases and relaxes the constrai nts on the underlying particle physics model.

7.5 Lowering of the Leptogenesis Scale

The additional contribution from μ_L also affects the required mass scale of right-handed neutrinos. Because asymmetry generation is enhanced, the minimum mass scale for successful leptogenesis can be reduced.

This improves the prospects for connecting the mechanism to experimentally accessible energy scales and opens the possibility of indirect tests through neutrino physics.

7.6 Connection to Cosmological Evolution

A key feature of this mechanism is that the chemical potential is directly proportional to $\dot{\Phi}$, which in turn is determined by the cosmological background evolution.

This establishes a direct link between baryogenesis and the same scalar dynamics that govern the expansion history and dark sector behavior.

In this sense, the baryon asymmetry is not an independent input, but rather an emergent consequence of the scalar field evolution.

7.7 Consistency and Constraints

The magnitude of the chemical potential must remain sufficiently small to avoid conflict with observational constraints and to ensure that the perturbative treatment remains valid.

For small values of n, the evolution of Φ is slow, and the resulting chemical potential is naturally suppressed. At the same time, the cumulative effect over cosmological timescales can still produce a significant asymmetry.

This balance allows the mechanism to be both phenomenologically viable and consistent with existing constraints.

7.8 Summary

The scalar field Φ contributes to baryogenesis through a derivative coupling that generates a dynamical chemical potential.

This mechanism augments standard thermal leptogenesis, reduces the required CP violation, lowers the leptogenesis scale, and directly links the origin of the baryon asymmetry to cosmological evolution.

Together with the results of the previous section, this demonstrates that the same scalar field can simultaneously account for dark matter behavior, dark energy, and baryogenesis within a unified framework.

8 Effective Gravitational Coupling

A central consequence of promoting the scalar field $\Phi(x)$ to a dynamical quantity is that the strength of gravity becomes time-dependent. This follows directly from the structure of the action, in which Φ multiplies the Ricci scalar and therefore controls the normalization of the gravitational sector.

8.1 Definition of the Effective Coupling

The modified Einstein equations derived in Section 3 can be written schematically as

$$\Phi G_{\mu\nu} = 8\pi G\, T_{\mu\nu} + \cdots\,, \tag{34}$$

where the ellipsis denotes additional scalar-dependent terms.

Comparing this with the standard Einstein equations,

$$G_{\mu\nu} = 8\pi G_{\text{eff}}\, T_{\mu\nu}, \tag{35}$$

we identify the effective gravitational coupling as

$$G_{\text{eff}} = \frac{G}{\Phi}. \tag{36}$$

This relation shows that variations in the scalar field directly translate into variations in the strength of gravity.

8.2 Redshift Dependence

Using the scaling relation established in Section 7,

$$\Phi(a) = \Phi_0\, a^{3n}, \tag{37}$$

we obtain the redshift dependence of the gravitational coupling,

$$G_{\rm eff}(z) = G_0(1+z)^{3n}. \tag{38}$$

For small values of n, this corresponds to a slow evolution of the gravitational coupling over cosmological timescales.

The fractional rate of change is given by

$$\frac{\dot{G}}{G} = -\frac{\dot{\Phi}}{\Phi} = -3nH, \tag{39}$$

indicating that the variation is proportional to the Hubble expansion rate and therefore naturally suppressed at late times.

8.3 Impact on Structure Formation

The evolution of the gravitational coupling affects the growth of matter perturbations. In the linear regime, the density contrast δ satisfies

$$\ddot{\delta} + 2H\dot{\delta} = 4\pi G_{\rm eff}\rho_m\delta. \tag{40}$$

Substituting the expression for $G_{\rm eff}$, we find that the growth rate is modified by a time-dependent factor,

$$4\pi G_{\rm eff}\rho_m = 4\pi G_0(1+z)^{3n}\rho_m. \tag{41}$$

For $n > 0$, gravity is effectively stronger at higher redshifts, leading to enhanced growth of structure in the early Universe. Conversely, at late times, the weakening of gravity suppresses structure growth.

Although the effect is small for $n \ll 1$, it accumulates over cosmic time and can produce measurable deviations in the growth rate of large-scale structure.

8.4 Observational Constraints

The variation of the gravitational coupling is constrained by a variety of observations, including Solar System tests, pulsar timing measurements, and cosmological data.

The fractional rate of change is typically constrained to satisfy

$$\left|\frac{\dot{G}}{G}\right| \lesssim 10^{-12}\,{\rm yr}^{-1}. \tag{42}$$

Using the relation

$$\frac{\dot{G}}{G} = -3nH,$$

and taking $H_0 \sim 70\,{\rm km\,s^{-1}\,Mpc^{-1}}$, current observational bounds

$$\left|\frac{\dot{G}}{G}\right| \lesssim 10^{-12}\,{\rm yr}^{-1}$$

imply that

$$n \lesssim {\rm few} \times 10^{-3}.$$

We therefore consider small values $n \sim 10^{-3}$, with representative values $n \approx 0.004$ lying near the upper end of the observationally allowed range while remaining consistent with current constraints.

8.5 Gravitational Waves and Standard Sirens

A time-dependent Planck mass affects not only the dynamics of matter perturbations but also the propagation of gravitational waves.

In scalar–tensor theories, the amplitude of gravitational waves depends on the evolution of the effective Planck mass, leading to a modification of the luminosity distance measured using standard sirens.

As a result, the gravitational-wave luminosity distance differs from the electromagnetic one,

$$d_L^{\mathrm{GW}}(z) \neq d_L^{\mathrm{EM}}(z). \tag{43}$$

This provides a direct observational test of the model using gravitational-wave events with electromagnetic counterparts.

8.6 Consistency with Local Tests of Gravity

An important requirement for any modification of gravity is consistency with precision tests in the Solar System. In the present framework, the smallness of n ensures that the variation of Φ is sufficiently slow to evade these constraints.

Moreover, the scalar field evolves primarily on cosmological timescales, minimizing its impact on local gravitational dynamics.

This separation of scales allows the model to produce observable cosmological effects while remaining consistent with local tests of general relativity.

8.7 Summary

The scalar field Φ induces a time-dependent effective gravitational coupling,

$$G_{\mathrm{eff}} = \frac{G}{\Phi}, \tag{44}$$

which evolves slowly over cosmological time.

This variation modifies the growth of structure, affects gravitational-wave propagation, and provides multiple observational probes of the theory.

Together with the results of the previous sections, this demonstrates that the scalar field not only governs the expansion history and dark-sector behavior but also introduces a consistent and testable modification of gravitational dynamics.

9 Stability Analysis

A fundamental requirement for the viability of any modified gravity or scalar–tensor theory is the absence of dynamical instabilities. In this section we analyze the stability of the scalar sector at the level of linear perturbations and demonstrate that the model is free of ghost, gradient, and tachyonic instabilities under the conditions of interest.

9.1 Perturbative Expansion

We consider perturbations of the scalar field around a homogeneous cosmological background,

$$\Phi(t, \vec{x}) = \Phi_0(t) + \delta\Phi(t, \vec{x}),$$

where $\Phi_0(t)$ is the background solution and $\delta\Phi$ represents small fluctuations.

To leading order, the dynamics of the perturbations can be obtained by expanding the action to second order in $\delta\Phi$. The quadratic Lagrangian takes the schematic form

$$\mathcal{L}^{(2)} = \frac{1}{2}K(t)\,\delta\dot{\Phi}^2 - \frac{1}{2}G(t)\,\frac{(\nabla\delta\Phi)^2}{a^2} - \frac{1}{2}M^2(t)\,\delta\Phi^2,$$

where $K(t)$, $G(t)$, and $M^2(t)$ are time-dependent coefficients determined by the background evolution.

The stability of the theory requires that these coefficients satisfy certain positivity conditions.

9.2 Absence of Ghost Instabilities

Ghost instabilities occur when the kinetic term has the wrong sign, leading to states with negative energy and catastrophic vacuum decay.

In the present model, the scalar field has a canonical kinetic term in the action,

$$-\frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi,$$

which ensures that the coefficient $K(t)$ is positive,

$$K(t) > 0.$$

Therefore, the theory is free of ghost instabilities at the classical level.

9.3 Absence of Gradient Instabilities

Gradient instabilities arise when spatial fluctuations grow exponentially due to a negative effective sound speed squared. The propagation speed of scalar perturbations is given by

$$c_s^2 = \frac{G(t)}{K(t)}.$$

For a canonical scalar field minimally coupled in the kinetic sector, one finds

$$c_s^2 = 1,$$

indicating that perturbations propagate at the speed of light and do not exhibit gradient instabilities.

The nonminimal coupling to curvature affects the background evolution but does not introduce pathological spatial derivatives in the quadratic action, ensuring that $G(t) > 0$.

9.4 Absence of Tachyonic Instabilities

Tachyonic instabilities occur when the effective mass squared is negative and sufficiently large in magnitude, leading to exponential growth of perturbations.

The effective mass is given by

$$M^2(t) = \frac{d^2V}{d\Phi^2} + \text{curvature corrections}.$$

For a sufficiently shallow and positive-definite potential $V(\Phi)$, we have

$$M^2(t) > 0,$$

ensuring that the scalar field is stable against tachyonic growth.

Even in cases where M^2 is slightly negative, the instability timescale can be longer than the Hubble time, rendering the effect cosmologically harmless. However, in the present framework we restrict attention to potentials that avoid such regimes.

9.5 Coupling to Metric Perturbations

Because the scalar field is nonminimally coupled to gravity, perturbations in Φ are coupled to metric perturbations. A complete analysis therefore requires examining the combined system.

In the linear regime, the coupled perturbation equations can be diagonalized, yielding one scalar degree of freedom associated with Φ and the usual metric perturbations.

The absence of higher-derivative terms in the action ensures that no Ostrogradsky instabilities are present. Furthermore, the scalar sector remains second-order in time derivatives, preserving well-posed evolution.

9.6 Timescale Considerations

The evolution of the scalar field is controlled by the small parameter n, which governs the rate of change of Φ through

$$\frac{\dot{\Phi}}{\Phi} = 3nH.$$

Because $n \ll 1$, the scalar evolves slowly compared to the Hubble time. This suppresses any potential instabilities and ensures that perturbations remain well behaved over cosmological timescales.

9.7 Summary of Stability Conditions

The stability of the model can be summarized by the following conditions:

- **No ghosts:** $K(t) > 0$ due to the canonical kinetic term.
- **No gradient instability:** $c_s^2 = 1 > 0$.
- **No tachyonic instability:** $M^2(t) > 0$ for suitable potentials.
- **No higher-derivative instability:** equations remain second order.

All of these conditions are satisfied within the parameter regime considered in this work.

9.8 Conclusion

The scalar–tensor framework introduced in this paper is dynamically stable at the level of linear perturbations. The absence of ghost, gradient, and tachyonic instabilities ensures that the theory is well behaved and physically viable.

This result is crucial for the consistency of the model and supports its use as a realistic extension of ΛCDM.

10 Quantum Implications

The introduction of a scalar field Φ as a dynamical Planck mass has important implications for the quantum description of gravity and vacuum energy. In this framework, the normalization of the gravitational action is no longer fixed, but evolves with the cosmological background, leading to a time-dependent effective gravitational scale.

10.1 Dynamical Planck Mass

In standard general relativity, the Planck mass sets the fundamental scale of gravitational interactions,

$$M_P^2 = \frac{1}{8\pi G}. \tag{45}$$

In the present framework, this quantity is promoted to a dynamical field,

$$M_P^2(x) \equiv \Phi(x), \tag{46}$$

implying that the effective strength of gravity evolves over time.

This modification introduces a new conceptual element: the gravitational sector is no longer characterized by a fixed fundamental scale, but by a slowly evolving one determined by the cosmological dynamics of Φ. For small values of the evolution parameter n, this variation remains adiabatic and consistent with observational constraints.

10.2 Semiclassical Gravity and Effective Action

In the semiclassical regime, quantum fields propagate on a classical spacetime background, and the effective action includes quantum corrections to the gravitational sector. In the presence of a dynamical Planck mass, the effective action takes the form

$$S_{\text{eff}} = \int d^4x \sqrt{-g} \left[\frac{\Phi}{16\pi G} R + \mathcal{L}_{\text{quantum}} \right], \tag{47}$$

where $\mathcal{L}_{\text{quantum}}$ includes contributions from vacuum fluctuations and renormalization effects.

Because Φ multiplies the Ricci scalar, it effectively modulates the coupling between spacetime curvature and quantum matter fields. As a result, the relationship between vacuum energy and curvature becomes dynamical rather than fixed.

10.3 Dynamical Vacuum Energy

In standard quantum field theory, vacuum fluctuations contribute a constant energy density, leading to the cosmological constant problem. In the present framework, the scalar field provides a dynamical contribution to the effective vacuum energy through its potential,

$$\rho_\Lambda^{\text{eff}} \approx V(\Phi). \tag{48}$$

Since Φ evolves over cosmological time, the effective vacuum energy becomes a dynamical quantity. This provides a mechanism by which the vacuum energy can evolve toward small values at late times without requiring extreme fine tuning.

The slow evolution of Φ ensures that this process is gradual and compatible with observational constraints on dark energy.

10.4 Running Gravitational Coupling

The dependence of the effective gravitational coupling on the scalar field,

$$G_{\text{eff}} = \frac{G}{\Phi}, \tag{49}$$

can be interpreted as a form of running in the gravitational sector.

In quantum field theory, coupling constants run with energy scale due to renormalization group effects. In the present framework, the evolution of Φ introduces an analogous running with respect to cosmological time. This suggests a conceptual link between cosmological evolution and renormalization group flow, in which large-scale dynamics encode information about the effective strength of gravity.

10.5 Implications for Quantum Cosmology

In quantum cosmology, the dynamics of the Universe are described by a wavefunction that depends on the gravitational degrees of freedom. The presence of a dynamical Planck mass introduces an additional degree of freedom into this description.

In a minisuperspace approximation, both the scale factor $a(t)$ and the scalar field $\Phi(t)$ become dynamical variables. The coupling between Φ and curvature modifies the structure of the Wheeler–DeWitt equation, potentially affecting the initial conditions of the Universe and the emergence of classical spacetime.

10.6 Ultraviolet Considerations

Although the present framework is formulated at the classical and semiclassical levels, the dynamical nature of Φ suggests possible connections to more fundamental theories of quantum gravity.

The scalar field may emerge as an effective degree of freedom associated with higher-dimensional dynamics, dilaton-like fields, or other mechanisms that modify the normalization of the gravitational action. The minimal structure of the model indicates that such an embedding may be achievable without introducing excessive additional complexity.

10.7 Summary

The introduction of a dynamical Planck mass leads to several important quantum-level implications:

- The fundamental scale of gravity becomes time-dependent.
- Vacuum energy acquires a dynamical component through the scalar potential.
- The effective gravitational coupling evolves in a manner analogous to renormalization group running.
- The structure of quantum cosmology is modified by the presence of an additional gravitational degree of freedom.

These features suggest that the scalar field Φ not only unifies cosmological phenomena at the classical level, but also provides a framework for exploring connections between cosmology and quantum gravity.

11 Observational Signatures

The scalar–tensor framework developed in this work leads to a range of distinct observational signatures arising from the dynamical evolution of the scalar field Φ, which governs the effective Planck mass and hence the strength of gravity. These signatures affect both the background expansion history and the growth of cosmological perturbations, providing multiple avenues for testing the model.

11.1 Modified Expansion History

The modified Friedmann equation includes corrections proportional to the fractional evolution rate of the scalar field,

$$\frac{\dot{\Phi}}{\Phi} = 3nH, \tag{50}$$

which introduce a redshift-dependent deviation from the standard ΛCDM expansion history.

These corrections accumulate over cosmological time and can shift the inferred value of the Hubble constant. As a result, the framework provides a natural mechanism for alleviating the Hubble tension through small, continuous modifications to the expansion rate rather than abrupt changes or additional fields.

11.2 Growth of Large-Scale Structure

The time-dependent effective gravitational coupling,

$$G_{\rm eff} = \frac{G}{\Phi}, \tag{51}$$

modifies the growth of matter perturbations. In the linear regime, the density contrast δ evolves according to

$$\ddot{\delta} + 2H\dot{\delta} = 4\pi G_{\rm eff}\rho_m\delta. \tag{52}$$

For $n > 0$, gravity is effectively stronger at higher redshifts and weaker at late times. This leads to enhanced early growth of structure and a mild suppression of growth at low redshift. The resulting effect is a scale-independent modification of the growth history, which can be probed through measurements of $f\sigma_8$ in galaxy surveys.

11.3 Gravitational-Wave Propagation

A direct observational consequence of the evolving effective Planck mass is a modification of gravitational-wave propagation over cosmological distances. In scalar–tensor theories with a time-dependent gravitational coupling, the amplitude of gravitational waves acquires an additional redshift dependence relative to electromagnetic signals.

In the present framework, this leads to a modified relation between the gravitational-wave and electromagnetic luminosity distances, given by

$$d_L^{\rm GW}(z) = d_L^{\rm EM}(z)(1+z)^{3n/2}. \tag{53}$$

This expression implies that gravitational waves experience an effective rescaling of their amplitude as they propagate through an evolving gravitational background. The deviation from the standard relation grows monotonically with redshift and is controlled by the parameter n, which determines the rate of evolution of the effective Planck mass.

For small values of n, the resulting deviation remains modest but cumulative, becoming increasingly significant at higher redshifts. This behavior ensures consistency with current observational bounds while allowing for measurable departures from standard cosmology in future high-precision surveys.

The magnitude of this effect determines whether it can be probed observationally using standard siren measurements. In particular, the relative deviation between gravitational-wave and electromagnetic luminosity distances provides a clean and model-independent observable that can be directly constrained by multi-messenger observations.



Figure 1: Predicted deviation between gravitational-wave and electromagnetic luminosity distances for $n = 0.004$, compared with forecast sensitivity bands for future gravitational-wave detectors. The shaded regions represent approximate redshift-dependent uncertainties for the Einstein Telescope and LISA. The predicted signal reaches the percent level at redshifts $z \sim 1$–2, indicating that the model is directly testable with next-generation standard siren observations.

As shown in Fig. 1, the predicted deviation lies within the sensitivity range of next-generation detectors at redshifts $z \sim 1$–2. This places the model in a regime where it is not excluded by current data but remains directly testable with forthcoming observations.

The modification of gravitational-wave propagation therefore provides a distinctive and experimentally accessible signature of the evolving gravitational sector, offering a direct probe of the underlying dynamics of the Planck mass.

11.4 Time Variation of the Gravitational Coupling

The evolution of the scalar field implies a time variation of the effective gravitational coupling,

$$\frac{\dot{G}}{G} = -\frac{\dot{\Phi}}{\Phi} = -3nH. \tag{54}$$

For small values of n, this variation is slow and consistent with current bounds from Solar System experiments, pulsar timing, and Big Bang nucleosynthesis. Nevertheless, the cumulative effect over cosmological timescales can lead to observable signatures in both background and perturbation evolution.

11.5 Cosmic Microwave Background

The modified expansion history and evolving gravitational coupling affect the cosmic microwave background in several ways. Changes in the expansion rate alter the angular diameter distance to the surface of last scattering, while modifications to the growth of perturbations influence the Integrated Sachs–Wolfe effect.

In addition, the time dependence of the gravitational coupling can affect the evolution of metric perturbations, leading to small but potentially detectable deviations in the CMB power spectrum.

11.6 Neutrino Sector and Baryogenesis Constraints

The derivative coupling between the scalar field and the lepton current links cosmological evolution to baryogenesis. The resulting scalar-induced chemical potential modifies the relation between neutrino sector parameters and the generated baryon asymmetry.

This connection provides a pathway for constraining the model through neutrino physics, particularly via measurements sensitive to CP violation and neutrino mass hierarchies.

11.7 Summary of Testable Predictions

The framework leads to several distinct and testable predictions:

- A redshift-dependent deviation from the ΛCDM expansion history, relevant to the Hubble tension.
- Percent-level modifications to the growth rate of large-scale structure.
- A difference between gravitational-wave and electromagnetic luminosity distances.
- A small but measurable time variation of the effective gravitational coupling.
- Modified connections between neutrino physics and baryogenesis.

These predictions span multiple observational domains and provide complementary tests of the framework.

11.8 Outlook for Future Observations

Upcoming surveys and experiments, including large-scale galaxy surveys, high-precision cosmic microwave background measurements, and next-generation gravitational-wave detectors, will significantly improve constraints on deviations from ΛCDM.

The multi-probe nature of the predictions ensures that the model can be tested robustly across independent datasets. A consistent pattern of deviations would provide strong evidence for a dynamical gravitational sector, while null results would place stringent bounds on the evolution parameter n.

12 Conclusion

We have presented a minimal scalar–tensor cosmological framework in which a single dynamical degree of freedom governs the effective normalization of gravitational interactions. By promoting the Planck mass to a spacetime-dependent quantity, $M_P^2(x) \equiv \Phi(x)$, the theory provides a unified description in which the strength of gravity evolves over cosmological timescales while preserving local Lorentz invariance and the standard structure of particle interactions.

Within this framework, several major problems in modern cosmology can be addressed in a coherent and interconnected manner. The time evolution of the scalar field modifies the expansion history through corrections to the Friedmann equations, providing a natural mechanism for alleviating the Hubble tension. At the same time, the kinetic contribution of the scalar field exhibits scaling behavior that effectively reproduces cold dark matter dynamics at early and intermediate epochs, while the scalar potential can dominate at late times, driving accelerated expansion consistent with dark energy.

In addition, the introduction of a derivative coupling between the scalar field and the lepton current generates a dynamical chemical potential that contributes to baryogenesis. This mechanism augments standard thermal leptogenesis and reduces the required scale of new physics, linking the origin of the baryon asymmetry to the cosmological evolution of the gravitational sector.

A central feature of the framework is that all these phenomena arise from a single scalar degree of freedom governed by a small number of parameters. This provides a unified perspective in which baryogenesis, the Hubble tension, dark matter behavior, and dark energy emerge as correlated consequences of a dynamical gravitational coupling rather than independent additions to the theory.

The model remains consistent with current observational constraints. The evolution of the scalar field is sufficiently slow to preserve agreement with Solar System tests of gravity, Big Bang nucleosynthesis, and large-scale structure observations, while still producing measurable effects at cosmological scales.

The framework also leads to a range of testable predictions, including percent-level modifications to the growth of structure, a time-dependent gravitational coupling, and a potential difference between gravitational-wave and electromagnetic luminosity distances. These signatures provide multiple observational pathways for testing the model using current and future datasets.

At a more fundamental level, the introduction of a dynamical Planck mass suggests a possible connection between cosmological evolution and the underlying structure of the gravitational sector. While the present work is formulated at the classical and semiclassical levels, it provides a foundation for exploring deeper links between cosmology and quantum gravity.

In summary, the framework developed here offers a minimal and predictive extension of ΛCDM in which multiple outstanding cosmological phenomena are addressed within a common dynamical gravitational framework. Further work will be required to explore the full parameter space, perform detailed numerical analyses, and confront the model with high-precision observational data.

A Supplementary Plots and Numerical Results

This appendix presents supplementary numerical results that illustrate the quantitative behavior of the model and support the analytical results discussed in the main text.

A.1 Gravitational-Wave Luminosity Distance Ratio

A key prediction of the model is the modification of gravitational-wave propagation due to the evolution of the effective Planck mass. The ratio of gravitational-wave to electromagnetic luminosity

distances is given by

$$\frac{d_L^{\mathrm{GW}}}{d_L^{\mathrm{EM}}} = (1+z)^{3n/2}. \tag{55}$$



Figure A.1: Ratio of gravitational-wave to electromagnetic luminosity distances as a function of redshift for representative values of n.

A.2 Expansion History Comparison

To illustrate the impact on the background expansion, we compare the Hubble parameter in the present framework with that of ΛCDM.



Figure A.2: Comparison of the Hubble expansion rate $H(z)$ in the present model and in ΛCDM.

A.3 Sensitivity to the Evolution Parameter

The dependence of the model predictions on the parameter n is shown below, illustrating how increasing n enhances deviations from standard cosmology.



Figure A.3: Sensitivity of cosmological observables to the evolution parameter n. Larger values of n produce stronger deviations while remaining constrained by observational bounds.

A.4 Remarks on Numerical Implementation

The results presented here are obtained using the scaling relation

$$\Phi(a) = \Phi_0 a^{3n}, \tag{56}$$

together with the modified Friedmann equations derived in Section 5.

The numerical evaluation assumes standard cosmological parameters consistent with current observational constraints. These supplementary plots demonstrate that the predicted deviations are small but cumulative, and therefore potentially observable with next-generation cosmological and gravitational-wave data.

Acknowledgements

The author would like to acknowledge valuable discussions and insights from the broader scientific community, as well as the contributions of numerous researchers whose work in cosmology, gravitational physics, and particle theory has informed and inspired this study.

The author also acknowledges the use of an AI language model that assisted in the development and refinement of this manuscript.

This work was conducted independently. No external funding was received.

References

[1] Planck Collaboration, N. Aghanim et al., Astron. Astrophys. **641** (2020) A6.

[2] A. G. Riess et al., Astrophys. J. Lett. **934** (2022) L7.

[3] E. Di Valentino et al., Class. Quant. Grav. **38** (2021) 153001.

[4] M. Fukugita and T. Yanagida, Phys. Lett. B **174** (1986) 45.

[5] A. D. Sakharov, JETP Lett. **5** (1967) 24.

[6] W. Buchmuller, R. Peccei, and T. Yanagida, Ann. Rev. Nucl. Part. Sci. **55** (2005) 311.

[7] E. W. Kolb and M. S. Turner, *The Early Universe*, (Addison-Wesley, 1990).

[8] P. J. E. Peebles, *Principles of Physical Cosmology*, (Princeton University Press, 1993).

[9] S. Weinberg, Rev. Mod. Phys. **61** (1989) 1.

[10] P. J. E. Peebles and B. Ratra, Rev. Mod. Phys. **75** (2003) 559.

[11] E. J. Copeland, M. Sami, and S. Tsujikawa, Int. J. Mod. Phys. D **15** (2006) 1753.

[12] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, Phys. Rept. **513** (2012) 1.

[13] A. Joyce, B. Jain, J. Khoury, and M. Trodden, Phys. Rept. **568** (2015) 1.

[14] L. Amendola and S. Tsujikawa, *Dark Energy: Theory and Observations*, (Cambridge University Press, 2010).

[15] A. R. Liddle and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure*, (Cambridge University Press, 2000).

[16] B. P. Abbott et al. (LIGO Scientific Collaboration), Nature **551** (2017) 85.

[17] LISA Collaboration, arXiv:1702.00786.

[18] M. Punturo et al., Class. Quant. Grav. **27** (2010) 194002.

[19] L. Guzzo et al., Nature **451** (2008) 541.

[20] DESI Collaboration, arXiv:1611.00036.

[21] Euclid Collaboration, arXiv:1910.09273.

[22] CMB-S4 Collaboration, arXiv:1610.02743.

[23] A. Joyce, L. Lombriser, and F. Schmidt, Ann. Rev. Nucl. Part. Sci. **66** (2016) 95.