

Dissipative charging of bosonic quantum batteries via nonequilibrium gain and loss

A. E. B. Costa,¹ I. Beder,^{2,2} and P. A. Brandão^{2,*}

¹*Instituto Federal de Alagoas, Maceió, Brasil*

²*Instituto de Física, Universidade Federal de Alagoas, Maceió, Brasil*

(Dated: September 4, 2025)

We investigate the dissipative charging dynamics of a bosonic quantum battery interacting with two Markovian reservoirs: one inducing energy gain and the other inducing loss, both described by identical rates γ , thereby realizing a PT -symmetric dissipative configuration. When the battery is initially empty, we observe a monotonic increase of the internal energy over time t while the ergotropy remains strictly zero during the charging protocol. For initially populated Fock states with k excitations, the ergotropy decreases and vanishes for $\gamma t \geq k$, indicating a complete loss of extractable work despite continuous energy input. We further examine the role of initial quantum coherences in the battery state and find that XXXX. Our results highlight fundamental limitations on work extraction from dissipatively charged quantum systems, even under symmetric and energy-preserving environmental configurations.

I. INTRODUCTION

A quantum battery (QB) is an energy-storage device that, distinct from its classical counterpart, takes advantage of quantum phenomena, such as entanglement, to boost its performance [1, 2]. One of the main figures of merit characterizing the state of a QB is the ergotropy \mathcal{E} . It quantifies the amount of work (useful energy) that can be extracted from the QB by a unitary cyclic process [3]. The role of coherence and entanglement in work extraction processes has been the central topic of intense research in the past few years [] and a host of physical systems that could act as a QB have been proposed, including spin-based QB [], (...)

A charging protocol is usually considered in which the QB, initially empty, interacts with another (quantum or classical) system, called the charger, that holds some energy before the interaction takes place. This energy is transferred to the QB in the hope that it becomes extractable, that is, with nonzero ergotropy.

II. THEORY

A. Collision Model

Consider a QB composed of a single-mode harmonic oscillator having frequency ω and Hamiltonian $H_S = \omega b^\dagger b$, where b is the annihilation operator and $\hbar = 1$ hereafter. The QB interacts simultaneously with two atomic baths composed of excited and ground state two-level systems, described by the Hamiltonians H_G and H_L , respectively, as depicted in Fig. 1. Each bath is formed by discrete ancillas that interact with the QB during a collision time τ . The initial state of the total system is given by

$$\rho_{SB}(0) = \rho_S(0) \otimes \rho_G(0) \otimes \rho_L(0), \quad (1)$$

where B indicates the baths G and L and $\rho_j(0)$ ($j = S, G, L$) is the initial density operator of system j . Each ancilla of the G bath starts in the excited state $|e_G\rangle$, so that $\rho_G(0) = \bigotimes_n |e_G^n\rangle \langle e_G^n|$ and each ancilla of the L bath starts in the ground state $|g_L\rangle$, so that $\rho_L(0) = \bigotimes_n |g_L^n\rangle \langle g_L^n|$.




FIG. 1: A bosonic quantum battery, with Hamiltonian $H_S = \omega b^\dagger b$, (a single-mode cavity field) interacts with two atomic nonequilibrium baths. The gain bath consists of a sequence of excited atoms, described by the Hamiltonian H_G , which may increase photon excitations. The loss bath, characterized by the Hamiltonian H_L , is formed by ground state atoms that may absorb photonic excitations. The ancillas composing both baths do not interact between each other and are coupled to the quantum battery during a collision time τ .

The interaction energy between each individual ancilla and the QB is given by

$$H_{Sj}^n = g_j (b^\dagger \sigma_{j,n}^- + b \sigma_{j,n}^+) \quad (j = G, L), \quad (2)$$

where g_j is the coupling parameter between the QB and the reservoir $j = G, L$ and σ^\pm are the Pauli raising and lowering operators for the qubits. The collision model assumes a discrete time $t_n = n\tau$ for the interaction between the QB and the ancillas. That is, the pair of ancillas n , one from each bath, interacts with the QB in the time interval $[(n-1)\tau, n\tau]$. In this interval, the total system

* paulo.brandao@fis.ufal.br

evolves as $\rho_{SB}(n) = U(n)\rho_{SB}[(n-1)]U^\dagger(n)$, where $U(n)$ is the time-evolution operator

$$U(n) = e^{-i\tau(H_S + H_{G,n} + H_{L,n} + H_{SG}^n + H_{SL}^n)} \quad (3)$$

and $H_{j,n} = (\omega/2)\sigma_{j,n}^z$ ($j = G, L$) is the atomic energy of the n th qubit in bath j . Notice that we assume a resonant interaction between each ancilla and the QB. The density operator $\rho(n)$ for the QB after n interactions can be obtained by taking the partial trace $\rho(n) = \text{Tr}_B[\rho_{SB}(n)]$.

Under these assumptions (one collision between each ancilla and the QB, forbidden ancilla-ancilla interaction and uncorrelated initially prepared ancillas), it is known that this collision model generates a Markovian dynamics where each collision is fully characterized by a CPTP map [4].

B. Ergotropy

The ergotropy \mathcal{E} is an important figure of merit that characterizes the maximum amount of work that can be extracted from a given quantum state $\hat{\rho}$ of the QB under a cyclic and unitary transformation [3]. It is defined by

$$\mathcal{E} = \text{Tr}(\rho H) - \min_U \text{Tr}(U \rho U^\dagger H),$$

where the minimization is over all possible unitary transformations U . A more tractable formula for \mathcal{E} can be written in terms of the passive state σ of ρ , defined by

$$\sigma = \sum_{n=0} \lambda_n |n\rangle \langle n| \quad (4)$$

where λ_n are the eigenvalues of $\hat{\rho}$ rearranged in decreasing order, $\lambda_n > \lambda_{n+1}$ and $\{|n\rangle\}$ is the Hamiltonian eigenbasis. The ergotropy can thus be evaluated as

$$\mathcal{E} = \text{Tr}(\rho H) - \text{Tr}(\sigma H). \quad (5)$$

In the next section, we explore the role of the ergotropy \mathcal{E} and the energy E in the charging protocol of the QB interacting with the gain and loss atomic baths.

III. RESULTS AND DISCUSSION

A. One-cell quantum battery

Let us start by assuming that the QB is initially in the vacuum state $\hat{\rho}(0) = |0\rangle \langle 0|$. As discussed in the previous section, the energy of the battery will then increase during the charging protocol and we are interested in the fraction of this energy that can be extracted as work, given by Eq. (5). The inset of Fig. 2 shows the energy E (gray line) and the ergotropy \mathcal{E} (red line) as a function of γt . It is clear from these plots that the energy increases but the ergotropy remains zero, so that no work can be extracted after the charging protocol is completed.




FIG. 2: Charging of a bosonic QB that is initially in the vacuum state $\hat{\rho}(0) = |0\rangle \langle 0|$. The plot displays the populations $\rho_n(t) = \langle n | \hat{\rho}(t) | n \rangle$ as a function of the occupation number n for $\gamma t = 0$ (gray dots), $\gamma t = 7.5$ (blue dots) and $\gamma t = 15$ (purple dots). The inset shows the energy E and ergotropy \mathcal{E} in units of ω as a function of γt .

To explain this behavior, we focus on the evolution of the populations $\rho_n(t) = \langle n | \hat{\rho}(t) | n \rangle$. Since the initial state of the QB is diagonal in the energy basis of the Hamiltonian, the eigenvalues of $\hat{\rho}$ are identified with the populations ρ_n . Figure 2 shows the populations ρ_n as a function of the mode number n for fixed charging times $\gamma t = 0$ (gray dots), $\gamma t = 0.75$ (blue dots) and $\gamma t = 1.5$ (purple dots). Initially, only the mode $n = 0$ is excited, as expected. As t increases, there is a redistribution of populations throughout excited states ($n > 0$), but, as is evident from the plots, the distribution follows an exponentially decreasing function of n , which suggests that the state of the quantum battery evolves to a thermal-like state distribution. Thus, a non-Hermitian-induced thermalization precludes the extraction of work. The same behavior was observed for the case $\gamma_G > \gamma_L$, except that the modes ρ_n are populated faster than the PT -symmetric case.

The exact thermal distribution can be obtained directly from Eq. (??) by fixing $k = 0$ and expanding $M_k(\mu, t)$ in a power series of $(1 - \mu)^n$. The populations are given by

$$\rho_n^{(k=0)}(t) = \frac{(\gamma t)^n}{(1 + \gamma t)^{n+1}}, \quad (6)$$

which describes a thermal distribution having mean photon number $\bar{n} = \gamma t$, which can be directly obtained from Eq. (??) by taking $\gamma_L = \gamma_G = \gamma$ and $\bar{n}(0) = 0$. Thus, the passive state is obviously $\hat{\sigma} = \rho_n^{(k=0)}$ and the ergotropy vanishes at all times if the QB is initially in the vacuum state.




FIG. 3: caption

We see that the energy deposited in the QB by the PT-symmetric reservoir is useless because it creates thermal-like states with zero ergotropy. What happens then if the QB has initially some energy and ergotropy? To explore this situation, let us consider the simplest case where the QB initially holds a single photon, that is, $\hat{\rho}(0) = |1\rangle\langle 1|$. Equation (??) can then be used with $k = 1$ to extract the populations,

$$\rho_n^{(k=1)}(t) = \left(\gamma t + \frac{n}{\gamma t} \right) \frac{(\gamma t)^n}{(1 + \gamma t)^{n+2}}. \quad (7)$$

Figure 3 plots the ergotropy \mathcal{E} as a function of γt . The plot clearly suggests an exponential *discharging* dynamics (in the sense that the ergotropy is being drained during the time period), even though the energy is increasing. Note, however, that this is not technically an exponential behavior for all times. The ergotropy exactly vanishes at $\gamma t_c = 1$. This can be seen by inspecting the plots of the populations and searching for the time t_c where the curves for ρ_0 and ρ_1 cross each other. The time t_c also happens to be the time where the ergotropy vanishes.

The exact time t_c for which $\mathcal{E}(t_c) = 0$ can be obtained directly from Eq. (??) by expanding $M_k(\mu, t)$ in powers of $(1 - \mu)^n$ and extracting the $n = 0$ and $n = 1$ coefficients.

They are given by the populations

$$\rho_0^{(k)}(t) = \frac{(\gamma t)^k}{(1 + \gamma t)^{k+1}} \quad (8)$$

and

$$\rho_1^{(k)}(t) = \frac{(\gamma t)^k}{(1 + \gamma t)^{k+1}} [\beta(k+1) - \alpha k]. \quad (9)$$

It is a simple exercise to demonstrate that the equation $\rho_0^{(k)}(t_c) = \rho_1^{(k)}(t_c)$ is satisfied only if $\gamma t_c = k$. Thus, if the battery initially holds k photons, the ergotropy will last for a time k (in units of γ).

It seems that the interaction between the QB and the PT-symmetric environment is detrimental to the performance of the QB in the sense that it always decreases the initial ergotropy (if any). So far, we have ignored the role of possible initial coherences present between the quantum states of the QB. To explore this scenario, let us assume that the initial state of the battery is given by

$$\hat{\rho}(0) = a |0\rangle\langle 0| + (1-a) |1\rangle\langle 1| + p |0\rangle\langle 1| + p^* |1\rangle\langle 0|, \quad (10)$$

where a is a positive number and p is the coherence parameter satisfying $|p|^2 \leq a(1-a) := f(a)$. The presence of nondiagonal terms modify the passive state $\hat{\sigma}$ of the QB and, consequently, the ergotropy.




FIG. 4: caption




FIG. 5: Colormap of ergotropy as a function of a and $|p|^2$ for $\gamma t = 2$. The red line indicates the upper limit of $|p|^2$ imposed by the positivity condition of the density matrix.

IV. CONCLUSIONS

ACKNOWLEDGMENTS

We wish to acknowledge the financial support of CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior).

-
- [1] R. Alicki and M. Fannes, Entanglement boost for extractable work from ensembles of quantum batteries, *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics* **87**, 042123 (2013).
 - [2] F. Campaioli, S. Gherardini, J. Q. Quach, M. Polini, and G. M. Andolina, Colloquium: quantum batteries, *Reviews of Modern Physics* **96**, 031001 (2024).
 - [3] A. E. Allahverdyan, R. Balian, and T. M. Nieuwenhuizen, Maximal work extraction from finite quantum systems, *Europhysics Letters* **67**, 565 (2004).
 - [4] F. Ciccarello, S. Lorenzo, V. Giovannetti, and G. M. Palma, Quantum collision models: Open system dynamics from repeated interactions, *Physics Reports* **954**, 1 (2022).