

GEOMETRY

Through Algebra

Errala Paulsonashish



Contents

Introduction	i
1 Triangle	1
1.1 Matrix	1
1.1.1 Vectors	2
1.1.2 Median	5
1.1.3 Altitude	10
1.1.4 Perpendicular Bisector	11
1.1.5 Angle Bisector	13

Introduction

This book shows how to solve problems in geometry using trigonometry and coordinate geometry.

Chapter 1

Triangle

Consider a triangle with vertices

$$\mathbf{A} = \begin{pmatrix} -5 \\ -4 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 3 \\ -3 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} 4 \\ 0 \end{pmatrix} \quad (1.1)$$

1.1. Matrix

The matrix of vertices of the triangle is defined as

$$\mathbf{P} = \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \quad (1.2)$$

$$= \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \quad (1.3)$$




Figure 1.1: $\triangle ABC$

1.1.1. Vectors

1.1.1.1. Obtain the direction matrix of the sides of $\triangle ABC$ defined as

$$M = \begin{pmatrix} \mathbf{A} - \mathbf{B} & \mathbf{B} - \mathbf{C} & \mathbf{C} - \mathbf{A} \end{pmatrix} \quad (1.1.1.1.1)$$

Solution:

$$\mathbf{M} = \begin{pmatrix} \mathbf{A} - \mathbf{B} & \mathbf{B} - \mathbf{C} & \mathbf{C} - \mathbf{A} \end{pmatrix} \quad (1.1.1.1.2)$$

$$= \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix} \quad (1.1.1.1.3)$$

$$= \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{pmatrix} \quad (1.1.1.1.4)$$

Using Matrix multiplication

$$\mathbf{M} = \begin{pmatrix} -8 & -1 & 9 \\ -1 & -3 & 4 \end{pmatrix} \quad (1.1.1.1.5)$$

where the second matrix above is known as a circulant matrix. Note that the 2^{nd} and 3^{rd} row of the above matrix are circular shifts of the 1^{st} row.

1.1.1.2. Obtain the normal matrix of the sides of $\triangle \mathbf{ABC}$

Solution:

Considering the rotation matrix

$$\mathbf{R} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad (1.1.1.2.1)$$

the normal matrix is obtained as

$$\mathbf{N} = \mathbf{R}\mathbf{M} \quad (1.1.1.2.2)$$

$$= \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -8 & -1 & 9 \\ -1 & -3 & 4 \end{pmatrix} \quad (1.1.1.2.3)$$

Using matrix multiplication

$$\mathbf{N} = \begin{pmatrix} 1 & 3 & -4 \\ -8 & -1 & 9 \end{pmatrix} \quad (1.1.1.2.4)$$

1.1.1.3. Obtain $\mathbf{a}, \mathbf{b}, \mathbf{c}$.

Solution:

The sides vector is obtained as

$$\mathbf{d} = \sqrt{\text{diag}(\mathbf{M}^\top \mathbf{M})} \quad (1.1.1.3.1)$$

$$\mathbf{M}^\top \mathbf{M} = \begin{pmatrix} -8 & -1 \\ -1 & -3 \\ 9 & 4 \end{pmatrix} \begin{pmatrix} -8 & -1 & 9 \\ -1 & -3 & 4 \end{pmatrix} \quad (1.1.1.3.2)$$

Using matrix multiplication

$$\mathbf{M}^\top \mathbf{M} = \begin{pmatrix} 65 & 11 & -76 \\ 11 & 10 & -21 \\ -76 & -21 & 97 \end{pmatrix} \quad (1.1.1.3.3)$$

$$\mathbf{d} = \sqrt{\text{diag} \left(\begin{pmatrix} 65 & 11 & -76 \\ 11 & 10 & -21 \\ -76 & -21 & 97 \end{pmatrix} \right)} \quad (1.1.1.3.4)$$

$$= \begin{pmatrix} \sqrt{65} & \sqrt{10} & \sqrt{97} \end{pmatrix} \quad (1.1.1.3.5)$$

1.1.1.4. Obtain the constant terms in the equations of the sides of the triangle.

Solution:

The constants for the lines can be expressed in vector form as

$$\mathbf{c} = \text{diag} \{ (\mathbf{N}^\top \mathbf{P}) \} \quad (1.1.1.4.1)$$

$$\mathbf{N}^\top \mathbf{P} = \begin{pmatrix} 1 & -8 \\ 3 & -1 \\ -4 & 9 \end{pmatrix} \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \quad (1.1.1.4.2)$$

$$(1.1.1.4.3)$$

Using matrix multiplication

$$= \begin{pmatrix} 27 & 27 & 4 \\ -11 & 12 & 12 \\ -16 & -39 & -16 \end{pmatrix} \quad (1.1.1.4.4)$$

$$\mathbf{c} = \text{diag} \left(\begin{pmatrix} 27 & 27 & 4 \\ -11 & 12 & 12 \\ -16 & -39 & -16 \end{pmatrix} \right) \quad (1.1.1.4.5)$$

$$= \begin{pmatrix} 27 & 12 & -16 \end{pmatrix} \quad (1.1.1.4.6)$$

1.1.2. Median

1.1.2.1. Obtain the midpoint matrix for the sides of the triangle

Solution:

$$\begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad (1.1.2.1.1)$$

$$= \frac{1}{2} \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix} \quad (1.1.2.1.2)$$

Using matrix multiplication

$$\begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} = \begin{pmatrix} \frac{7}{2} & -\frac{1}{2} & -1 \\ -\frac{3}{2} & -2 & -\frac{7}{2} \end{pmatrix} \quad (1.1.2.1.3)$$




Figure 1.2: mid-points

1.1.2.2. Obtain the median direction matrix.

Solution:

The median direction matrix is given by

$$\mathbf{M}_1 = \begin{pmatrix} \mathbf{A} - \mathbf{D} & \mathbf{B} - \mathbf{E} & \mathbf{C} - \mathbf{F} \end{pmatrix} \quad (1.1.2.2.1)$$

$$= \begin{pmatrix} \mathbf{A} - \frac{\mathbf{B}+\mathbf{C}}{2} & \mathbf{B} - \frac{\mathbf{C}+\mathbf{A}}{2} & \mathbf{C} - \frac{\mathbf{A}+\mathbf{B}}{2} \end{pmatrix} \quad (1.1.2.2.2)$$

$$= \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & 1 & -\frac{1}{2} \\ -\frac{1}{2} & -\frac{1}{2} & 1 \end{pmatrix} \quad (1.1.2.2.3)$$

$$= \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & 1 & -\frac{1}{2} \\ -\frac{1}{2} & -\frac{1}{2} & 1 \end{pmatrix} \quad (1.1.2.2.4)$$

Using matrix multiplication

$$\mathbf{M}_1 = \begin{pmatrix} -\frac{17}{2} & \frac{7}{2} & 5 \\ -\frac{5}{2} & -1 & \frac{7}{2} \end{pmatrix} \quad (1.1.2.2.5)$$

1.1.2.3. Obtain the median normal matrix.

Solution:

Considering the rotation matrix

$$\mathbf{R} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad (1.1.2.3.1)$$

the normal matrix is obtained as

$$\mathbf{N}_1 = \mathbf{R}\mathbf{M}_1 \quad (1.1.2.3.2)$$

$$= \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -\frac{17}{2} & \frac{7}{2} & 5 \\ -\frac{5}{2} & -1 & \frac{7}{2} \end{pmatrix} \quad (1.1.2.3.3)$$

$$\mathbf{N}_1 = \begin{pmatrix} \frac{5}{2} & 1 & -\frac{7}{2} \\ -\frac{17}{2} & \frac{7}{2} & 5 \end{pmatrix} \quad (1.1.2.3.4)$$

1.1.2.4. Obtain the median equation constants.

$$\mathbf{c}_1 = \text{diag} \left(\left(\mathbf{N}_1^\top \begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} \right) \right) \quad (1.1.2.4.1)$$

$$\mathbf{N}_1^\top \begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} = \begin{pmatrix} \frac{5}{2} & -\frac{17}{2} \\ 1 & \frac{7}{2} \\ -\frac{7}{2} & 5 \end{pmatrix} \begin{pmatrix} \frac{7}{2} & -\frac{1}{2} & -1 \\ -\frac{3}{2} & -2 & -\frac{7}{2} \end{pmatrix} \quad (1.1.2.4.2)$$

Using matrix multiplication

$$= \begin{pmatrix} \frac{43}{2} & \frac{63}{4} & \frac{109}{4} \\ -\frac{7}{4} & -\frac{15}{2} & -\frac{53}{4} \\ -\frac{79}{4} & -\frac{33}{4} & -14 \end{pmatrix} \quad (1.1.2.4.3)$$

$$\mathbf{c}_1 = \text{diag} \left(\begin{pmatrix} \frac{43}{2} & \frac{63}{4} & \frac{109}{4} \\ -\frac{7}{4} & -\frac{15}{2} & -\frac{53}{4} \\ -\frac{79}{4} & -\frac{33}{4} & -14 \end{pmatrix} \right) \quad (1.1.2.4.4)$$

$$\mathbf{c}_1 = \begin{pmatrix} \frac{43}{2} & -\frac{15}{2} & -14 \end{pmatrix} \quad (1.1.2.4.5)$$

1.1.2.5. Obtain the centroid by finding the intersection of the medians.

Solution:

$$\left(\mathbf{N}_1^\top \mid \mathbf{c}^\top \right) = \left(\begin{array}{cc|c} \frac{5}{2} & -\frac{17}{2} & \frac{43}{2} \\ 1 & \frac{7}{2} & -\frac{15}{2} \\ -\frac{7}{2} & 5 & -14 \end{array} \right) \quad (1.1.2.5.1)$$

Using Gauss-Elimination method:

$$\left(\begin{array}{cc|c} \frac{5}{2} & -\frac{17}{2} & \frac{43}{2} \\ 1 & \frac{7}{2} & -\frac{15}{2} \\ -\frac{7}{2} & 5 & -14 \end{array} \right) \xleftrightarrow{R_1 \leftarrow \frac{2R_1}{5}} \left(\begin{array}{cc|c} 1 & -\frac{17}{5} & \frac{43}{5} \\ 1 & \frac{7}{2} & -\frac{15}{2} \\ -\frac{7}{2} & 5 & -14 \end{array} \right) \quad (1.1.2.5.2)$$

$$\xleftrightarrow{R_2 \leftarrow R_2 - R_1} \left(\begin{array}{cc|c} 1 & -\frac{17}{5} & \frac{43}{5} \\ 0 & \frac{69}{10} & -\frac{161}{10} \\ -\frac{7}{2} & 5 & -14 \end{array} \right) \quad (1.1.2.5.3)$$

$$\xleftrightarrow{R_3 \leftarrow R_3 + \frac{7R_1}{2}} \left(\begin{array}{cc|c} 1 & -\frac{17}{5} & \frac{43}{5} \\ 0 & \frac{69}{10} & -\frac{161}{10} \\ 0 & -\frac{69}{10} & \frac{161}{10} \end{array} \right) \quad (1.1.2.5.4)$$

$$\xleftrightarrow{R_2 \leftarrow \frac{10R_2}{69}} \left(\begin{array}{cc|c} 1 & -\frac{17}{5} & \frac{43}{5} \\ 0 & 1 & -\frac{7}{3} \\ 0 & -\frac{69}{10} & \frac{161}{10} \end{array} \right) \quad (1.1.2.5.5)$$

$$\xleftrightarrow{R_1 \leftarrow R_1 + \frac{17R_2}{5}} \left(\begin{array}{cc|c} 1 & 0 & \frac{2}{3} \\ 0 & 1 & -\frac{7}{3} \\ 0 & -\frac{69}{10} & \frac{161}{10} \end{array} \right) \quad (1.1.2.5.6)$$

$$\xleftrightarrow{R_3 \leftarrow R_3 + \frac{69R_2}{10}} \left(\begin{array}{cc|c} 1 & 0 & \frac{2}{3} \\ 0 & 1 & -\frac{7}{3} \\ 0 & 0 & 0 \end{array} \right) \quad (1.1.2.5.7)$$

$$\text{Therefore } \mathbf{G} = \left(\begin{array}{c} \frac{2}{3} \\ -\frac{7}{3} \end{array} \right) \quad (1.1.2.5.8)$$




Figure 1.3: centroid of triangle ABC

1.1.3. Altitude

1.1.3.1. Find the normal matrix for the altitudes

Solution: The desired matrix is

$$\mathbf{M}_2 = \begin{pmatrix} \mathbf{B} - \mathbf{C} & \mathbf{C} - \mathbf{A} & \mathbf{A} - \mathbf{B} \end{pmatrix} \quad (1.1.3.1.1)$$

$$= \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{pmatrix} \quad (1.1.3.1.2)$$

$$= \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \begin{pmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{pmatrix} \quad (1.1.3.1.3)$$

Using Matrix multiplication

$$\mathbf{M}_2 = \begin{pmatrix} -1 & 9 & -8 \\ -3 & 4 & -1 \end{pmatrix} \quad (1.1.3.1.4)$$

1.1.3.2. Find the constant vector for the altitudes.

Solution:

The desired vector is

$$\mathbf{c}_2 = \text{diag} \{ (\mathbf{M}^\top \mathbf{P}) \} \quad (1.1.3.2.1)$$

$$\mathbf{M}^\top \mathbf{P} = \begin{pmatrix} -1 & -3 \\ 9 & 4 \\ -8 & -1 \end{pmatrix} \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \quad (1.1.3.2.2)$$

$$(1.1.3.2.3)$$

Using matrix multiplication

$$\mathbf{M}^\top \mathbf{P} = \begin{pmatrix} 17 & 6 & -4 \\ -61 & 15 & 36 \\ 44 & -21 & -32 \end{pmatrix} \quad (1.1.3.2.4)$$

$$\mathbf{c}_2 = \text{diag} \left(\begin{pmatrix} 17 & 6 & -4 \\ -61 & 15 & 36 \\ 44 & -21 & -32 \end{pmatrix} \right) \quad (1.1.3.2.5)$$

$$\mathbf{c}_2 = \begin{pmatrix} 17 & 15 & -32 \end{pmatrix} \quad (1.1.3.2.6)$$




Figure 1.4: Orthocentre of $\triangle ABC$

1.1.4. Perpendicular Bisector

1.1.4.1. Find the normal matrix for the perpendicular bisectors

Solution:

The normal matrix is \mathbf{M}_2

$$\mathbf{M}_2 = \begin{pmatrix} -1 & 9 & -8 \\ -3 & 4 & -1 \end{pmatrix} \quad (1.1.4.1.1)$$




Figure 1.5: plot of perpendicular bisectors

1.1.4.2. Find the constants vector for the perpendicular bisectors.

Solution: The desired vector is

$$\mathbf{c}_3 = \text{diag} \left\{ \mathbf{M}_2^\top \begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} \right\} \quad (1.1.4.2.1)$$

Solution:

$$\mathbf{c}_3 = \text{diag} \left\{ \mathbf{M}_2^\top \begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} \right\} \quad (1.1.4.2.2)$$

$$\mathbf{M}_2^\top \begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} = \begin{pmatrix} -1 & -3 \\ 9 & 4 \\ -8 & -1 \end{pmatrix} \begin{pmatrix} \frac{7}{2} & -\frac{1}{2} & -1 \\ -\frac{3}{2} & -2 & -\frac{7}{2} \end{pmatrix} \quad (1.1.4.2.3)$$

$$(1.1.4.2.4)$$

Using matrix multiplication

$$\mathbf{M}_2^\top \begin{pmatrix} \mathbf{D} & \mathbf{E} & \mathbf{F} \end{pmatrix} = \begin{pmatrix} 1 & \frac{13}{2} & \frac{23}{2} \\ \frac{51}{2} & -\frac{25}{2} & -23 \\ -\frac{5}{32} & 6 & \frac{23}{2} \end{pmatrix} \quad (1.1.4.2.5)$$

$$\mathbf{c}_3 = \text{diag} \left(\begin{pmatrix} 1 & \frac{13}{2} & \frac{23}{2} \\ \frac{51}{2} & -\frac{25}{2} & -23 \\ -\frac{5}{32} & 6 & \frac{23}{2} \end{pmatrix} \right) \quad (1.1.4.2.6)$$

$$\mathbf{c}_3 = \begin{pmatrix} 1 & -\frac{25}{2} & \frac{23}{2} \end{pmatrix} \quad (1.1.4.2.7)$$




Figure 1.6: circumcentre and circumcircle of $\triangle ABC$

1.1.5. Angle Bisector

1.1.5.1. Find the points of contact.

Solution:

The points of contact are given by

$$\begin{pmatrix} \frac{n\mathbf{A}+p\mathbf{C}}{n+p} & \frac{p\mathbf{B}+m\mathbf{A}}{p+m} & \frac{m\mathbf{C}+n\mathbf{B}}{m+n} \end{pmatrix} = \begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} \frac{n}{b} & \frac{m}{c} & 0 \\ 0 & \frac{p}{c} & \frac{n}{a} \\ \frac{p}{b} & 0 & \frac{m}{a} \end{pmatrix} \quad (1.1.5.1.1)$$

$$\begin{pmatrix} \mathbf{p} & \mathbf{m} & \mathbf{n} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} \end{pmatrix} \begin{pmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{pmatrix} \quad (1.1.5.1.2)$$

$$= \frac{1}{2} \begin{pmatrix} \sqrt{10} & \sqrt{97} & \sqrt{65} \end{pmatrix} \begin{pmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{pmatrix} \quad (1.1.5.1.3)$$

$$= \frac{1}{2} \begin{pmatrix} 3.16227766 & 9.84885780 & 8.062257748 \end{pmatrix} \begin{pmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{pmatrix} \quad (1.1.5.1.4)$$

Using matrix multiplication

$$\begin{pmatrix} \mathbf{p} & \mathbf{m} & \mathbf{n} \end{pmatrix} = \begin{pmatrix} 7.374418944 & 0.687838804 & 2.474438856 \end{pmatrix}$$

(1.1.5.1.5)

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} & \mathbf{C} \end{pmatrix} \begin{pmatrix} \frac{n}{b} & \frac{m}{c} & 0 \\ 0 & \frac{p}{c} & \frac{n}{a} \\ \frac{p}{b} & 0 & \frac{m}{a} \end{pmatrix} = \begin{pmatrix} -5 & 3 & 4 \\ -4 & -3 & 0 \end{pmatrix} \begin{pmatrix} \frac{2.474438856}{\sqrt{97}} & \frac{0.687838804}{\sqrt{65}} & 0 \\ 0 & \frac{7.374418944}{\sqrt{65}} & \frac{2.474438856}{\sqrt{10}} \\ \frac{7.374418944}{\sqrt{97}} & 0 & \frac{0.687838804}{\sqrt{10}} \end{pmatrix} \quad (1.1.5.1.6)$$

Using matrix multiplication We get the points of contact

$$= \begin{pmatrix} 1.7388292 & 2.31747277 & 3.21751372 \\ -1.0049648 & -3.0853159 & -2.34745882 \end{pmatrix} \quad (1.1.5.1.7)$$




Figure 1.7: Contact points of incircle of *triangle* ABC




Figure 1.8: Incircle and Incentre of $\triangle ABC$