

d

The Dirac Axioms for Quantization

It is well established that classical mechanics has its mathematical formalisms in the context of symplectic and Poisson geometry via Hamiltonian mechanics, and quantum mechanics has its mathematical formalisms in the context of Hilbert spaces via the Dirac-von Neumann axioms for quantum mechanics, which unifies the wave-function approach of Schrödinger with the operator approach of Heisenberg.

To elaborate, the Hamiltonian view of classical mechanics postulates that, given a physical system, there is a *configuration space* given by a smooth manifold Q that represents the space of possible states of the physical system; we may then obtain its *phase space* by considering its cotangent bundle, T^*Q , which carries a natural symplectic structure¹. Observables of this physical system are given by smooth functions $T^*Q \rightarrow \mathbf{R}$; the space of smooth functions $C^\infty(T^*Q)$ forms a Poisson algebra. From this function space, there is a distinguished² observable H called the *Hamiltonian* that representing the energy of the system; Hamilton's equations then state that the physical dynamics of the system are governed by the equation

$$\frac{f}{t} = \{f, H\}$$

for $f \in C^\infty(T^*Q)$ an observable quantity, and $\{\cdot, \cdot\}$ the standard Poisson bracket on $C^\infty(T^*Q)$. For further details and a more comprehensive treatment, see the classic text by Arnol'd³; see also the related appendix.

On the other hand, the mathematical formulation of quantum mechanics (following the Dirac-von Neumann axioms) calls for a Hilbert space \mathcal{H} to represent the space of states of a physical quantum system⁴, with observables given by self-adjoint operators $O : \mathcal{H} \rightarrow \mathcal{H}$. Similar to the classical case, the collection of all such observables forms an algebra, albeit a non-commutative one. Dynamics are determined by Schrödinger's equation

$$i\hbar \frac{\psi}{t} = H\psi$$

or equivalently by the Heisenberg equation

$$\frac{O}{t} = \frac{i}{\hbar} [H, O]$$

depending on whether one considers the states to be time-varying or the observables to be time-varying. Here, H is a distinguished operator on the Hilbert space called the Hamiltonian, O is an observable,

¹ Additionally, we may consider a *reduced* phase space if there is an relevant Lie group acting on the phase space via symplectomorphisms — i.e. via “physical symmetries”. This, and the construction of a quantum analogue of the classical reduction process, is considered in a later chapter.

² (Due primarily to its importance in a physical context rather than any intrinsic mathematical property.)

³

⁴ Actually, we may also consider the projectivization $\mathbf{P}\mathcal{H}$ of this Hilbert space as the space of states, since we have to account for equivalence modulo scalar multiples — $\lambda \cdot |\psi\rangle$ represents the same state as $|\psi\rangle$, for any complex nonzero λ .

and $[\cdot, \cdot]$ is the commutator bracket on the algebra of observables. Detailed discussions on the mathematical formulations of quantum mechanics may be found in von Neumann's classic text ⁵ as well as in the relevant appendix.

It is natural to seek a way of relating these two frameworks. Historically, the development of the theory of quantization began with Dirac, who introduced (what is now called) canonical quantization for the purpose of explaining quantum phenomena by way of a "classical analogy" in his doctoral thesis ⁶; in particular, the correspondence between the Poisson bracket and the commutator is developed. In Dirac's later text ("*Principle of Quantum Mechanics*", ⁷), the axiomatic framework for quantization was proposed.

In general, we have what are known as the *Dirac axioms* for quantization: for a symplectic manifold (M, Ω) , a quantization \mathcal{Q} of M should give us an associated Hilbert space \mathcal{H} as well as an association between smooth functions $C^\infty(M) \ni f : M \rightarrow \mathbb{R}$ and self-adjoint operators $\mathcal{Q}(f) = \mathcal{O}_f : \mathcal{H} \rightarrow \mathcal{H}$ such that the following properties hold:

Definition 1. • *Poisson brackets correspond to commutators, i.e. $\mathcal{Q}(\{f, g\}) = [\mathcal{Q}(f), \mathcal{Q}(g)]$;*

- *$\mathcal{Q}(1) = i \cdot \text{Id}$, i.e. the constant function 1 should be mapped to the identity operator times the scalar $\sqrt{-1}$;*
- *for $\alpha, \beta \in \mathbb{C}$, $\mathcal{Q}(\alpha f + \beta g) = \alpha \mathcal{Q}(f) + \beta \mathcal{Q}(g)$ (Linearity); and*
- *a Minimality Condition: any complete family of functions maps to a complete family of operators.*

These four ideas correspond to intuitive points regarding the structures we might wish to see preserved through the quantization. The first point states, essentially, that Lie-algebraic structures should be preserved; the second point states that the trivial classical observable should correspond to the trivial quantum observable⁸; the third point states that quantization is a linear map; and the last point, the minimality condition, states that complete families of functions (i.e., families of functions that separate points on the manifold M) should correspond to families of operators (i.e., operators that act irreducibly on \mathcal{H}).

⁸ ...So that "non-observation" is quantized to "non-observation".

Additionally, physical intuition stipulates that the position and momentum functions $x^i : M \rightarrow \mathbb{R}$ and $p_j : M \rightarrow \mathbb{R}$ must be quantized to the operators $\psi \mapsto x^i \psi$ and $\psi \mapsto -i\hbar_j \psi$, respectively; we shall see, however, that these conditions are rarely a point of difficulty during the quantization process and serve mainly as a guide. The important structural content of a quantization scheme is codified mainly in the above four axioms.

Unfortunately, it is not possible to quantize the entire Poisson algebra of smooth functions, and there are many obstructions against quantization; see *Obstruction Results in Quantization Theory*⁹ for a survey. In particular, we have the *Groenewald-van Hove theorem*, which informally states that the algebra of polynomials on the phase space \mathbb{R}^{2n} has no unitary representation ρ such that ρ extends the Schrödinger representation of the Heisenberg algebra. Consequently, this provides a counterexample to the claim that it is possible to quantize the entire algebra of observables on a classical phase space. This obstruction, as well as others, are discussed in the chapter dedicated to obstruction results in the theory of quantization.

Some of the more common approaches to quantization are *canonical* (first) and *second* quantization. The former refers to a semiclassical quantization of mechanical systems, where the potentials are treated classically and particles are quantized into elements of a Hilbert space of functions (for example, an L^2 space); the latter then refers to a functor from the category of Hilbert spaces (i.e., state spaces of single-particle physical systems) to a category of Fock spaces (a direct sum of tensor products of Hilbert spaces, representing state spaces of physical systems that may have multiple particles).

The current survey will not discuss either of those approaches; standard references such as^{10, 11} or¹² may be consulted for their mathematical developments and references. Instead, the current survey will present *geometric* quantization and *deformation* quantization, for two main reasons: first, the mathematical techniques involved in these two theories are less understood, and the scope of their prerequisites often clouds their intuitive structural properties, which form a rich theory that attempts to preserve as much geometric and algebraic insight as possible; and second, because much active work is being done on furthering these geometric and algebraic approaches

9

10

11

12

—

To motivate the development of the geometric and deformation theories of quantization, we note that the basic intuition behind these theories is to deal with the inability to satisfy all of Dirac's axioms¹³ by either compromising on the correspondence of the Poisson bracket and the commutator, or by limiting the observables that we are "allowed" to quantize. Deformation quantization does the former by treating \hbar as a formal parameter that "adjusts" the bracket, allowing us to obtain Poisson brackets in the "limit" $\hbar \rightarrow 0$. Geometric quantization does the latter by preserving geometric structure through the quantization and choosing a structure (called a *polarization*) on the quantized geometry that limits which operators are able to be meaningfully quantized.

¹³ In fact, it is provable that it is only possible to satisfy two of them at most!