

1. Find whether the following pair of linear equations are consistent or inconsistent:

$$5x - 3y = 11, -10x + 6y = 22.$$

2. Solve for x and y :

$$x + y = 6, 2x - 3y = 4.$$

3. Find out whether the pair of equations $2x + 3y = 0$ and $2x - 3y = 26$ is consistent or inconsistent.
4. For what values of k , does the pair of linear equations $kx - 2y = 3$ and $3x + y = 5$ have a unique solution?
5. What type of lines will you get by drawing the graph of the pair of equations $x - 2y + 3 = 0$ and $2x - 4y = 5$?
6. The sum of the numerator and the denominator of a fraction is 18. If the denominator is increased by 2, the fraction reduces to $\frac{1}{3}$. Find the fraction.
7. Find the value of k for which the system of equations $x + 2y = 5$ and $3x + ky + 15 = 0$ has no solution.
8. If 2 tables and 2 chairs cost ₹ 700 and 4 tables and 3 chairs cost ₹ 1,250, then find the cost of one table.
9. If the graph of a pair of lines $x - 2y + 3 = 0$ and $2x - 4y = 5$ be drawn, then what type of lines are drawn?

10. $\begin{bmatrix} 12 \\ 34 \end{bmatrix}$

11. If $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$, then A^2 equals

(a) $\begin{bmatrix} 2 & -2 \\ -2 & 2 \end{bmatrix}$

(b) $\begin{bmatrix} 2 & -2 \\ -2 & -2 \end{bmatrix}$

(c) $\begin{bmatrix} -2 & -2 \\ -2 & 2 \end{bmatrix}$

(d) $\begin{bmatrix} -2 & 2 \\ 2 & -2 \end{bmatrix}$

12. $\begin{vmatrix} 43 & 44 & 45 \\ 44 & 45 & 46 \\ 45 & 46 & 47 \end{vmatrix}$

- (a) 0

(b) -1

(c) 1

(d) 2

13. A square matrix A is said to be singular if _____.

If $A = \begin{bmatrix} 3 & -5 \\ 2 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 17 \\ 0 & -10 \end{bmatrix}$, then $|AB| = \underline{\hspace{2cm}}$.

14. If $\begin{bmatrix} 4 & x+2 \\ 2x-3 & x+1 \end{bmatrix}$ is a symmetric matrix, then find the value of x .

If A is a square matrix such that $A^2 = A$, then find $(2 + A)^3 - 19A$.

15. For matrix $A = \begin{bmatrix} 2 & 3 \\ -4 & -6 \end{bmatrix}$, verify the following: $A(adj A) = (adj A)A = |A|I$

16. Using properties of determinants show that $\begin{vmatrix} 1+a^2-b^2 & 2ab & -2b \\ 2ab & 1-a^2 & 2a \\ 2b & -2a & 1-a^2-b^2 \end{vmatrix} = (1+a^2+b^2)^3$

17. Find the equation of the line joining $A(1, 3)$ and $B(0, 0)$, using determinants. Also, find k if $D(k, 0)$ is a point such that the area of $\triangle ABD$ is 3 square units.

18. Solve the system of linear equations, using the matrix method:

$$7x + 2y = 114x - 7 = 2$$

19. Find the Value of x , if $\begin{bmatrix} x & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix} \begin{bmatrix} x \\ 3 \end{bmatrix} = 0$

20. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, then $A^4 = \underline{\hspace{2cm}}$.

21. Given $A = \begin{bmatrix} 1-1 & 1 & \\ 3 & -2 & 1 \\ -2 & 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 1 & -2 \end{bmatrix}$, the order of the matrix AB is _____.

22. if $A = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$ ($i^2 = -1$) and $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$, then AB is equal to

(a) $\begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}$

(b) $\begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}$

(c) $\begin{bmatrix} i & -i \\ 0 & 1 \end{bmatrix}$

(d) $\begin{bmatrix} 0 & 0 \\ i & 0 \end{bmatrix}$

23. If A is a $5 \times p$ matrix, B is a $2 \times q$ matrix, then the order of the matrix AB is 5×4 . What are the values of p and q ?
- (a) $p = 2, q = 4$
 - (b) $p = 4, q = 2$
 - (c) $p = 2, q = 2$
 - (d) $p = 4, q = 4$
24. Value of k , for which $A = \begin{bmatrix} k & 8 \\ 1 & 2k \end{bmatrix}$ is a singular matrix is:
- (a) 4
 - (b) -4
 - (c) ± 4
 - (d) 0
25. If $A = [a_{ij}]$ is a square matrix of order 2 such that $a_i = \begin{cases} 1, \text{ when } i + j \\ 0, \text{ when } i - j \end{cases}$, then A^2 is:
- (a) $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$
 - (b) $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$
 - (c) $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$
 - (d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
26. Given that A is a square matrix of order 3 and $|A| = -4$, then $|adj A|$ is equal to:
- (a) -4
 - (b) 4
 - (c) -16
 - (d) 16
27. If $\begin{bmatrix} 2a+b & a-2b \\ 5c-d & 4c+3d \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 11 & 24 \end{bmatrix}$, then the value of $a + b - c + 2d$ is:
- (a) 8
 - (b) 10
 - (c) 4
 - (d) -8

28. Given that matrices A and B are of order $3 \times n$ and $m \times 5$ respectively, then the order of matrix $C = 5A + 3B$ is:
- 3×5
 - 5×3
 - 3×3
 - 5×5
29. For matrix $A = \begin{bmatrix} 2 & 5 \\ -11 & 7 \end{bmatrix}$, $(\text{adj}A)'$ is equal to:
- $\begin{bmatrix} -2 & -5 \\ 11 & -7 \end{bmatrix}$
 - $\begin{bmatrix} 7 & 5 \\ 11 & 2 \end{bmatrix}$
 - $\begin{bmatrix} 7 & 11 \\ -5 & 2 \end{bmatrix}$
 - $\begin{bmatrix} 7 & -5 \\ 11 & 2 \end{bmatrix}$
30. Given that $A = [a_{ij}]$ is a square matrix of order 3×3 and $|A| = -7$, then the value of $\sum_{i=1}^3 a_{i2}A_{i2}$, where A_{ij} denotes the cofactor of element a_{ij} is:
- 7
 - 7
 - 0
 - 49
31. If $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -4 \\ 2 & -1 & 5 \end{bmatrix}$, then
- $A^{-1} = B$
 - $A^{-1} = 6B$
 - $B^{-1} = B$
 - $B^{-1} = \frac{1}{6}A$
32. Given that A is a non-singular matrix of order 3 such that $A^2 = 2A$, then the value of $|2A|$ is:
- 4
 - 8
 - 64
 - 16

33. If $A = \begin{bmatrix} 0 & 2 \\ 3 & -4 \end{bmatrix}$ and $kA = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$, then the values of k, a and b respectively are:

- (a) $-6, -12, -18$
- (b) $-6, -4, -9$
- (c) $-6, 4, 9$
- (d) $-6, 12, 18$

34. If A is square matrix such that $A^2 = A$, then $(I + A)^3 - 7A$ is equal to:

- (a) A
- (b) $I + A$
- (c) $I - A$
- (d) I

35. For $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, then 14^{-1} is given by:

- (a) $14 \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$
- (b) $\begin{bmatrix} 4 & -2 \\ 2 & 6 \end{bmatrix}$
- (c) $2 \begin{bmatrix} 2 & -1 \\ 1 & -3 \end{bmatrix}$
- (d) $2 \begin{bmatrix} -3 & -1 \\ 1 & -2 \end{bmatrix}$

36. Given that $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ and $A^2 = 3I$, then:

- (a) $1 + \alpha^2 + \beta\gamma = 0$
- (b) $1 - \alpha^2 - \beta\gamma = 0$
- (c) $3 - \alpha^2 - \beta\gamma = 0$
- (d) $3 + \alpha^2 + \beta\gamma = 0$

- (a) $|A| = 0$
- (b) $|A| \in (2, \infty)$
- (c) $|A| \in (2, 4)$
- (d) $|A| \in [2, 4]$