

# EE5609: Matrix Theory

## Assignment-10

M Pavan Manesh  
EE20MTECH14017

**Abstract**—This document explains regarding the use of Cayley-Hamilton theorem

Download all latex-tikz codes from

<https://github.com/pavanmanesh/EE5609/tree/master/Assignment10>

### 1 PROBLEM

For the linear operator  $\mathbf{T}$

$$\mathbf{T} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 3x_1 \\ x_1 - x_2 \\ 2x_1 + x_2 + x_3 \end{pmatrix} \quad (1.0.1)$$

Prove that

$$(\mathbf{T}^2 - I)(\mathbf{T} - 3I) = 0 \quad (1.0.2)$$

### 2 SOLUTION

Expressing (1.0.1) in matrix form

$$\mathbf{T} = \begin{pmatrix} 3 & 0 & 0 \\ 1 & -1 & 0 \\ 2 & 1 & 1 \end{pmatrix} \quad (2.0.1)$$

The characteristic equation of  $\mathbf{T}$  is given as follows,

$$|\mathbf{T} - \lambda \mathbf{I}| = \begin{vmatrix} 3 - \lambda & 0 & 0 \\ 1 & -1 - \lambda & 0 \\ 2 & 1 & 1 - \lambda \end{vmatrix} = 0 \quad (2.0.2)$$

$$\begin{aligned} \implies (3 - \lambda)(-1 - \lambda)(1 - \lambda) &= 0 \\ \implies (\lambda - 3)(1 + \lambda)(1 - \lambda) &= 0 \\ \implies (\lambda - 3)(1 - \lambda^2) &= 0 \\ \implies (\lambda^2 - 1)(\lambda - 3) &= 0 \end{aligned} \quad (2.0.3)$$

By the Cayley-Hamilton theorem, We can write (2.0.3) as

$$(\mathbf{T}^2 - I)(\mathbf{T} - 3I) = 0 \quad (2.0.4)$$