Name	:	**Mrs. BHARATHI K JOSHI**			
Age/Gender	:	63 Years / Female	Vial ID	:	2950084
Ref.By	:	Self	Collected On	:	03-Feb-2026 02:52 PM
Req No.	:	**BNG2611421**	Registered On	:	03-Feb-2026 02:40 PM
Sample Type	:	Serum	Reported On	:	03-Feb-2026 06:11 PM
Client Name	:	VMEDO RETAIL-CMLKAD56	Client Code	:	CMLKAD56

C-Reactive Protien (CRP) (Nephlometry)

Test Name	Observed Values	Units	Biological Reference Intervals
* C-REACTIVE PROTIEN(CRP) Method: Immunoturbidimetric	3.69	mg/L	Up to 6.0

INTERPRETATION:

- C - Reactive Protein (CRP) is an acute phase protein that is involved in the activation of complement, acceleration of phagocytosis and detoxification of substances released from damaged tissue. As such, CRP is considered to be one of the most sensitive indicators of inflammation. In response to an inflammatory stimulus, a rise in CRP may be detected with 6 hours. CRP is sensitive, though considered to be a non-specific indicator of acute phase reactants.
- Measurement of C - Reactive Protein is most frequently used for the evaluation of injury to body tissue or for the detection of an inflammatory event somewhere in the body. CRP levels in serum are typically elevated in patients with arthritis or liver diseases such as Hepatitis A, Hepatitis B or biliary cirrhosis and after severe infections such as septic shock.

Dr Faeeza Begum MD,
KMC.No:103490
MD PATHOLOGY

-------------- End Of The Report --------------

Name	: **Mrs. BHARATHI K JOSHI**		
Age/Gender	: 63 Years / Female	Vial ID	: 2950083
Ref.By	: Self	Collected On	: 03-Feb-2026 02:52 PM
Req No.	: **BNG2611421**	Registered On	: 03-Feb-2026 02:40 PM
Sample Type	: WB-EDTA	Reported On	: 03-Feb-2026 03:17 PM
Client Name	: VMEDO RETAIL-CMLKAD56	Client Code	: CMLKAD56

Complete Blood Count (CBC)

Test Name	Observed Values	Units	Biological Reference Intervals
* HAEMOGLOBIN Method:Colorimetric	12.3	gm%	12.0 - 15.0
* RBC Count Method:Electrical Impedence	4.56	millions/cumm	4.50 - 5.50
* WBC Count Method:Electrical Impedence	9100	cells/cumm	4000-11000
* PLATELET COUNT Method:Impedence and Light Scattering	2.53	lakhs/cumm	1.5 - 4.5
* Packed Cell Volume(PCV) Method:Cum.RBC Pulse High detection	38.4	%	40 - 50
* MCV Method:Calculated	88	fL	80 - 100
* MCH Method:Calculated	28.0	pg	27 - 33
* M C H C Method:Calculated	34.0	g/dL	32.0 - 34.0
DIFFERENTIAL COUNT			
* NEUTROPHILS,, Method:Manual - Leishman Stain & Smear Microscopy	75	%	50-75
* LYMPHOCYTES Method:Flowcytometry	20	%	20 - 40
* EOSINOPHILS Method:Manual - Leishman Stain & Smear Microscopy	03	%	02 - 06
* MONOCYTES Method:Flowcytometry	02	%	02 - 08
* BASOPHILS Method:Manual - Leishman Stain & Smear Microscopy	00	%	0 - 1

Dr Faeeza Begum MD,
KMC.No:103490
MD PATHOLOGY

-------------- End Of The Report --------------

Name	: **Mrs. BHARATHI K JOSHI**		
Age/Gender	: 63 Years / Female	Vial ID	: 2950084
Ref.By	: Self	Collected On	: 03-Feb-2026 02:52 PM
Req No.	: **BNG2611421**	Registered On	: 03-Feb-2026 02:40 PM
Sample Type	: Serum	Reported On	: 03-Feb-2026 06:11 PM
Client Name	: VMEDO RETAIL-CMLKAD56	Client Code	: CMLKAD56

Kidney Basic Screen

Test Name	Observed Values	Units	Biological Reference Intervals
* SERUM CREATININE Method:Enzymatic Method (sarcosine oxidase, Peroxidase)	0.83	mg/dL	0.6 - 1.2
* UREA Method:Urease UV	20.0	mg/dL	16.8-43.2
* UREA NITROGEN (BUN) Method:Calculated	9.35	mg/dL	8 - 23
* BUN/CREATININE RATIO	11.26		
* URIC ACID Method:Uricase-Peroxidase	3.46	mg/dL	Male:3.6 -8.2 Female:2.3 - 6.1
* SODIUM Method:ISE Direct	140	mmol/L	136-145
* POTASSIUM Method:ISE Direct	**3.19**	mmol/L	3.5 - 5.5
* CHLORIDE Method:ISE Direct	102	mmol/L	95-105

INTERPRETATION::

kidney disease usually does not have signs or symptoms. Testing is the only way to know how your kidneys are doing. It is important to get checked for kidney disease if an individual having the key risk factors - diabetes, high blood pressure, heart disease, or a family history of kidney failure.

Dr Faeeza Begum MD,
KMC.No:103490
MD PATHOLOGY

-------------- End Of The Report --------------