

Comparison chart

Written by [Pavel Sobolev](#) and located [here](#)

Used abbreviations:

KIC – ID from Kepler Input Catalog;

KID – Kernel Identifier;

KL – Kernel Link (for *george*);

CNL – Notebook Link (*celerite*);

GNL – Notebook Link (*george*);

TP – 'True' value of the period of a star [days];

IP – Inferred value of the period obtained by
minimizing the negative marginalized likelihood [days];

RCL – Result Link (PDF, *celerite*);

RGL – Result Link (PDF, *george*)

KIC	KID	KL	CNL	GNL	TP	IP	RCL	RGL
1430163	Base	link	link	link	3.88 ± 0.58	≈ 3.7897	link	link
	DS	link	link	link	3.88 ± 0.58	≈ 3.8751	link	link

Base

$$k(\tau) = \frac{a}{2+b} e^{-\tau/c} \left[\cos\left(\frac{2\pi\tau}{P}\right) + (1+b) \right]$$

Parameters:

a \rightarrow log_amp;

b \rightarrow log_factor;

c \rightarrow log_timescale;

P \rightarrow log_period

Bounds:

log_amp: (-10.0, 0.0);

log_timescale: (1.5, 5.0);

log_period: (-3.0, 5.0);

log_factor: (-5.0, 5.0)

MCMC (george):

nwalkers: 8;

nburn: 50;

nsteps: 500

MCMC (celerite):

nwalkers: 32;

nburn: 500;

nsteps: 5000

Execution time:

Sigma clipping: 10m 18s;

MCMC sampling: 7h 50m 15s

Execution time:

Sigma clipping: 7.46s;

MCMC sampling: 1m 56.7s

DS

$$k(\tau) = e^{-a\tau} \cos\left(\frac{2\pi\tau}{p}\right)$$

Parameters:

$a \rightarrow \log_a$;

$p \rightarrow \log_p$

MCMC (george):

nwalkers: 6;

nburn: 50;

nsteps: 500

Execution time:

Sigma clipping: 5m 16s;

MCMC sampling: 10h 3m 32s

Bounds:

\log_a : (-5.0, -1.5);

\log_p : (-3.0, 5.0);

MCMC (celerite):

nwalkers: 32;

nburn: 500;

nsteps: 5000

Execution time:

Sigma clipping: 6.20s;

MCMC sampling: 1m 29.5s