


BIOXPLO^R LTD

INTELLECTUAL PROPERTY ACQUISITION OPPORTUNITY

Bioinformatics software, biomedical literature search technologies, ontology-assisted analysis and structured data assets



CONFIDENTIAL PRODUCT PROPOSITION

AUGUST 2026

Prepared by Pavels Kartasevs, CTO of BioXplor Ltd

High-level acquisition overview

OPPORTUNITY OVERVIEW

A portfolio of biomedical search and analysis technologies

BioXplor Ltd developed software for drug indication expansion, biomedical literature analysis and processing of biomedical terms identified in research publications.

TRANSACTION CONTEXT

Due to the company's financial position and to support an orderly closure, BioXplor Ltd management intends to sell the company's intellectual property portfolio.

INDICATIVE ASKING PRICE

Approximately EUR 10,000 for the intellectual property portfolio, subject to buyer review and a definitive agreement.

Portfolio at a glance

01 Internal scientific literature tool Searches for specific scientific terms and identifies direct co-occurrence and indirect relationships among two, three or more terms.	02 Indication expansion platform Finds literature from user queries, applies ontology-based co-occurrence filters, searches at sentence level and presents visualisations.
03 Structured databases Includes BioXplor Ltd's remaining databases of co-occurring scientific terms and literature sentences.	04 Publication text and image search Searches publication sentences, full text and figures, including image-text, category and similarity-based retrieval.

Core capabilities represented in the portfolio

- Biomedical literature discovery and sentence-level retrieval
- Scientific term co-occurrence and indirect-relationship analysis
- Ontology-assisted filtering and interactive visualisation
- Full-text, sentence and publication-figure search

TECHNOLOGY ASSET 02

Indication expansion platform

The indication expansion technology combines user-defined literature queries with ontology co-occurrence filtering, sentence-level search and visual presentation of results.

QUERY	FILTER	ANALYSE	VISUALISE
User-defined literature search	Ontology co-occurrence filtering	Sentence-level evidence discovery	Interactive result presentation

Available product versions

VERSION	SCOPE
Academic tool	General academic indication expansion and biomedical literature exploration.
COVID-19 tool	A version focused on COVID-19-related literature and terminology.
OncoExplorer	An oncology-focused version with literature filtering and relationship visualisation.
Product tool	Adds the ability to identify, search and display products mentioned in scientific literature.

Representative interface: academic tool




Figure 1. Academic indication expansion interface with term labels, result counts and filter controls.

Scientific term analysis, structured databases and publication search

01	Internal scientific literature tool Finds specific terms in scientific literature and supports both direct co-occurrence and indirect relationship discovery. Searches may involve two, three or more scientific terms.
03	BioXplor Ltd databases Includes the company's remaining databases of co-occurring terms and literature sentences.
04	Publication sentence, text and image search Searches publication sentences and text, and retrieves publication images using image text, image category and image similarity.

[bloxplor](#)

🔍 https://bloxplor.com/search

🏠 📁 ⌕ 🔖 ⚙️

Close

Serum Interleukin 17 Levels in Patients with Crohn's Disease: Real Life Data

► Authors

▼ Contents

- ▶ Protein (1)
- ▶ Gene (10)
- ▶ product (1)
- ▼ Druggable-Proteins (1)
 - kit
- ▼ Disease (91)
 - CD (38)
 - anti-TNF (7)
 - UC (5)
 - albumin (5)
 - Albuminuria
 - Albuminuria and retinopathy
 - Albuminuria and uraemia
 - AUR
 - ALU
 - Albuminuria
 - ALI
 - Albuminuria increased
 - Albuminuria reduction
 - Albuminuria in HIV-infected
 - Alb
 - ALB
 - Albumin-deficient
 - Albumin deficiency
 - A-LI
 - AA
 - AD
 - Albumin dialysis
 - AAS
 - AL
 - AN
 - AR
 - AS
 - Ab-T
 - Ab-T

Serum Interleukin 17 Levels in Patients with Crohn's Disease: Real Life Data

1. Introduction

Crohn's disease (CD) is a chronic relapsing inflammatory disease affecting the gastrointestinal tract and presenting with extraintestinal manifestations as well. Although the etiologyogenesis of this disease is not completely understood, it seems to be influenced by several environmental factors in genetically predisposed individuals [1, 2]. T-helper lymphocytes (Th) play an important role in the pathogenesis of CD. Crohn's disease is postulated to be an immunoregulatory disease mediated by Th1 cells and recently also by Th17 cells [3]. Th17 cells are major contributors to several autoimmune diseases that were previously thought to be Th1-cell predominant diseases. These autoimmune diseases can be listed as rheumatoid arthritis, systemic lupus, spondyloarthritis, multiple sclerosis, ulcerative colitis, idiopathic thrombocytopenic purpura, autoimmune myocarditis, and endometriosis [4]. Th17 cells also play an important role in maintaining intestinal barrier function by affecting innate and adaptive responses. Mucosal Th17 cells prevent migration of pathogens from breaking mucosa to the systemic circulation through the lamina propria of neutrophils and macrophages. One of the important features of Th17 cells is to balance mucosal inflammation by regulating the immunogenic response against self-antigens or intestinal pathogens due to their relationship with regulatory T cells. The other is plasticity, the ability of these cells to differentiate to other T cell subgroups under various types of stimulation. Th17 cells that are induced *in vivo* by transforming growth factor α (TGFα) and interleukin (IL)-6 alone produce both IL-17 and the inhibitory cytokine IL-22 and do not induce autoimmunity, whereas Th17 cells that have been stimulated with IL-23 after stimulation with TGFα and IL-6 produce other proinflammatory cytokines and chemokines, but not IL-17, and are involved in inflammation [5]. The cytokines secreted by Th17 cells are IL-1, IL-2, IL-23, IL-26, tumor necrosis factor α, interferon γ, and mainly IL-17. IL-17 is produced by T cells, macrophages, dendritic cells, natural killer, and other T cells. Thereby, IL-17 influences several types of immune cells, particularly B cells and antigen-presenting cells, to induce the expression of several genes. IL-17 has been shown to stimulate the production of matrix metalloproteinases [6, 7, 8]. The role of Th17 cells and IL-17 cytokine in CD is not completely understood, and data on this issue has been obtained from animal models and tissue sample studies from inflamed mucosa [9–11]. On the other hand, the IL-17 level in sera of CD patients has not yet been studied. In the present study, IL-17 levels of CD patients were investigated. The relation of IL-17 levels with disease activity and other proinflammatory markers was also evaluated.

2. Materials and Methods

2.1. Patients

Fifty patients diagnosed with CD and 40 controls were included in the study between January 2012 and August 2012. Demographic characteristics, disease features (location, behavior, disease duration), and extraintestinal manifestations, medications, and disease activity of CD patients were recorded. Disease activity in CD patients was assessed by the Crohn's disease activity index (CDAI) [12]. A score under 150 was accepted as inactive disease, while a CDAI score over 150 was considered as active disease [12]. The medications were classified as "tumor necrosis factor blocker (anti-TNF) group," "corticosteroids without immunomodulatory and without mesalazine," "without immunomodulatory and without corticosteroids," "with immunomodulatory group (immunomodulatory drug without corticosteroids and without mesalazine)," and "mesalazine group (mesalazine without corticosteroids). The control group consisted of 40 age- and sex-matched subjects coming to the hospital with dyspeptic complaints, who had no any inflammatory disease or disorder with a characteristic of changed IL-17 levels were excluded from the control group. Pregnancy and ages younger than 18 years old were other exclusion criteria for patients and controls. The protocol was approved by the local ethics committee. Informed consent was obtained from all subjects before obtaining samples. Samples of peripheral blood were allowed to clot and were then centrifuged. The sera were frozen at -80°C immediately after sample collection.

APPLICATION GALLERY

Search and product-discovery interfaces

Screenshots below illustrate the functionality and interface concepts represented in the BioXplor Ltd portfolio.




Figure 3. Publication image search, including image-text and category-based exploration.




Figure 4. Product tool interface, which identifies and displays products mentioned in scientific literature.

APPLICATION GALLERY

OncoExplorer and relationship visualisation

The oncology-focused interfaces demonstrate filtered literature discovery and network-style visualisation of biomedical relationships.




Figure 5. Network visualisation of relationships identified through literature analysis.




Figure 6. OncoExplorer search results with date distribution, filters and biomedical term labels.

APPLICATION GALLERY

OncoExplorer result detail

A further view of the OncoExplorer interface showing disease filters, publication records and extracted term labels.




Figure 7. OncoExplorer result-detail view with disease filters and extracted biomedical concepts.

Acquisition proposition

BioXplor Ltd is presenting its intellectual property portfolio for acquisition in connection with the company's planned orderly closure. The portfolio comprises the software tools, platform variants and databases described in this document. The indicative asking price for the portfolio is approximately EUR 10,000, subject to buyer review and definitive transaction documents.

BIOXPLOL LTD INTELLECTUAL PROPERTY

Confidential high-level acquisition overview

PAVELS KARTASEVS

CTO, BioXplor Ltd