


Extreme Kinetic Pressure Near Dynamical Black Holes in the PPC Law of Gravity

Author: Pawan Upadhyay

Affiliation: Independent Researcher

Email: pawanupadhyay28@hotmail.com

ORCID ID: <https://orcid.org/0009-0007-9077-5924>



Abstract

This paper examines the role of kinetic pressure near the event horizons of dynamical black holes within the framework of Pawan Upadhyay's Pressure Curvature Law of Gravity (PPC Law of Gravity). The theory proposes that total gravitational pressure consists of two components: rest pressure and kinetic pressure. The kinetic-pressure component is related to kinetic energy density through a variable equation-of-state parameter. Since most astrophysical black holes are dynamical systems that continuously interact with surrounding matter and radiation, large kinetic energy densities may arise near their event horizons. The PPC framework proposes that the equation-of-state parameter approaches unity in this region, causing kinetic pressure to become approximately equal to kinetic energy density. Consequently, extreme kinetic energy

densities produce extreme kinetic pressures, making kinetic pressure a dominant contributor to gravitational pressure near the event horizon.

1. Introduction

Black holes are among the most extreme gravitational objects in the universe. While idealized theoretical models often describe stationary black holes, most astrophysical black holes are dynamical systems. They accrete matter, interact with surrounding plasma, emit energetic outflows, and evolve through continuous exchanges of energy and momentum with their environments.

The PPC Law of Gravity emphasizes the role of pressure in the generation of spacetime curvature. Within this framework, gravitational pressure is separated into distinct components associated with different forms of energy. The purpose of this paper is to investigate the behavior of these pressure components near the event horizons of dynamical black holes.

2. Gravitational Pressure in the PPC Framework

The PPC Law proposes that total gravitational pressure is given by

$$P_g = P_{\text{rest}} + P_{\text{kinetic}},$$

where P_g denotes the total gravitational pressure, P_{rest} denotes the pressure associated with rest-energy density, and P_{kinetic} denotes the pressure associated with kinetic-energy density.

The kinetic-pressure component is defined as

$$P_{\text{kinetic}} = \omega K_d,$$

where K_d is the kinetic energy density and ω is a dimensionless variable equation-of-state parameter.

Unlike conventional models that employ constant equation-of-state parameters, the PPC framework allows ω to vary with position and local physical conditions.

3. Dynamical Black Holes and Kinetic Energy Density

Most black holes observed in nature are dynamical rather than perfectly static. Matter falling toward a black hole accelerates under the influence of gravity and acquires increasingly large kinetic energy.

Accretion disks, infalling plasma, and energetic particle flows contribute to the growth of kinetic energy density near the event horizon. Consequently, the kinetic energy density may become extremely large in the immediate vicinity of a dynamical black hole.

Within the PPC framework, this increase in kinetic energy density directly influences the kinetic-pressure component of the gravitational field.

4. Near-Horizon Pressure Regime

The PPC Law proposes that near the event horizon the equation-of-state parameter approaches unity,

$$\omega \approx 1.$$

Under this condition,

$$P_{\text{kinetic}} \approx K_d.$$

The theory further proposes that the rest-pressure component approaches the rest-energy density,

$$P_{\text{rest}} \approx E_d,$$

where E_d denotes the rest-energy density.

Substituting these relations into the gravitational-pressure equation yields

$$P_g \approx E_d + K_d.$$

This result indicates that both rest-energy density and kinetic-energy density contribute significantly to the total gravitational pressure near the event horizon.

5. Extreme Kinetic Pressure Near the Event Horizon

Because kinetic energy density becomes extremely large near the event horizon of a dynamical black hole, the corresponding kinetic pressure also becomes extremely large.

When

$K_d \rightarrow \text{extreme values,}$

the relation

$$P_{\text{kinetic}} \approx K_d$$

implies that

$P_{\text{kinetic}} \rightarrow \text{extreme values.}$

Therefore, the event horizon may be characterized as an extreme kinetic-pressure region.

In the PPC framework, kinetic pressure may become the dominant component of gravitational pressure in strong gravitational fields. As a result, the curvature of spacetime near the event horizon may be strongly influenced by the kinetic state of matter.

6. Physical Interpretation

The PPC Law of Gravity suggests that gravitational pressure depends not only upon rest energy but also upon the dynamical motion of matter. Matter possessing large kinetic energy contributes additional gravitational pressure through the kinetic-pressure component.

Near the event horizon, where kinetic energy densities become exceptionally large, kinetic pressure approaches the same order of magnitude as kinetic energy density itself. Consequently, the strong gravitational environment surrounding black holes may represent a regime in which kinetic pressure plays a fundamental role in the generation of spacetime curvature.

7. Conclusion

The PPC Law of Gravity decomposes gravitational pressure into rest-pressure and kinetic-pressure components. The kinetic-pressure component is related to kinetic-energy density through a variable equation-of-state parameter.

Most astrophysical black holes are dynamical systems that generate large kinetic-energy densities near their event horizons. The PPC framework proposes that the equation-of-state parameter approaches unity in this regime, leading to the relation .

$$P_{\text{kinetic}} \approx K_d.$$

As kinetic energy density becomes extreme, kinetic pressure likewise becomes extreme. Consequently, the event horizon of a dynamical black hole may be regarded as an extreme kinetic-pressure environment. This framework suggests that kinetic pressure becomes a significant, and potentially dominant, contributor to gravitational pressure and spacetime curvature in the strongest gravitational fields known in nature.

Keywords: PPC Law of Gravity, Dynamical Black Holes, Event Horizon, Kinetic Pressure, Gravitational Pressure, Kinetic Energy Density, Equation of State Parameter, Spacetime Curvature.

Rest Pressure and Kinetic Pressure Near the Event Horizon

In the PPC Law of Gravity, the total gravitational pressure is proposed to consist of two distinct components: rest pressure and kinetic pressure. The total gravitational pressure is expressed as

$$P_g = P_{\text{rest}} + P_{\text{kinetic}}.$$

The kinetic-pressure component is related to the kinetic energy density by

$$P_{\text{kinetic}} = \omega K_d,$$

where K_d denotes the kinetic energy density and ω is a dimensionless variable equation-of-state parameter. The parameter ω is not constant but varies with position and local physical conditions.

Near the event horizon of a black hole, the kinetic energy density becomes extremely large due to the intense gravitational environment. Within the PPC framework, it is proposed that the equation-of-state parameter approaches unity in this regime:

$$\omega \approx 1.$$

Consequently, the kinetic pressure becomes approximately equal to the kinetic energy density:

$$P_{\text{kinetic}} \approx K_d.$$

The theory further proposes that near the event horizon the rest-pressure component approaches the rest-energy density:

$$P_{\text{rest}} \approx E_d,$$

where E_d is the rest-energy density.

Substituting these limiting relations into the gravitational pressure equation gives

$$P_g \approx E_d + K_d.$$

This result suggests that, in the strong-gravity region near the event horizon, both rest-energy density and kinetic-energy density contribute significantly to the total gravitational pressure. As the kinetic energy density increases to extreme values, the corresponding kinetic pressure also becomes extreme. Therefore, kinetic pressure may become a dominant component of the total gravitational pressure in the vicinity of the event horizon.

Within the PPC Law of Gravity, this behavior implies that gravitational pressure is determined not only by rest energy but also by the kinetic state of matter. Consequently, the extreme gravitational environment near the event horizon is characterized by substantial pressure contributions arising from both rest-energy density and kinetic-energy density. Such a regime may play an important role in the curvature of spacetime and the gravitational dynamics of matter near black holes.

Extreme Kinetic Pressure Near the Event Horizon

Within the framework of the PPC Law of Gravity, the kinetic-pressure component is defined by

$$P_{\text{kinetic}} = \omega K_d,$$

where K_d is the kinetic energy density and ω is a dimensionless variable equation-of-state parameter.

Near the event horizon of a black hole, matter experiences an extremely strong gravitational environment. As matter accelerates toward the event horizon, its kinetic energy density increases dramatically. The PPC framework further proposes that, in this regime,

$$\omega \approx 1.$$

Consequently,

$$P_{\text{kinetic}} \approx K_d.$$

Since the kinetic energy density becomes extremely large near the event horizon, the corresponding kinetic pressure also becomes extremely large. Thus, the vicinity of the event horizon is characterized by an extreme kinetic-pressure regime.

In the PPC Law of Gravity, this extreme kinetic pressure contributes significantly to the total gravitational pressure,

$$P_g = P_{\text{rest}} + P_{\text{kinetic}},$$

and may become the dominant pressure component in the strong-field region surrounding the event horizon. Therefore, the event horizon can be viewed as a region where kinetic pressure reaches its highest physically attainable values, reflecting the extreme dynamical state of matter and energy in the gravitational field of a black hole.

Most astrophysical black holes are dynamical systems. Matter and radiation continuously interact with the black hole environment, leading to ongoing changes in energy density, kinetic energy density, pressure, and spacetime curvature. Consequently, the gravitational pressure near the event horizon is generally not static but evolves with the dynamical state of the surrounding matter.

Because most black holes are dynamical, the kinetic energy density K_d near their event horizons can be substantial. Since the PPC Law relates kinetic pressure to kinetic energy density through

$P_{\text{kinetic}} = \omega K_d$, dynamical black holes may naturally generate extreme kinetic pressures in the near-horizon region. If ω approaches unity near the event horizon, then $P_{\text{kinetic}} \approx K_d$, implying that extreme kinetic energy densities produce correspondingly extreme kinetic pressures.

As kinetic energy density becomes extreme, kinetic pressure likewise becomes extreme. Consequently, the event horizon of a dynamical black hole may be regarded as an extreme kinetic-pressure environment. Within the PPC Law of Gravity, this suggests that the **gravitational pressure field** near the event horizon is dominated by kinetic pressure, making kinetic energy density a major contributor to spacetime curvature in the strongest gravitational environments known in nature.

Definition: In the PPC Law of Gravity, the term gravitational pressure field refers to the pressure distribution generated by rest-energy density and kinetic-energy density. This pressure field acts as the source of spacetime curvature and gravitational phenomena.

Pressure Gradients Near the Event Horizon

In the PPC Law of Gravity, the gravitational pressure field is determined by the distribution of rest pressure and kinetic pressure throughout spacetime. Since kinetic energy density becomes extremely large near the event horizon, the associated kinetic pressure also becomes extremely large. Consequently, the spatial variation of pressure becomes increasingly steep in the near-horizon region.

Mathematically, if

$$P_g = P_{\text{rest}} + P_{\text{kinetic}},$$

and

$$P_{\text{kinetic}} = \omega K_d,$$

with

$$\omega \approx 1$$

near the event horizon, then

$$P_g \approx E_d + K_d.$$

As E_d and K_d change rapidly with position in the strong-gravity region, the pressure gradients

$$\nabla P_g$$

may become extremely large.

Near the event horizon of a dynamical black hole, both rest-energy density and kinetic-energy density may attain extreme values. Within the PPC framework, these energy densities generate correspondingly large rest pressure and kinetic pressure. As a consequence, the gravitational pressure field exhibits very steep spatial variations. The resulting pressure gradients become extremely large in the vicinity of the event horizon. Therefore, the near-horizon region may be characterized not only by extreme gravitational pressure but also by extreme pressure gradients. These pressure gradients may play a significant role in the generation of spacetime curvature and the dynamics of matter surrounding black holes.

Strong spacetime curvature corresponds to strong gravitational pressure gradients.

Maximum Energy and Gravitational Pressure Near the Event Horizon

Within the framework of the PPC Law of Gravity, a fundamental **Maximum Energy Limit** is proposed. The total local energy density, consisting of rest-energy density and kinetic-energy density, satisfies

$$E_d + K_d \leq E_{\text{max}},$$

where E_{max} denotes the maximum physically attainable energy density.

Near the event horizon of a dynamical black hole, both the rest-energy density E_d and the kinetic-energy density K_d increase substantially. As a consequence, the combined energy density approaches the Maximum Energy Limit:

$$E_d + K_d \rightarrow E_{\text{max}}.$$

The PPC Law further proposes that the total gravitational pressure is given by

$$P_g = P_{\text{rest}} + P_{\text{kinetic}},$$

where P_{rest} is the rest-pressure component and P_{kinetic} is the kinetic-pressure component.

Near the event horizon, the theory assumes

$$P_{\text{rest}} \approx E_d,$$

and

$$P_{\text{kinetic}} \approx K_d.$$

Therefore,

$$P_g \approx E_d + K_d.$$

As the total energy density approaches the Maximum Energy Limit, the total gravitational pressure correspondingly approaches its maximum value:

$$P_g \rightarrow P_{g,\text{max}}.$$

Within the PPC framework, the event horizon of a dynamical black hole represents an extreme gravitational-pressure environment where energy density, gravitational pressure, and pressure gradients simultaneously approach their maximum physical limits. The theory therefore predicts that the strongest spacetime curvature occurs in regions where the gravitational pressure field and its gradients attain their greatest magnitudes.

$$E_{\max} \Rightarrow P_{g,\max} \Rightarrow (\nabla P_g)_{\max}.$$

Maximum Energy and Gravitational Pressure Near Neutron Stars

Within the framework of the PPC Law of Gravity, neutron stars represent some of the most extreme gravitational-pressure environments in the universe outside of black holes. The enormous matter density inside and near a neutron star produces very large rest-energy densities and significant kinetic-energy densities.

The total local energy density is given by

$$E_{\text{total}} = E_d + K_d,$$

where E_d is the rest-energy density and K_d is the kinetic-energy density.

The PPC Law proposes that the total gravitational pressure is

$$P_g = P_{\text{rest}} + P_{\text{kinetic}},$$

with

$$P_{\text{kinetic}} = \omega K_d,$$

where ω is a dimensionless variable equation-of-state parameter.

The theory further proposes that near neutron stars,

$$\omega \approx 1.$$

Consequently,

$$P_{\text{kinetic}} \approx K_d.$$

In addition, the rest-pressure component is assumed to satisfy

$$P_{\text{rest}} \approx E_d.$$

Therefore, near neutron stars,

$$P_g \approx E_d + K_d.$$

Because both E_d and K_d attain very large values in the intense gravitational-pressure environment of a neutron star, the total gravitational pressure also becomes very large:

$$P_g \gg 0.$$

$$P_{\text{rest}} \gg 0,$$

$$P_{\text{kinetic}} \gg 0.$$

As a result,

$$P_g \gg 0.$$

Furthermore, the gravitational pressure field varies rapidly with position near the neutron star surface. As a consequence, the pressure gradients become extremely large:

$$|\nabla P_g| \gg 0.$$

Within the PPC framework, neutron stars may therefore be regarded as high-pressure gravitational systems in which rest pressure and kinetic pressure contribute almost equally to the total gravitational pressure field. The near-unity value of the equation-of-state parameter implies that kinetic energy density contributes nearly directly to gravitational pressure.

Although the gravitational pressure field surrounding neutron stars is extraordinarily strong, it remains below the maximum-energy and maximum-pressure regime proposed for black-hole event horizons. Thus, neutron stars occupy an intermediate position between ordinary stellar objects and black holes in the hierarchy of gravitational-pressure environments.

Ordinary Stars < White Dwarfs < Neutron Stars < Black Hole Event Horizons.

In this sequence, the quantities E_d , K_d , P_g , and $|\nabla P_g|$ increase progressively, reaching their greatest values in the vicinity of black-hole event horizons, where the PPC framework predicts the approach to maximum energy density and maximum gravitational pressure.

Thus, the PPC hierarchy of compact objects can be expressed qualitatively as

Ordinary Stars < White Dwarfs < Neutron Stars < Black Hole Event Horizons,

The extreme density and internal kinetic motion of matter within neutron stars generate large kinetic energy densities. Within the PPC framework, these large kinetic energy densities produce substantial kinetic pressures, making kinetic pressure an important contributor to the gravitational pressure field surrounding neutron stars.

Neutron stars are characterized by extremely strong internal kinetic motion. The dense neutron matter, rapid particle motion, and intense gravitational-pressure environment contribute to a large kinetic energy density K_d . Within the PPC framework, where the equation-of-state parameter approaches unity ($\omega \approx 1$) near neutron stars, the kinetic pressure satisfies $P_{\text{kinetic}} \approx K_d$. Consequently, the strong kinetic motion of matter inside and around neutron stars generates a substantial kinetic-pressure component, which contributes significantly to the total gravitational pressure field.

The kinetic motion of matter within neutron stars is extremely strong due to their enormous density and intense gravitational-pressure field. As a result, neutron stars possess large kinetic energy densities and correspondingly large kinetic pressures. In the PPC framework, with $\omega \approx 1$, the kinetic pressure becomes approximately equal to the kinetic energy density, making kinetic pressure a major contributor to the gravitational pressure field near neutron stars.