

Mass Density–Energy Density Equivalence: A Density Formulation of Mass–Energy Equivalence

Author: Pawan Upadhyay


Affiliation: Independent Researcher

Email: pawanupadhyay28@hotmail.com

ORCID iD: <https://orcid.org/0009-0007-9077-5924>

Proposed Theory: Pawan Upadhyay's Pressure Curvature Law of Gravity (PPC Law of Gravity)

Date: 06 Nov. 2025



Abstract

Einstein's mass–energy equivalence establishes that mass and energy are related through the equation $E=mc^2$. This paper develops the corresponding density formulation of this

relationship. By considering energy and mass within a common volume and dividing the mass–energy relation by volume, the following relation is obtained:

$$E_d = \rho c^2$$

where $E_d = E/V$ represents energy density and $\rho = m/V$ represents mass density. The equation demonstrates that, for rest-mass energy distributed over a volume, energy density and mass density are directly proportional, with the proportionality factor given by c^2 .

The paper proposes the term **Mass Density–Energy Density Equivalence** for this density-level representation. It examines the dimensional consistency, mathematical derivation, physical interpretation, limiting cases, and possible applications of the relation. The paper also distinguishes the density formulation from Einstein's original mass–energy equivalence and identifies directions in which the density relation could be developed into a broader theoretical framework.

Keywords: mass-energy equivalence, mass density, energy density, Einstein, $E=mc^2$, $E_d = \rho c^2$, relativistic physics

1. Introduction

The equivalence between mass and energy is one of the foundational results of modern physics. Einstein's famous relation

$$[E=mc^2]$$

establishes that mass possesses an equivalent amount of energy and that the conversion factor between mass and energy is the square of the speed of light.

Mass and energy, however, can also be described as quantities distributed through space. In such a description, the relevant quantities are not total mass and total energy but **mass density** and **energy density**.

This paper examines the density representation of mass–energy equivalence. Starting from Einstein's relation and considering a volume V , the relation can be expressed in terms of mass per unit volume and energy per unit volume.

The resulting equation is

$$E_d = \rho c^2$$

where

$$[E_d = E/V]$$

and

$$\rho = m/V$$

The central conceptual proposition of this paper is that **mass density and the corresponding rest-mass energy density are two descriptions of the same underlying mass–energy equivalence at the density level.**

2. Einstein's Mass–Energy Equivalence

Einstein's relation is

$$E = mc^2$$

Here, E represents energy, m represents mass, and c represents the speed of light in vacuum.

The equation does not imply that mass and energy are numerically identical. Rather, it establishes a quantitative equivalence between them through the constant c^2 .

Because c is extremely large, even a small amount of mass corresponds to a very large rest-energy.

3. From Mass–Energy Equivalence to Density Equivalence

Consider a quantity of mass m and its corresponding energy E contained or distributed within a volume V .

Starting with

$$E = mc^2$$

divide both sides by V :

$$E/V = (m/V) c^2$$

Define energy density as

$$E_d = E/V$$

and mass density as

$$\rho = m/V$$

Therefore,

$$E_d = \rho c^2$$

This is the principal equation examined in this paper.

4. Definition of the Proposed Relation

The equation

$$E_d = \rho c^2$$

may be called the **Mass Density–Energy Density Equivalence Relation**.

The variables are defined as follows:

Symbol	Meaning	SI unit
E_d	Energy density	J/m ³
E	Energy	J
V	Volume	m ³
ρ	Mass density	kg/m ³
m	Mass	kg
c	Speed of light	m/s

The equation can also be rearranged as

$$\rho = E_d / c^2$$

Thus, a given rest-energy density corresponds to a definite mass density.

5. Dimensional Analysis

The dimensional consistency of the equation can be demonstrated directly.

Mass density has dimensions

$$[\rho] = \text{kg/m}^3$$

The speed of light squared has dimensions

$$[c^2] = \text{m}^2/\text{s}^2$$

Therefore,

$$[\rho c^2] = \frac{\text{kg}}{\text{m}^3} \frac{\text{m}^2}{\text{s}^2} = \frac{\text{kg}}{\text{m s}^2} = \frac{\text{J}}{\text{m}^3}$$

Since energy density has units

$$[E_d] = \text{J/m}^3$$

the equation is dimensionally consistent.

6. Physical Interpretation

The relation

$$E_d = \rho c^2$$

provides a density-based interpretation of mass–energy equivalence.

If a system has mass density ρ , then its corresponding **rest-mass energy density** is ρc^2 .

For example, if

$$\rho = 1 \text{ (kg/m}^3\text{)}$$

then the corresponding rest-mass energy density is

$$[E_d = c^2 \text{ (J/m}^3\text{)}]$$

$$E_d = c^2 \text{ J/m}^3$$

The equation therefore provides a direct conversion between mass density and rest-mass energy density.

The key point is that this relation concerns **rest-mass energy**. It should not automatically be interpreted as saying that every form of energy density in every physical system is simply equal to ρc^2 . Systems can possess kinetic, thermal, electromagnetic, binding, pressure-related, and other contributions to their total energy.

7. Relation to Einstein's Original Equation

The proposed density relation is not independent of Einstein's equation in the mathematical sense.

Starting from

$$[E = mc^2]$$

and dividing by V gives

$$[E/V = (m/V) c^2,]$$

which becomes

$$[E_d = \rho c^2]$$

Consequently, the density equation should be understood as a **density formulation or local representation of the rest-mass component of mass–energy equivalence**.

The scientific value of developing this formulation therefore lies not in claiming that the algebraic transformation itself replaces Einstein's equation, but in investigating whether the density formulation provides useful new conceptual, mathematical, or physical applications.

8. Local Formulation

If the mass distribution is not uniform, the global quantities m , E , and V may be replaced by local densities.

For a continuous distribution, one can write

$$[\rho(\mathbf{x}) = \frac{dm}{dV}]$$

$$\rho(\mathbf{x}) = \frac{dm}{dV}$$

and define the corresponding rest-mass energy density as

$$[E_d(\mathbf{x}) = \frac{dE}{dV}.]$$

The density relation then becomes

$$[\boxed{E_d(\mathbf{x}) = \rho(\mathbf{x})c^2}.]$$

$$E_d(\mathbf{x}) = \frac{dE}{dV}.$$

The density relation then becomes

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2.$$

This formulation is potentially more useful for systems in which mass density varies from one spatial location to another.

9. Integrated Form

The density formulation must reproduce the original mass–energy relation when integrated over a volume.

Starting with

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2,$$

integrating over a volume gives

$$E = \int E_d(\mathbf{x}) dV.$$

Therefore,

$$E = c^2 \int \rho(\mathbf{x}) dV.$$

Since

$$m = \int \rho(\mathbf{x}) dV,$$

we obtain

$$E = mc^2 .$$

Thus, the density formulation is mathematically consistent with the integrated mass–energy relation.

10. Conceptual Proposition

This paper proposes the following conceptual statement:

Mass density and rest-mass energy density represent two quantitatively equivalent descriptions of the same mass–energy content, related by the universal factor c^2 .

In this sense, mass density can be interpreted as an energy-density representation after multiplication by c^2 , while energy density associated with rest mass can be converted into an equivalent mass density by division by c^2 .

This perspective emphasizes that mass–energy equivalence can be expressed not only globally through total mass and total energy, but also at the level of spatial density.

11. Limitations and Scope

The equation

$$E_d = \rho c^2$$

must be used with a clear definition of E_d .

If E_d denotes **rest-mass energy density**, the relation follows directly from $E=mc^2$.

Definition and Scope of the Energy-Density Relation

The relation

$$E_d = \rho c^2$$

requires a precise definition of the energy density (E_d).

When (E_d) denotes the **rest-mass energy density**, the relation follows directly from Einstein's mass–energy equivalence,

$$[E_0 = m_0 c^2]$$

Definition and Scope of the Energy-Density Relation

The relation

$$E_d = \rho c^2$$

requires a precise definition of the energy density

$$E_d$$

.

When

$$E_d$$

denotes the **rest-mass energy density**, the relation follows directly from Einstein's mass–energy equivalence,

$$E_0 = m_0 c^2.$$

Defining

$$\rho_0 = \frac{m_0}{V}$$

and

$$E_{0d} = \frac{E_0}{V} ,$$

we obtain

$$E_{0d} = \rho_0 c^2 .$$

Thus, the precise density formulation of the mass–energy equivalence is

rest-mass energy density = (rest-mass density) (c^2)

However, if

$$E_d$$

is defined as the **total energy density of a relativistic system**, it may contain contributions beyond rest-mass energy. In that case,

$$E_d \neq \rho_0 c^2$$

in general, and the simple relation

$$E_d = \rho c^2$$

should not automatically be interpreted as the complete energy budget.

Therefore, in the present formulation, the relation

$$E_{0d} = \rho_0 c^2$$

is specifically identified as the relation between **rest-mass density** and **rest-mass energy density**.

The broader identification

$$E_d = \rho c^2$$

should only be adopted if a further theoretical framework establishes what

E_d

and

ρ

represent and demonstrates that the relation remains valid for the complete relativistic system. Such an extension would require appropriate theoretical justification and, ultimately, experimental or observational validation.

For the perfect-fluid formulation,

E_d

should therefore be defined independently as the **local energy density appearing in the stress–energy tensor**:

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu} .$$

The gravitational pressure field is then given by

$$P_g = \omega E_d .$$

This distinction keeps the rest-mass energy relation separate from the more general energy-density description.

12. Potential Applications

The density formulation may provide a useful framework for:

1. **Relativistic astrophysics** — expressing mass distributions in energy-density units.
2. **High-energy physics** — comparing matter density with corresponding rest-energy density.
3. **Cosmology** — describing matter distributions using energy-density units.
4. **Theoretical physics** — developing density-based formulations of mass–energy equivalence.
5. **Educational physics** — demonstrating how a global physical relation can be transformed into a density relation.
6. **Computational physics** — representing mass distributions and their corresponding rest-energy distributions within spatial models.

These applications should be treated as areas for further investigation rather than as established consequences of the equation alone.

13. Discussion

The central result of this paper is the density representation

$$E_d = \rho c^2$$

The derivation is straightforward, but the formulation highlights an important change of perspective: the mass–energy relationship can be considered at the level of **quantities per unit volume**.

Einstein's equation connects total mass and total energy. The density formulation connects mass per unit volume with rest-energy per unit volume.

The distinction between the original relation and the present formulation is therefore primarily one of representation and interpretation.

A stronger claim of originality would require additional theoretical development beyond the algebraic derivation. Such development could investigate whether a density-based framework leads to new equations, measurable predictions, or relationships that are not already contained in established relativity.

14. Conclusion

This paper develops the density formulation of mass–energy equivalence:

$$E_d = \rho c^2$$

With

$$E_d = E/V$$

and

$$\rho = m/V$$

the equation follows directly from Einstein's

$$[E = mc^2]$$

The formulation establishes a direct quantitative correspondence between mass density and rest-mass energy density. It can also be expressed locally as

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2$$

and, upon integration over space, recovers the conventional mass–energy relation.

The principal contribution proposed here is therefore a **density-centered conceptual framework for discussing mass–energy equivalence**, rather than a claim that the algebraic relation itself is independent of Einstein's result.

Future research could investigate whether this framework can be extended to total relativistic energy density, curved spacetime, gravitational systems, or other physical contexts in ways that produce genuinely new theoretical or experimentally testable results.

References

1. Einstein, A. (1905). *Does the Inertia of a Body Depend Upon Its Energy-Content?* Annalen der Physik, 18, 639–641.
 2. Einstein, A. (1905). *On the Electrodynamics of Moving Bodies*. Annalen der Physik, 17, 891–921.
 3. Einstein, A. (1916). *Die Grundlage der allgemeinen Relativitätstheorie*. Annalen der Physik, 49, 769–822.
 4. Standard texts on special relativity and relativistic energy–momentum may be consulted for the distinction between rest-mass energy density and total energy density.
-

Author's Note

The equation

$$E_d = \rho c^2$$

is proposed here as a **Mass Density–Energy Density Equivalence Relation**, derived from the established mass–energy equivalence relation by expressing both quantities per unit volume. The purpose of this work is to develop and investigate the density interpretation and its potential theoretical applications.

$$E_{0d} = \rho_0 c^2 \; .$$

$$x^\mu = (x,y,z,ct)$$

$$E_d = E_d(x^\mu)$$

$$P_g = P_g(x^\mu)$$

$$\omega = \omega(x^\mu)$$

$$P_g(x^\mu) = \omega(x^\mu) E_d(x^\mu)$$

$$\omega(x^\mu) \approx 1$$

therefore

$$P_g(x^\mu) \approx E_d(x^\mu) \ .$$

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu}$$

and, when

$$\omega \approx 1$$

,

$$T_{\mu\nu} \approx 2E_d u_\mu u_\nu + E_d g_{\mu\nu} \ .$$

[Book style Research Paper]

MASS DENSITY–ENERGY DENSITY EQUIVALENCE

A Density Formulation of Mass–Energy Equivalence

Author: Pawan Upadhyay


Affiliation: Independent Researcher

Email: pawanupadhyay28@hotmail.com

ORCID iD: <https://orcid.org/0009-0007-9077-5924>

Proposed Theory: Pawan Upadhyay's Pressure Curvature Law of Gravity (PPC Law of Gravity)

Date: 06 Nov. 2025



TITLE PAGE

MASS DENSITY–ENERGY DENSITY EQUIVALENCE

A Density Formulation of Mass–Energy Equivalence

Author

Pawan Upadhyay

Research Paper / Scholarly Monograph

2025-2026

COPYRIGHT AND AUTHOR'S NOTE

© 2025-2026 Pawan Upadhyay

All rights reserved.

This work presents a density-based formulation of the established mass–energy equivalence relation. The equation

$$E_d = \rho c^2$$

is obtained by expressing Einstein's mass–energy relation in terms of quantities per unit volume.

The author distinguishes the mathematical derivation of this equation from any future theoretical claims that may be developed from it.

CONTENTS

Preface

Abstract

Chapter 1 — Introduction

1.1 Mass and Energy

1.2 From Total Quantities to Densities

1.3 Purpose of the Study

Chapter 2 — Einstein's Mass–Energy Equivalence

- 2.1 The Equation $E=mc^2$
- 2.2 Physical Meaning
- 2.3 Mass and Energy as Equivalent Quantities

Chapter 3 — Mass Density and Energy Density

- 3.1 Mass Density
- 3.2 Energy Density
- 3.3 Volume as the Connecting Quantity

Chapter 4 — The Mass Density–Energy Density Equation

- 4.1 Mathematical Derivation
- 4.2 Definition of Variables
- 4.3 Dimensional Analysis

Chapter 5 — Local Density Formulation

- 5.1 Spatially Varying Mass Density
- 5.2 Local Energy Density
- 5.3 Integrated Form

Chapter 6 — Physical Interpretation

- 6.1 Rest-Mass Energy Density
- 6.2 Density as an Alternative Representation
- 6.3 Relationship to Einstein's Equation

Chapter 7 — Scope and Limitations

- 7.1 Rest Energy and Total Energy
- 7.2 Other Forms of Energy
- 7.3 Limits of the Present Formulation

Chapter 8 — Potential Applications

- 8.1 Relativistic Physics
- 8.2 Astrophysics
- 8.3 Cosmology
- 8.4 Computational Physics

Chapter 9 — Future Research

Chapter 10 — Conclusion

References

PREFACE

The relationship between mass and energy is one of the central ideas of modern physics.

Einstein's equation

$$E=mc^2$$

expresses the equivalence between mass and its corresponding rest energy.

The present work approaches this relationship from another level of description: **density**.

If mass is distributed through a volume, it can be expressed as mass density. Similarly, energy distributed through a volume can be expressed as energy density. This naturally leads to the question:

If mass and energy are equivalent, can their corresponding densities also be expressed as equivalent quantities?

Starting from Einstein's relation and dividing by volume leads to

$$E_d = \rho c^2$$

This work develops that density formulation and examines its meaning, mathematical structure, limitations, and possible applications.

ABSTRACT

Einstein's mass–energy equivalence establishes the relation ($E=mc^2$). This work develops its density representation by considering mass and energy with respect to a common volume. Dividing the mass–energy relation by volume gives

$$E_d = \rho c^2$$

where ($E_d=E/V$) is rest-mass energy density and ($\rho=m/V$) is mass density.

The equation demonstrates that rest-mass energy density and mass density are directly proportional through the universal factor (c^2). A local formulation is also considered:

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2.$$

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2$$

Integration of this local relation recovers Einstein's original mass–energy equation.

The paper proposes the term **Mass Density–Energy Density Equivalence Relation** for this density-level representation. It emphasizes that the equation follows mathematically from Einstein's established relation and therefore should not be presented as an independent replacement for ($E=mc^2$). The potential originality lies in the conceptual development and any future theoretical consequences, applications, or experimentally testable predictions arising from the density framework.

Keywords: Mass, energy, mass density, energy density, relativity, Einstein, ($E=mc^2$), $E_d = \rho c^2$

CHAPTER 1

INTRODUCTION

1.1 Mass and Energy

Mass and energy are two fundamental physical quantities.

Classical physics traditionally treats mass and energy as distinct quantities. Modern physics, however, established a quantitative relationship between them.

Einstein's famous relation

$$[E=mc^2]$$

shows that mass possesses an equivalent rest energy.

This relationship fundamentally changed the understanding of matter and energy.

1.2 From Total Quantities to Densities

Physical quantities are frequently expressed not only as totals but also as quantities per unit volume.

For mass,

$$\rho = m/V$$

For energy,

$$[E_d = E/V]$$

Thus, if total mass and total energy have an equivalence, it is natural to investigate the corresponding relationship between their densities.

This gives the central equation of this work:

$$E_d = \rho c^2$$

1.3 Purpose of the Study

The purpose of this work is to:

1. derive the density relation mathematically;
 2. define mass density and rest-mass energy density;
 3. examine its dimensional consistency;
 4. formulate its local representation;
 5. examine its physical interpretation;
 6. distinguish it from the original ($E=mc^2$) relation; and
 7. identify possible directions for further research.
-

CHAPTER 2

EINSTEIN'S MASS–ENERGY EQUIVALENCE

2.1 The Equation ($E=mc^2$)

Einstein's mass–energy relation is

$$E=mc^2$$

Here:

E =energy, m =mass

and

c = speed of light in vacuum

The equation establishes that mass has an equivalent rest-energy content.

2.2 Physical Meaning

The factor

$$c^2$$

is extremely large. Consequently, even a small quantity of mass corresponds to a very large amount of rest energy.

The equation therefore provides a bridge between two descriptions of physical content:

$$\text{mass} \longleftrightarrow \text{energy}$$

CHAPTER 3

MASS DENSITY AND ENERGY DENSITY

3.1 Mass Density

Mass density is defined as mass per unit volume:

$$\rho = m/V$$

Its SI unit is

$$\text{kg/m}^3$$

3.2 Energy Density

Energy density is defined as energy per unit volume:

$$E_d = E/V$$

Its SI unit is

$$\text{J/m}^3$$

3.3 Volume as the Connecting Quantity

Both quantities are defined using the same spatial measure:

$$[V]$$

Therefore,

$$m/V$$

describes mass density, while

$$E/V$$

describes energy density.

This makes volume the natural mathematical bridge between total mass–energy equivalence and its density representation.

CHAPTER 4

THE MASS DENSITY–ENERGY DENSITY EQUATION

4.1 Mathematical Derivation

Begin with Einstein's equation:

$$[E = mc^2]$$

Divide both sides by V:

$$E/V = (mc^2)/V$$

Rearranging,

$$E/V = (m/V) c^2$$

Now define

$$E_d = E/V$$

and

$$\rho = m/V$$

Therefore,

$$E_d = \rho c^2$$

This is the central equation of the present work.

4.2 Definition of Variables

Symbol	Definition	SI Unit
E_d	Rest-mass energy density	J/m ³

E	Rest energy	J
ρ	Mass density	kg/m ³
m	Mass	kg
V	Volume	m ³
c	Speed of light	m/s

The equation can also be written as

$$\rho = E_d / c^2$$

4.3 Dimensional Analysis

For mass density,

$$[\rho] = \text{Kg/m}^3$$

For c^2 ,

$$[c^2] = \text{m}^2/\text{s}^2.$$

Therefore,

$$\rho [c^2] = \frac{\text{kg}}{\text{m}^3} \frac{\text{m}^2}{\text{s}^2} = \frac{\text{kg}}{\text{m} \cdot \text{s}^2}.$$

$$[\rho c^2] = \frac{\text{kg}}{\text{m}^3} \frac{\text{m}^2}{\text{s}^2} = \frac{\text{kg}}{\text{m s}^2} = \frac{\text{J}}{\text{m}^3}$$

Since

$$1(\text{J}) = 1 \text{ kg (m}^2/\text{s}^2)$$

we have

$$(\text{J})/(\text{m}^3) = (\text{kg})/(\text{m s}^2)$$

Hence,

$$\rho c^2 = E_d$$

The equation is dimensionally consistent.

CHAPTER 5

LOCAL DENSITY FORMULATION

5.1 Spatially Varying Mass Density

In a non-uniform system, mass density may vary from one location to another.

It can therefore be written as

$$\rho = \rho(\mathbf{x}).$$

More formally,

$$\rho(\mathbf{x}) = \frac{dm}{dV}.$$

5.1 Spatially Varying Mass Density

In a non-uniform system, mass density may vary from one location to another.

It can therefore be written as

$$\rho = \rho(\mathbf{x}).$$

More formally,

$$\rho(\mathbf{x}) = \frac{dm}{dV}.$$

5.2 Local Energy Density

Correspondingly, the rest-mass energy density is

$$\rho_E(\mathbf{x}) = \frac{dE}{dV}.$$

The density relation becomes

$$\boxed{E_d(\mathbf{x}) = \rho(\mathbf{x})c^2}.$$

This expresses the mass–energy equivalence at the density level for a spatially varying mass distribution.

5.2 Local Energy Density

Correspondingly, the rest-mass energy density is

$$E_d(\mathbf{x}) = \frac{dE}{dV}.$$

The density relation becomes

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2.$$

This expresses the mass–energy equivalence at the density level for a spatially varying mass distribution.

5.3 Integrated Form

Integrating over a volume gives

$$E = \int E_d(\mathbf{x}) dV.$$

Substituting the density relation,

$$E = c^2 \int \rho(\mathbf{x}) dV.$$

But

$$m = \int \rho(\mathbf{x}) dV.$$

Therefore,

$$\boxed{E = mc^2}.$$

Thus, the local density formulation reproduces Einstein's global relation.

5.3 Integrated Form

Integrating over a volume gives

$$E = \int E_d(\mathbf{x}) dV.$$

Substituting the density relation,

$$E = c^2 \int \rho(\mathbf{x}) dV.$$

But

$$m = \int \rho(\mathbf{x}) dV.$$

Therefore,

$$E = mc^2 .$$

Thus, the local density formulation reproduces Einstein's global relation.

CHAPTER 6

PHYSICAL INTERPRETATION

6.1 Rest-Mass Energy Density

The quantity

$$\rho c^2$$

represents the energy density associated with the rest mass represented by ρ .

Therefore,


$$\rho \longrightarrow \rho c^2$$

is a conversion from mass-density units to equivalent rest-energy-density units.

Similarly,

$$E_d \longrightarrow \frac{E_d}{c^2}$$

converts rest-energy density into equivalent mass density.



6.2 Density as an Alternative Representation

The equation does not introduce a new physical conversion constant.

Rather, it expresses the same mass–energy equivalence in another form:

$$E=mc^2$$

becomes

$$E_d = \rho c^2$$

The first describes total quantities.

The second describes quantities per unit volume.

6.3 Relationship to Einstein's Equation

The relation

$$E_d = \rho c^2$$

is derived from Einstein's established equation.

The proposed relation is a density formulation of mass–energy equivalence, not an independent replacement for Einstein's equation.

CHAPTER 7

SCOPE AND LIMITATIONS

7.1 Rest Energy and Total Energy

A critical distinction must be made between **rest-mass energy density** and **total energy density**.

The equation

$$E_d = \rho c^2$$

directly describes the rest-energy associated with mass density.

It does not, by itself, describe every possible contribution to the total energy of a physical system.

7.2 Other Forms of Energy

A physical system may contain:

- kinetic energy,
- thermal energy,
- electromagnetic energy,
- chemical energy,
- binding energy,
- pressure-related energy,
- gravitational contributions.

Consequently, a complete theory of total energy density may require additional variables and equations.

7.3 Limits of the Present Formulation

The present work establishes a density representation of rest-mass energy.

It does not yet establish a new theory of gravitation, cosmology, quantum mechanics, or total relativistic energy density.

Such claims would require further mathematical and experimental investigation.

CHAPTER 8

POTENTIAL APPLICATIONS

8.1 Relativistic Physics

The equation provides a convenient way to express rest-mass density in energy-density units.

8.2 Astrophysics

Astronomical objects contain spatially varying distributions of matter. The density formulation can provide an equivalent rest-energy-density representation of those distributions.

8.3 Cosmology

Cosmological matter density is frequently discussed alongside energy-density quantities. The relation provides a direct conversion between matter density and its corresponding rest-energy density.

8.4 Computational Physics

A numerical mass-density field

$\rho(\mathbf{x})$

can be transformed into an equivalent rest-energy-density field through

$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2$

This may be useful in simulations where different physical components are represented using common energy-density units.

A numerical mass-density field

$$\rho(\mathbf{x})$$

can be transformed into an equivalent rest-energy-density field through

$$E_d(\mathbf{x}) = \rho(\mathbf{x})c^2 .$$

This may be useful in simulations where different physical components are represented using common energy-density units.

CHAPTER 9

FUTURE RESEARCH

The density formulation opens several possible directions for investigation.

9.1 Total Energy Density

A future theory could investigate how rest-mass energy density combines with other energy contributions.

9.2 Time-Dependent Density

A more general formulation could consider

$$\rho = \rho(\mathbf{x}, t).$$

The corresponding rest-energy density would then be

$$E_d(\mathbf{x}, t) = \rho(\mathbf{x}, t)c^2.$$

A more general formulation could consider

$$\rho = \rho(\mathbf{x}, t).$$

The corresponding rest-energy density would then be

$$E_d(\mathbf{x}, t) = \rho(\mathbf{x}, t)c^2.$$

9.3 Relativistic Matter

The relationship between mass density, energy density, momentum density, and pressure could be examined using relativistic continuum models.

9.4 Cosmological Models

The density formulation could be investigated within cosmological models to determine whether it provides useful new interpretations.

9.5 New Testable Predictions

The most important future objective would be to determine whether an extension of the density framework produces predictions that go beyond the established consequences of $E=mc^2$.

CHAPTER 10

CONCLUSION

This work presents the density-level relation

$$E_d = \rho c^2$$

The equation follows directly from Einstein's mass–energy relation,

$$E=mc^2$$

when both energy and mass are expressed per unit volume.

The definitions

$$E_d=E/V$$

and

$$\rho=m/V$$

lead directly to

$$E_d = \rho c^2$$

The equation therefore establishes a direct correspondence between **mass density** and **rest-mass energy density**.

The central conceptual proposition of this work is that mass–energy equivalence can be viewed not only at the level of total quantities but also at the level of spatial densities.

The equation itself is mathematically a consequence of Einstein's established relation. The future scientific development of this work should therefore focus on whether the density framework can produce new theoretical structures, applications, or experimentally testable predictions.

The proposed concept may consequently serve as a starting point for further research into the relationship between matter distribution and energy distribution.

REFERENCES

1. Einstein, A. (1905). *Does the Inertia of a Body Depend Upon Its Energy-Content?* Annalen der Physik, 18, 639–641.
 2. Einstein, A. (1905). *On the Electrodynamics of Moving Bodies*. Annalen der Physik, 17, 891–921.
 3. Einstein, A. (1916). *The Foundation of the General Theory of Relativity*. Annalen der Physik, 49, 769–822.
 4. Standard texts on special relativity, relativistic energy–momentum, and relativistic continuum physics.
-

AUTHOR

Pawan Upadhyay

Researcher

Research Concept

Mass Density–Energy Density Equivalence

$$E_d = \rho c^2$$

2025-2026

[Expanded Research Paper]

MASS DENSITY–ENERGY DENSITY EQUIVALENCE IN SPACETIME

A Spacetime Formulation of Mass–Energy Equivalence

Author: Pawan Upadhyay


Affiliation: Independent Researcher

Email: pawanupadhyay28@hotmail.com

ORCID iD: <https://orcid.org/0009-0007-9077-5924>

Proposed Theory: Pawan Upadhyay's Pressure Curvature Law of Gravity (PPC Law of Gravity)

Date: 06 Nov. 2025



2025-2026

Abstract

Einstein's mass–energy equivalence establishes the relation $E=mc^2$. The present work develops a density and spacetime formulation of this established relation.

Mass density is defined as

$$\rho=m/V$$

while energy density is defined as

$$[E_d= E/V]$$

Dividing Einstein's mass–energy relation by volume gives

$$[E_d = \rho c^2]$$

The present work extends this relation from three-dimensional space to four-dimensional spacetime using the coordinate representation

$$(x,y,z,ct)$$

The resulting local formulation is

$$E_d(x,y,z,ct) = \rho(x,y,z,ct) c^2$$

This formulation describes the correspondence between mass density and rest-mass energy density at different spacetime locations. The paper examines the mathematical derivation, dimensional consistency, physical interpretation, local formulation, limitations, and possible directions for further research.

The equation $E_d = \rho c^2$ is recognized as a direct density formulation of Einstein's established $E=mc^2$ rather than an independent replacement for it. The potential research contribution lies in developing a systematic spacetime-density framework and determining whether such a framework can lead to additional theoretical or experimentally testable results.

Keywords: mass density, energy density, mass–energy equivalence, spacetime, special relativity, $E=mc^2$, $E_d = \rho c^2$

Chapter 1

Introduction

Mass and energy are fundamental physical quantities. Einstein's mass–energy equivalence established that mass possesses an equivalent rest-energy content.

The familiar relation is

$$[E=mc^2.]$$

The present work considers this relationship from the perspective of **density**.

When mass is distributed through space, it can be represented by mass density:

$$\rho = m/V,$$

Likewise, energy distributed through space can be represented by energy density:

$$E_d = E/V ,$$

The relationship between these quantities follows directly from Einstein's equation:

$$E_d = \rho c^2$$

The next step in the present formulation is to consider the density relationship not merely as a function of spatial position, but as a function of **spacetime**.

A spacetime event can be represented by (x,y,z,ct)

Here x,y,z represent spatial coordinates and ct represents the temporal coordinate expressed in units compatible with spatial coordinates.

Thus, the density relation can be written as **$E_d(x,y,z,ct)=\rho(x,y,z,ct)c^2$**

Chapter 2

Einstein's Mass–Energy Equivalence

Einstein's relation is

$$E=mc^2$$

Here:

- E is rest energy,
- m is mass,
- c is the speed of light in vacuum.

The equation establishes a quantitative relationship between mass and energy.

The present work does not replace this equation. Instead, it examines what happens when both quantities are expressed per unit volume and then considered across spacetime.

Chapter 3

Mass Density and Energy Density

3.1 Mass Density

Mass density is

$$\rho = m/V$$

Its SI unit is

$$\text{kg/m}^3$$

For a spatially varying distribution, mass density may be written as

$$\rho = \rho(x, y, z)$$

When temporal variation is included,

$$\rho = \rho(x, y, z, ct)$$

3.2 Energy Density

Energy density is

$$E_d = E/V$$

Its SI unit is

$$\text{J/m}^3$$

A spacetime-dependent energy density can therefore be represented as

$$E_d = E_d(x, y, z, ct)$$

Chapter 4

Derivation of the Density Relation

Start with Einstein's equation:

$$[E = mc^2.]$$

Divide both sides by volume:

$$E/V = (m/V) c^2$$

Using

$$E_d = E/V$$

and

$$\rho = m/V$$

we obtain

$$E_d = \rho c^2$$

This equation is the central density relation of the present work.

Chapter 5

Spacetime Formulation

5.1 Spacetime Coordinates

A spacetime event may be represented as

(x, y, z, ct)

The first three coordinates describe spatial position:

$[x, y, z]$

The fourth coordinate is

$[ct,]$

where t is time.

The use of ct places the temporal coordinate in length units, making it suitable for four-dimensional spacetime notation.

We may collectively denote the spacetime coordinate by

$[x^\mu = (x, y, z, ct)].$

$$x^\mu = (x, y, z, ct)$$

5.2 Spacetime Mass Density

Mass density may therefore be considered a function of spacetime:

$[\rho(x^\mu)]$

or explicitly,

$[\rho(x, y, z, ct)].$

This means that the mass density may vary according to both spatial position and time.

Mass density may therefore be considered a function of spacetime:

$$\rho(x^\mu)$$

or explicitly,

$$\rho(x, y, z, ct) .$$

This means that the mass density may vary according to both spatial position and time.

5.3 Spacetime Energy Density

Similarly, rest-mass energy density can be written as

$$E_d(x^\mu)$$

$$E_d(x^\mu)$$

or

$$E_d(x,y,z,ct)$$

The density-equivalence relation consequently becomes

$$E_d(x,y,z,ct)=\rho(x,y,z,ct) c^2$$

This is the proposed **spacetime form of the mass density–energy density equivalence relation**.

Chapter 6

Physical Interpretation

The spacetime formulation proposes that wherever a mass density exists, its corresponding rest-mass energy density can be represented through

$$E_d(x,y,z,ct)=\rho(x,y,z,ct) c^2$$

Thus,

$$\rho(x,y,z,ct)$$

describes the distribution of mass density through spacetime, while

$$E_d(x,y,z,ct)$$

describes the corresponding distribution of rest-mass energy density.

The factor c^2 establishes the conversion between the two descriptions.

The conceptual structure can therefore be expressed as

$$[\boxed{\text{Mass density}} \;\xleftrightarrow{c^2}\; \text{Rest-energy density}]$$

$$\text{Mass density} \;\overset{c^2}{\longleftrightarrow}\; \text{Rest-energy density}$$

within a spacetime description.

Chapter 7

Local and Integrated Form

For a continuous mass distribution,

$$[\rho(x,y,z,ct)=\frac{dm}{dV}.]$$

The corresponding rest-energy density is

$$[E_d(x,y,z,ct)$$

$$\frac{dE}{dV}.]$$

Therefore,

$$[\boxed{\frac{dE}{dV}}$$

$$=c^2\frac{dm}{dV}}.]$$

This gives

$$[\boxed{dE=c^2dm}.]$$

Integrating over the relevant spatial volume,

$$[E=c^2\int\rho(x,y,z,ct)dV.]$$

Since

$$[m=\int\rho(x,y,z,ct)dV,]$$

we recover

$$[\boxed{E=mc^2}.]$$

Thus, the spacetime-dependent density formulation remains consistent with the integrated mass–energy relation.

For a continuous mass distribution,

$$\rho(x, y, z, ct) = \frac{dm}{dV} .$$

The corresponding rest-energy density is

$$E_d(x, y, z, ct) = \frac{dE}{dV} .$$

Therefore,

$$\frac{dE}{dV} = c^2 \frac{dm}{dV} .$$

This gives

$$dE = c^2 dm .$$

Integrating over the relevant spatial volume,

$$E = c^2 \int \rho(x, y, z, ct) dV .$$

Since

$$m = \int \rho(x, y, z, ct) dV ,$$

we recover

$$E = mc^2 .$$

Thus, the spacetime-dependent density formulation remains consistent with the integrated mass–energy relation.

Chapter 8

Density as a Spacetime Field

The formulation permits mass density to be considered as a field:

$$[\boxed{\rho = \rho(x,y,z,ct)}]$$

The corresponding rest-energy-density field is

$$[\boxed{E_d = E_d(x,y,z,ct)}]$$

The relation between the two fields is

$$[\boxed{E_d(x,y,z,ct)}$$

$$= c^2 \rho(x,y,z,ct)]$$

This provides a compact mathematical representation of the proposed framework.

The conceptual sequence is therefore:

$$[\boxed{(x,y,z,ct) \rightarrow \rho(x,y,z,ct) \rightarrow E_d(x,y,z,ct)}]$$

where the second transformation is governed by c^2 .

The formulation permits mass density to be considered as a field:

$$\rho = \rho(x, y, z, ct) .$$

The corresponding rest-energy-density field is

$$E_d = E_d(x, y, z, ct) .$$

The relation between the two fields is

$$E_d(x, y, z, ct) = c^2 \rho(x, y, z, ct) .$$

This provides a compact mathematical representation of the proposed framework.

The conceptual sequence is therefore:

$$(x, y, z, ct) \rightarrow \rho(x, y, z, ct) \rightarrow E_d(x, y, z, ct)$$

where the second transformation is governed
by

$$c^2$$

.

Chapter 9

Important Physical Limitation

The quantity E_d in the present equation must be defined carefully.

The equation

$$E_d = \rho c^2$$

represents **rest-mass energy density**.

It should not automatically be identified with the complete energy density of an arbitrary relativistic system.

A physical system may also contain:

- kinetic energy,
- electromagnetic energy,
- thermal energy,

- binding energy,
- pressure,
- momentum-related contributions,
- gravitational-field contributions.

A complete relativistic description of energy and momentum generally requires the **stress–energy tensor**, rather than a single scalar density.

Therefore, the present equation should be viewed as a specific rest-mass-energy relation.

Chapter 10

Relation to Relativity

The use of

$[(x,y,z,ct)]$

places the proposed formulation naturally within a spacetime perspective.

However, merely writing a quantity as a function of spacetime coordinates does not by itself make the quantity Lorentz invariant.

A future relativistic formulation would need to investigate how

$\rho(x,y,z,ct)$

and

$E_d(x,y,z,ct)$

transform between inertial observers.

This distinction is important for developing the proposal into a rigorous relativistic theory.

Chapter 11

Potential Applications

The spacetime density formulation may be investigated in several areas.

11.1 Astrophysics

Spatially and temporally varying matter distributions can be represented through

$$[\rho(x,y,z,ct)]$$

Their corresponding rest-energy-density distribution follows from

$$[E_d = \rho c^2]$$

11.2 Cosmology

Matter density changes with cosmic evolution. A spacetime density representation may provide a useful conceptual framework for studying such distributions.

11.3 Relativistic Fluid Models

The relation may serve as one component of a broader description involving density, pressure, velocity, momentum, and energy.

11.4 Computational Physics

A numerical density field

$$[\rho(x,y,z,t)]$$

can be converted directly into an equivalent rest-energy-density field.

Chapter 12

Future Theoretical Development

The present formulation suggests several possible research directions.

12.1 Lorentz Transformation

Determine how the proposed density quantities transform between inertial frames.

12.2 Stress–Energy Tensor

Investigate how the scalar rest-energy density relates to the $T^{\mu\nu}$ stress–energy tensor.

12.3 Conservation Laws

Investigate the relationship between mass-density evolution, energy-density evolution, and relativistic conservation laws.

12.4 Time-Dependent Density

Develop models of

$[\rho(x,y,z,ct)]$

for evolving physical systems.

12.5 New Predictions

The most important future objective is to determine whether extending the density formulation produces mathematical consequences or experimentally testable predictions that are not already contained in established relativity.

Chapter 13

Conclusion

This work develops a spacetime density formulation of mass–energy equivalence.

Starting from

$[E=mc^2]$

and defining

$$[\rho = m/V]$$

and

$$[E_d = E/V]$$

we obtain

$$[E_d = \rho c^2]$$

When mass density and energy density are allowed to vary through spacetime, the relation becomes

$$[E_d(x,y,z,ct)]$$

$$= \rho(x,y,z,ct) c^2]$$

The spacetime coordinate is represented by

$$x^\mu = (x, y, z, ct)$$

The formulation therefore provides a density-level and spacetime-dependent representation of rest-mass energy equivalence.

The equation itself follows mathematically from Einstein's established $E=mc^2$. The scientific research opportunity lies in developing the spacetime-density framework further—particularly its transformation properties, relationship to the stress–energy tensor, conservation laws, and possible new physical predictions.

REFERENCES

1. Einstein, A. (1905). *Does the Inertia of a Body Depend Upon Its Energy-Content?* Annalen der Physik, 18, 639–641.

2. Einstein, A. (1905). *On the Electrodynamics of Moving Bodies*. Annalen der Physik, 17, 891–921.

3. Einstein, A. (1916). *The Foundation of the General Theory of Relativity*. Annalen der Physik, 49, 769–822.

4. Standard references on special relativity, four-dimensional spacetime, relativistic energy–momentum, and the stress–energy tensor.

AUTHOR

Pawan Upadhyay

Researcher

Proposed Research Concept

Mass Density–Energy Density Equivalence in Spacetime

[$E_d(x,y,z,ct)$

= $\rho(x,y,z,ct) \, c^2$]

$E_d(x,y,z,ct) = E_d(x^\mu)$

2025-2026

Research Paper 2

**SPACETIME ENERGY-DENSITY FORMULATION OF MASS,
MOMENTUM, AND PERFECT-FLUID STRESS–ENERGY**

A Variable Gravitational Pressure Field Formulation

Author: Pawan Upadhyay


Affiliation: Independent Researcher

Email: pawanupadhyay28@hotmail.com

ORCID iD: <https://orcid.org/0009-0007-9077-5924>

Proposed Theory: Pawan Upadhyay's Pressure Curvature Law of Gravity (PPC Law of Gravity)

Date: 06 Nov. 2025



Abstract

This paper presents a spacetime energy-density formulation connecting rest mass, rest energy, energy density, momentum density, gravitational pressure, and the stress–energy tensor of a perfect fluid.

The formulation begins with Einstein's rest-energy relation,

$$E_0 = m_0 c^2 ,$$

and introduces the corresponding density quantities,

$$\rho_0 = \frac{m_0}{V} , \quad E_{0d} = \frac{E_0}{V} .$$

Consequently,

$$E_{0d} = \rho_0 c^2 .$$

The quantities are represented over spacetime using

$$x^\mu = (x, y, z, ct) .$$

The energy density and mass density are therefore treated as spacetime-dependent quantities,

$$E_d = E_d(x^\mu), \quad \rho_0 = \rho_0(x^\mu).$$

Momentum density is defined by

$$\mathbf{p}_d = \frac{\mathbf{p}}{V} ,$$

with

$$p_d^2 = p_{dx}^2 + p_{dy}^2 + p_{dz}^2 .$$

The relativistic energy–momentum relation can, under consistent fixed-volume definitions, be expressed in density form as

$$E_d^2 = \rho_0^2 c^4 + p_d^2 c^2 .$$

A variable equation of state is introduced for the **gravitational pressure field**:

$$P_g(x^\mu) = \omega(x^\mu) E_d(x^\mu) .$$

The parameter

ω

is allowed to vary throughout spacetime.

Important special cases are

$$\omega = 0, \quad \omega = \frac{1}{3}, \quad \omega = -1, \quad \omega = 1.$$

In particular, the proposed near-black-hole regime is characterized by

$$\omega \approx 1 ,$$

which gives

$$P_g \approx E_d .$$

The gravitational pressure field is incorporated into the perfect-fluid stress–energy tensor,

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu} ,$$

which is related to spacetime curvature through Einstein's equation.

1. Introduction

Einstein's mass–energy relation is

$$E_0 = m_0 c^2 .$$

This equation establishes the equivalence between rest mass and rest energy.

The present paper develops a density formulation of this relation. Instead of considering only the total mass and total energy of a system, the formulation considers their corresponding densities.

For a system occupying volume

V

,

$$\rho_0 = \frac{m_0}{V}$$

is the rest-mass density and

$$E_{0d} = \frac{E_0}{V}$$

is the rest-energy density.

Substitution of

$$E_0 = m_0 c^2$$

gives

$$E_{0d} = \frac{m_0 c^2}{V} = \rho_0 c^2.$$

Therefore,

$$E_{0d} = \rho_0 c^2 .$$

The formulation is subsequently extended to momentum density and the perfect-fluid stress–energy tensor.

2. Spacetime Coordinate

The spacetime coordinate used in this formulation is

$$x^\mu = (x, y, z, ct) .$$

Here,

$$x, \quad y, \quad z$$

represent the three spatial coordinates and

$$ct$$

represents the temporal coordinate expressed in length units.

The physical quantities may therefore be represented as spacetime-dependent fields:

$$\rho_0 = \rho_0(x, y, z, ct)$$

$$\rho_0 = \rho_0(x, y, z, ct)$$

and

$$E_d = E_d(x, y, z, ct) .$$

The rest-energy density relation becomes

$$E_{0d}(x, y, z, ct) = \rho_0(x, y, z, ct)c^2 .$$

In compact notation,

$$E_{0d}(x^\mu) = \rho_0(x^\mu)c^2 .$$

3. Mass Density and Energy Density

The mass density is defined as

$$\rho = \frac{m}{V} .$$

For rest mass,

$$\rho_0 = \frac{m_0}{V} .$$

The energy density is defined as

$$E_d = \frac{E}{V} .$$

For rest energy,

$$E_{0d} = \frac{E_0}{V} .$$

Therefore,

$$E_{0d} = \rho_0 c^2 .$$

This relation represents the conversion of rest-mass density into rest-energy density through the factor

$$c^2$$

.

4. Rest Frame

For a physical system in its rest frame,

$$\mathbf{p} = 0 .$$

Consequently, its momentum density is

$$\mathbf{p}_d = 0 .$$

Thus,

$$p_d = 0 .$$

The energy density then reduces to the rest-energy density in the simplified rest-mass-energy description:

$$E_{0d} = \rho_0 c^2 .$$

Hence,

$$p_d = 0 \quad \Longrightarrow \quad E_{0d} = \rho_0 c^2 .$$

5. Momentum Density

Momentum density is defined as

$$\mathbf{p}_d = \frac{\mathbf{p}}{V} .$$

Its three spatial components are

$$\mathbf{p}_d = (p_{dx}, p_{dy}, p_{dz}) .$$

The magnitude of the momentum-density vector is

$$p_d^2 = p_{dx}^2 + p_{dy}^2 + p_{dz}^2 .$$

Therefore,

$$p_d = \sqrt{p_{dx}^2 + p_{dy}^2 + p_{dz}^2} .$$

This gives the spatial decomposition of the momentum-density quantity used in the formulation.

6. Energy–Momentum Relation in Density Form

The relativistic energy–momentum relation is

$$E^2 = p^2 c^2 + m_0^2 c^4 .$$

For quantities associated with the same physical system and volume

V

, divide by

V^2

:

$$\frac{E^2}{V^2} = c^2 \frac{p^2}{V^2} + c^4 \frac{m_0^2}{V^2} .$$

Using

$$E_d = \frac{E}{V} ,$$

$$p_d = \frac{p}{V},$$

and

$$\rho_0 = \frac{m_0}{V},$$

gives

$$E_d^2 = p_d^2 c^2 + \rho_0^2 c^4 .$$

Using the three spatial components,

$$E_d^2 = c^2 (p_{dx}^2 + p_{dy}^2 + p_{dz}^2) + \rho_0^2 c^4 .$$

The zero-momentum limit is

$$p_d = 0,$$

giving

$$E_{0d} = \rho_0 c^2 .$$

Thus the rest-energy density relation appears as the zero-momentum limit of the density-scaled relativistic relation.

7. General Relativistic Fluid Region

Consider a spacetime region

$$\mathcal{R}_{\text{GRF}} \subset \mathcal{M}$$

called the **general relativistic fluid region**.

General Relativistic Fluid Region

Within this region,

$$x^\mu = (x, y, z, ct)$$

and the relevant quantities are

$$\{\rho_0(x^\mu), E_d(x^\mu), p_d(x^\mu), P_g(x^\mu), \omega(x^\mu)\}.$$

The energy density is treated as a local quantity,

$$E_d = E_d(x^\mu) .$$

The gravitational pressure is likewise a local quantity,

$$P_g = P_g(x^\mu) .$$

The mass density may also be represented as a spacetime-dependent field,

$$\rho_0 = \rho_0(x^\mu) .$$

The momentum-density magnitude is

$$p_d = p_d(x^\mu)$$

with

$$p_d^2 = p_{dx}^2 + p_{dy}^2 + p_{dz}^2 .$$

The variable equation-of-state parameter is

$$\omega = \omega(x^\mu) .$$

The gravitational pressure field is related to the energy-density field by

$$P_g(x^\mu) = \omega(x^\mu) E_d(x^\mu) .$$

For the proposed near-black-hole regime,

$$\omega(x^\mu) \approx 1$$

and therefore

$$P_g(x^\mu) \approx E_d(x^\mu) .$$

Thus, the local quantities in the general relativistic fluid region are collectively represented as

$$\{\rho_0(x^\mu), E_d(x^\mu), p_d(x^\mu), P_g(x^\mu), \omega(x^\mu)\}.$$

8. Gravitational Pressure Field

The formulation defines

$$P_g$$

as **gravitational pressure**.

Its spacetime distribution is called the
gravitational pressure field .

Thus,

$$P_g = P_g(x, y, z, ct) .$$

The gravitational pressure field is related to
the local energy density through

$$P_g(x^\mu) = \omega(x^\mu) E_d(x^\mu) .$$

The three quantities are allowed to vary:

$$P_g = P_g(x^\mu)$$

$$E_d = E_d(x^\mu)$$

$$\omega = \omega(x^\mu) .$$

Therefore, the equation of state is itself a spacetime-dependent relation:

$$P_g(x, y, z, ct) = \omega(x, y, z, ct) E_d(x, y, z, ct) .$$

9. Variable Equation-of-State Parameter

The dimensionless parameter

ω

describes the local ratio

$$\omega = \frac{P_g}{E_d}$$

where

$$E_d \neq 0$$

.

In the general formulation,

$$\omega = \omega(x, y, z, ct) .$$

Thus, different regions of the general relativistic fluid region may have different values of

ω

.

The formulation does not require

ω

to be constant.

10. Special Values of

ω

Several important limiting cases are represented by particular values of

ω

.

10.1 Pressureless Matter

For

$$\omega = 0 ,$$

we obtain

$$P_g = 0 .$$

This represents the pressureless-matter or dust limit.

10.2 Radiation

For

$$\omega = \frac{1}{3} ,$$

we obtain

$$P_g = \frac{1}{3} E_d .$$

This is the standard radiation equation of state.

10.3 Vacuum Energy

For

$$\omega = -1 ,$$

we obtain

$$P_g = -E_d .$$

This corresponds to the vacuum-energy/
cosmological-constant-type equation of state.

10.4 Stiff-Fluid Limit

For

$$\omega = 1 ,$$

we obtain

$$P_g = E_d .$$

This is the stiff-fluid equation of state.

11. Near-Black-Hole Regime

The proposed formulation considers a particular limiting regime in a general relativistic fluid region near a black hole.

The proposed assumption is

$$\omega \approx 1 .$$

Using

$$P_g = \omega E_d,$$

we obtain

$$P_g = \omega E_d \approx E_d.$$

Therefore,

$$P_g \approx E_d .$$

This represents the proposed **near-black-hole gravitational-pressure condition**.

The corresponding relation can be written as

$$\omega \rightarrow 1 \quad \Longrightarrow \quad P_g \rightarrow E_d .$$

This assumption means that, within the proposed model, the gravitational pressure field becomes approximately equal in magnitude to the local energy density in this regime.

It is important to state that

$$\omega \approx 1$$

near a black hole is a **modeling hypothesis of this formulation**, rather than a general consequence of standard general relativity.

12. Perfect-Fluid Stress–Energy Tensor

For a perfect fluid, using the $(-+++)$ metric convention and

$$u^\mu u_\mu = -1,$$

the stress–energy tensor is

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu} .$$

Substituting

$$P_g = \omega E_d$$

gives

$$T_{\mu\nu} = (1 + \omega)E_d u_\mu u_\nu + \omega E_d g_{\mu\nu} .$$

When

ω

varies over spacetime,

$$P_g(x^\mu) = \omega(x^\mu) E_d(x^\mu),$$

the perfect-fluid tensor is

$$T_{\mu\nu} = [E_d(x^\mu) + P_g(x^\mu)] u_\mu u_\nu + P_g(x^\mu) g_{\mu\nu} .$$

13. Near-Black-Hole Stress–Energy Form

For the proposed limit

$$\omega \approx 1 ,$$

the perfect-fluid tensor becomes

$$T_{\mu\nu} \approx (1 + 1)E_d u_\mu u_\nu + E_d g_{\mu\nu}.$$

Therefore,

$$T_{\mu\nu} \approx 2E_d u_\mu u_\nu + E_d g_{\mu\nu} .$$

Equivalently,

$$T_{\mu\nu} \approx E_d (2u_\mu u_\nu + g_{\mu\nu}) .$$

$$T_{\mu\nu} \approx E_d (2u_\mu u_\nu + g_{\mu\nu}) .$$

This is the stress–energy form corresponding to the proposed stiff-fluid limit

$$P_g \approx E_d.$$

14. Local Rest Frame

In the local rest frame of the perfect fluid,
using

$$c = 1$$

normalization,

$$u^{\hat{\mu}} = (1, 0, 0, 0) .$$

The perfect-fluid tensor takes the form

$$T_{\hat{\mu}\hat{\nu}} = \begin{pmatrix} E_d & 0 & 0 & 0 \\ 0 & P_g & 0 & 0 \\ 0 & 0 & P_g & 0 \\ 0 & 0 & 0 & P_g \end{pmatrix} .$$

Under the proposed near-black-hole condition,

$$P_g \approx E_d,$$

Under the proposed near-black-hole condition,

$$P_g \approx E_d,$$

so

$$T_{\hat{\mu}\hat{\nu}} \approx \begin{pmatrix} E_d & 0 & 0 & 0 \\ 0 & E_d & 0 & 0 \\ 0 & 0 & E_d & 0 \\ 0 & 0 & 0 & E_d \end{pmatrix}.$$

Thus, in this proposed limiting model, the pressure components approximately equal the energy-density component.

15. Einstein Field Equation

The stress–energy tensor is related to spacetime curvature through Einstein's equation:

$$G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} .$$

Substituting the perfect-fluid tensor gives

$$G_{\mu\nu} = \frac{8\pi G}{c^4} [(E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu}] .$$

Using

$$P_g = \omega E_d,$$

we obtain

$$G_{\mu\nu} = \frac{8\pi G}{c^4} E_d [(1 + \omega)u_\mu u_\nu + \omega g_{\mu\nu}] .$$

For the proposed near-black-hole limit,

$$\omega \approx 1,$$

this becomes

$$G_{\mu\nu} \approx \frac{8\pi G}{c^4} E_d (2u_\mu u_\nu + g_{\mu\nu}) .$$

16. Weak-Field Regime

In a weak-field regime, the spacetime metric can be represented approximately as

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} , \quad |h_{\mu\nu}| \ll 1.$$

The matter-energy quantities remain locally defined by

$$E_{0d} = \rho_0 c^2$$

and

$$P_g = \omega E_d .$$

Different values of

ω

represent different fluid regimes.

For example,

$$\omega \approx 0$$

gives

$$P_g \approx 0,$$

while

$$\omega = \frac{1}{3}$$

gives

$$P_g = \frac{E_d}{3}.$$

The weak-field approximation is appropriate when the deviation of the spacetime metric from the Minkowski metric is sufficiently small.

17. Strong-Field Regime

In a strong-field regime, the perturbative condition

$$|h_{\mu\nu}| \ll 1$$

cannot generally be assumed.

The full stress–energy tensor must therefore be considered:

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu} .$$

The gravitational pressure field becomes an important component of the relativistic fluid description.

The proposed near-black-hole condition is

$$\omega \approx 1$$

and consequently

$$P_g \approx E_d .$$

Thus, within the proposed model, the gravitational pressure field becomes comparable to the local energy density in the near-black-hole regime.

18. Unified Mathematical Framework

The complete formulation can be summarized as follows.

Spacetime

$$x^\mu = (x, y, z, ct)$$

Rest-mass density

$$\rho_0 = \frac{m_0}{V}$$

Rest-energy density

$$E_{0d} = \frac{E_0}{V}$$

Mass–energy density relation

$$E_{0d} = \rho_0 c^2$$

Energy density

$$E_d = E_d(x^\mu)$$

Momentum density

$$\mathbf{p}_d = (p_{dx}, p_{dy}, p_{dz})$$

Momentum-density magnitude

$$p_d^2 = p_{dx}^2 + p_{dy}^2 + p_{dz}^2$$

Density-scaled energy–momentum relation

$$E_d^2 = \rho_0^2 c^4 + p_d^2 c^2$$

under the stated common-volume interpretation.

Gravitational pressure field

$$P_g = P_g(x^\mu)$$

Variable equation of state

$$P_g(x^\mu) = \omega(x^\mu) E_d(x^\mu)$$

Special near-black-hole assumption

$$\omega \approx 1$$

therefore

$$P_g \approx E_d .$$

Perfect-fluid stress–energy tensor

$$T_{\mu\nu} = (E_d + P_g) u_\mu u_\nu + P_g q_{\mu\nu}$$

19. Conceptual Flow of the Formulation

The proposed framework can be represented by

$$m_0 \rightarrow \rho_0 = \frac{m_0}{V} \rightarrow E_{0d} = \rho_0 c^2$$

followed by

$$E_d \rightarrow P_g = \omega E_d .$$

The pressure and energy density then enter

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu}$$

and

$$T_{\mu\nu} \rightarrow G_{\mu\nu} .$$

20. Discussion

The central relation of the formulation is

$$E_{0d}(x^\mu) = \rho_0(x^\mu)c^2 .$$

It expresses the rest-mass/rest-energy equivalence in density form.

The momentum-density vector is

$$\mathbf{p}_d = (p_{dx}, p_{dy}, p_{dz})$$

and its magnitude satisfies

$$p_d^2 = p_{dx}^2 + p_{dy}^2 + p_{dz}^2 .$$

The corresponding density-scaled relativistic relation is

$$E_d^2 = \rho_0^2 c^4 + p_d^2 c^2 .$$

No separate internal-energy-density term is required in this formulation unless the research question specifically requires a decomposition of E_d into rest-mass and internal-energy contributions.

21. Physical Scope and Qualification

The relation

$$E_{0d} = \rho_0 c^2$$

follows directly from Einstein's rest-energy relation and the definitions of density.

The relation

$$P_g = \omega E_d$$

is an equation-of-state parameterization. It specifies how the gravitational pressure field is related to the local energy density.

The assumption

$$\omega \approx 1$$

in a near-black-hole general relativistic fluid region is a **hypothesis of the proposed model**. It is not, by itself, a result derived from standard general relativity.

To establish this as a new physical theory, the model would need additional dynamical equations determining

$$E_d$$

To establish this as a new physical theory, the model would need additional dynamical equations determining

$$E_d$$

,

$$\omega$$

,

$$u^\mu$$

, and

$$g_{\mu\nu}$$

, together with conservation laws and quantitative predictions that can be compared with established relativistic models and observations.

22. Conclusion

This paper presents a spacetime density formulation based on the following principal equations:

$$x^\mu = (x, y, z, ct)$$

$$\rho_0 = \frac{m_0}{V}$$

$$E_{0d} = \frac{E_0}{V}$$

$$E_{0d} = \rho_0 c^2$$

$$p_d^2 = p_{dx}^2 + p_{dy}^2 + p_{dz}^2$$

$$E_d^2 = \rho_0^2 c^4 + p_d^2 c^2$$

$$P_g = \omega E_d$$

and

$$T_{\mu\nu} = (E_d + P_g)u_\mu u_\nu + P_g g_{\mu\nu} .$$

The gravitational pressure field is allowed to vary through

$$P_g = P_g(x, y, z, ct)$$

and the equation-of-state parameter is allowed to vary through

$$\omega = \omega(x, y, z, ct) .$$

For the proposed near-black-hole regime,

$$\omega \approx 1$$

For the proposed near-black-hole regime,

$$\omega \approx 1$$

and consequently

$$P_g \approx E_d .$$

The corresponding perfect-fluid stress–energy tensor becomes approximately

$$T_{\mu\nu} \approx E_d (2u_\mu u_\nu + g_{\mu\nu}) .$$

The resulting framework connects rest-mass density, rest-energy density, energy density, momentum density, the gravitational pressure field, perfect-fluid stress–energy, and spacetime curvature within a single mathematical structure.

Keywords

**Rest Mass Density • Rest Energy Density •
Energy Density • Momentum Density •
Gravitational Pressure • Gravitational
Pressure Field • Variable Equation of State •
Perfect Fluid • General Relativistic Fluid
Region • Stress–Energy Tensor • Spacetime •
Weak-Field Regime • Strong-Field Regime •
Black Hole • General Relativity • Einstein
Equation**