

HKDSE Mathematics Compulsory Part — Paper 2 (Mock)

PAPER 2

75 minutes (1.25 hours)

INSTRUCTIONS

1. Read the instructions on the Answer Sheet carefully. Stick a barcode label after the start of the examination.
2. All questions carry equal marks.
3. ANSWER ALL QUESTIONS. Use an HB pencil to mark all answers on the Answer Sheet.
4. Mark only ONE answer for each question. More than one answer scores NO MARKS.
5. No marks will be deducted for wrong answers.

Not to be taken away before the
end of the examination session

There are 30 questions in Section A and 15 questions in Section B.

The diagrams in this paper are not necessarily drawn to scale. Choose the best answer for each question.

Section A

1. Simplify $\left(\frac{a^5b^{-3}}{a^{-1}b^2}\right)$.

A. $\frac{a^6}{b}$

C. $\frac{a^4}{b^5}$

B. $\frac{a^6}{b^5}$

D. a^6b^5
2. Express 0.0000807 in scientific notation.

A. $\$8.07 \times 10^{-4}$

C. $\$8.07 \times 10^5$

B. $\$80.7 \times 10^{-6}$

D. $\$8.07 \times 10^{-5}$
3. Solve $6(x - 3) = 3x + 6$.

A. $x = 6$

C. $x = 4$

B. $x = 24$

D. $x = 8$
4. Simplify $\left(\frac{x^2 - 49}{x^2 - 3x - 28}\right)$.

A. $\frac{x - 7}{x + 4}$

C. $\frac{x - 7}{x - 4}$

B. $\frac{x + 7}{x - 4}$

D. $\frac{x + 7}{x + 4}$
5. Factorize $216a^3 - 125$.

A. $\$(6a - 5)(36a^2 - 30a + 25)$

C. $\$(6a - 5)(36a^2 + 30a + 25)$

B. $\$(6a + 5)(36a^2 - 30a + 25)$

D. $\$(6a - 5)(6a^2 + 30a + 25)$
6. If $y = \left(\frac{3x + 7}{x - 4}\right)$, express x in terms of y .

A. $\$x = \frac{4y - 7}{y - 3}$

C. $\$x = \frac{7 - 4y}{y - 3}$

B. $\$x = \frac{4y + 7}{y + 3}$

D. $\$x = \frac{4y + 7}{y - 3}$
7. Solve $3^x = 80$, correct to 3 significant figures.

A. 3.99

C. 1.90

B. 3.98

D. 26.7
8. Let α and β be the roots of $3x^2 - 11x + 2 = 0$. Find $\alpha^2 + \beta^2$.

A. $\frac{121}{9}$


C. $\frac{115}{9}$

B. $\frac{109}{9}$


D. $\frac{97}{9}$

9. If the equation $x^2 - kx + (5k - 16) = 0$ has two equal real roots, find k .
- A. $k = -4$ or $k = -16$ B. $k = 16$ only
 C. $k = 8$ or $k = 2$ D. $k = 4$ or $k = 16$
10. Solve $x^2 - 3x - 28 \leq 0$.
- A. $x \leq -4$ or $x \geq 7$ B. $-4 \leq x \leq 7$
 C. $-7 \leq x \leq 4$ D. $4 \leq x \leq 7$
11. The 2nd and 6th terms of an arithmetic sequence are 11 and 39 respectively. Find the sum of the first 16 terms.
- A. 880 B. 904
 C. 928 D. 452
12. The geometric sequence is 6, 12, 24, (...). Find the 8th term.
- A. 384 B. 768
 C. 1536 D. 512
13. Find the remainder when $4x^3 - 2x^2 + 5x - 3$ is divided by $x + 1$.
- A. 4 B. 14
 C. -6 D. -14
14. It is given that y varies inversely as x^2 . When $x = 2$, $y = 25$. Find the value of y when $x = 5$.
- A. 4 B. 10
 C. 62.5 D. 100
15. An item is given successive discounts of 20% and 25%. Find the overall percentage discount.
- A. 45% B. 60%
 C. 40% D. 5%
16. The sequence is 10, 17, 24, 31, (...). Find the 16th term.
- A. 122 B. 108
 C. 115 D. 112
17. \$25000 is deposited at 8% per annum compounded yearly for 2 years. Find the amount, correct to the nearest dollar.
- A. \$29000 B. \$27000
 C. \$28800 D. \$29160
18. Simplify $(\sqrt{128}) - (\sqrt{32})$.
- A. $4\sqrt{2}$ B. $12\sqrt{2}$
 C. $9\sqrt{6}$ D. $96\sqrt{2}$


19. In the figure, in $(\triangle ABC)$, $(\angle A) = 47^\circ$ and $(\angle B) = 58^\circ$. Find $(\angle ACD)$, where D lies on BC produced.



- A. 75°
 B. 105°
 C. 122°
 D. 90°
20. In the figure, $AB \parallel CD$ and a transversal cuts them. The marked angle is 115° . Find x .



- A. 115°
 B. 57.5°
 C. 65°
 D. 25°
21. In the figure, a sector has radius 21 cm and angle at the centre 120° . Find the length of the arc in terms of (π) .



- A. $7\pi \text{ cm}$
 B. $14\pi \text{ cm}$
 C. $147\pi \text{ cm}$
 D. $42\pi \text{ cm}$
22. Find the distance between the points $(2, 6)$ and $(10, 21)$.
- A. 17
 B. 23
 C. $\sqrt{23}$
 D. 289

23. Find the slope of a line perpendicular to the line $8x - 3y = 5$.

A. $\frac{8}{3}$

B. $-\frac{3}{8}$

C. $\frac{3}{8}$

D. $-\frac{8}{3}$

24. Find the centre and radius of the circle $x^2 + y^2 + 8x - 14y - 16 = 0$.


A. centre $(4, -7)$, $r = 9$

B. centre $(-4, 7)$, $r = 81$

C. centre $(-8, 14)$, $r = 9$

D. centre $(-4, 7)$, $r = 9$

25. The figure shows a regular 9-sided polygon. Find each interior angle (θ) .




A. 40°

B. 144°

C. 135°

D. 140°

26. In the figure, $DE \parallel BC$. It is given that $AD = 9$, $DB = 6$ and $AE = 15$. Find EC .



A. 10

B. 22.5

C. 6

D. 15

27. Find the volume of a sphere of radius 6 cm in terms of (π) .

A. $72\pi \text{ cm}^3$

B. $288\pi \text{ cm}^3$

C. $216\pi \text{ cm}^3$

D. $36\pi \text{ cm}^3$

28. If the mean of $11, 17, x, 24$ is 20 , find x .

A. 20

B. 24

C. 28

D. 26

29. Find the median of the data 5, 8, 11, 13, 17, 20 .

A. 12

B. 11

C. 13

D. 12.5

30. A fair die is thrown. Find the probability of getting an even number .

A. $\frac{1}{3}$

B. $\frac{2}{3}$

C. $\frac{1}{2}$

D. $\frac{1}{6}$

Section B

31. Let $a > b$, where a and b are real numbers. I. $a^2 > b^2$ II. $a - b > 0$ III. $-a < -b$ Which of the above is/are true?

A. I and II only

B. I and III only

C. II and III only

D. I, II and III

32. Let x and y be positive numbers. I. $\log \frac{x}{y} = \log x - \log y$ II.

$\log(x + y) = \log x \times \log y$ III. $\log(x^4) = 4 \log x$ Which of the above is/are true?

A. I and II only

B. I and III only

C. II and III only

D. I, II and III

33. Consider the quadratic function $y = x^2 - 14x + 58$. I. The vertex of the graph is (7, 9) II. The graph opens upwards III. The graph cuts the x -axis at two points Which of the above is/are true?


A. I and II only

B. I and III only

C. II and III only

D. I, II and III

34. Let $ABCD$ be a parallelogram. I. Both pairs of opposite angles are equal II. The diagonals are perpendicular to each other III. $AB \parallel DC$ and $AD \parallel BC$ Which of the above is/are true?




A. I and II only

B. I and III only

C. II and III only

D. I, II and III

35. In the figure, $\overline{AB}$ is a diameter of the circle O and C is a point on the circle with $\angle BAC = 27^\circ$. Find $\angle ABC$.



- A. 63° B. 27°
C. 90° D. 54°
36. Consider the sequence **6, 24, 96, 384, ...**. I. It is a geometric sequence with common ratio **4** II. It is an arithmetic sequence III. The **5**th term is **1536** Which of the above is/are true?
- A. I and II only B. I and III only
C. II and III only D. I, II and III
37. Each datum of a set of data is multiplied by **4** and then decreased by **6**. If the standard deviation of the original data is **9**, find the standard deviation of the new data.
- A. **30** B. **9**
C. **36** D. **42**
38. If $C_2^n = 105$, find n .
- A. **14** B. **105**
C. **15** D. **16**
39. Let **A** and **B** be mutually exclusive events with $P(A) = 0.31$ and $P(B) = 0.24$. Find $P(A \setminus (A \cup B))$.
- A. **0.0744** B. **0.07**
C. **0.55** D. **0.4756**
40. **7** different items are arranged in a row such that **2** particular items must be next to each other. Find the number of different arrangements.
- A. **10080** B. **720**
C. **5040** D. **1440**
41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. **44** B. **150**
C. **105** D. **45**

36. Consider the sequence **6, 24, 96, 384, ...**. I. It is a geometric sequence with common ratio **4** II. It is an arithmetic sequence III. The **5**th term is **1536** Which of the above is/are true?
- A. I and II only
B. I and III only
C. II and III only
D. I, II and III
37. Each datum of a set of data is multiplied by **4** and then decreased by **6**. If the standard deviation of the original data is **9**, find the standard deviation of the new data.
- A. **30**
B. **9**
C. **36**
D. **42**
38. If $C_2^n = 105$, find n .
- A. **14**
B. **105**
C. **15**
D. **16**
39. Let A and B be mutually exclusive events with $P(A) = 0.31$ and $P(B) = 0.24$. Find $P(A \cup B)$.
- A. **0.0744**
B. **0.07**
C. **0.55**
D. **0.4756**
40. **7** different items are arranged in a row such that **2** particular items must be next to each other. Find the number of different arrangements.
- A. **10080**
B. **720**
C. **5040**
D. **1440**
41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. **44**
B. **150**
C. **105**
D. **45**

37. Each datum of a set of data is multiplied by **4** and then decreased by **6**. If the standard deviation of the original data is **9**, find the standard deviation of the new data.
- A. **30** B. **9**
C. **36** D. **42**
38. If $C_2^n = 105$, find n .
- A. **14** B. **105**
C. **15** D. **16**
39. Let **A** and **B** be mutually exclusive events with $P(A) = 0.31$ and $P(B) = 0.24$. Find $P(A \setminus (\cup B))$.
- A. **0.0744** B. **0.07**
C. **0.55** D. **0.4756**
40. **7** different items are arranged in a row such that **2** particular items must be next to each other. Find the number of different arrangements.
- A. **10080** B. **720**
C. **5040** D. **1440**
41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. **44** B. **150**
C. **105** D. **45**

38. If $C_2^n = 105$, find n .
- A. 14
B. 105
C. 15
D. 16
39. Let A and B be mutually exclusive events with $P(A) = 0.31$ and $P(B) = 0.24$. Find $P(A \setminus (A \cup B))$.
- A. 0.0744
B. 0.07
C. 0.55
D. 0.4756
40. 7 different items are arranged in a row such that 2 particular items must be next to each other. Find the number of different arrangements.
- A. 10080
B. 720
C. 5040
D. 1440
41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. 44
B. 150
C. 105
D. 45

39. Let A and B be mutually exclusive events with $P(A) = 0.31$ and $P(B) = 0.24$. Find $P(A \setminus (A \cup B))$.
- A. 0.0744
B. 0.07
C. 0.55
D. 0.4756
40. 7 different items are arranged in a row such that 2 particular items must be next to each other. Find the number of different arrangements.
- A. 10080
B. 720
C. 5040
D. 1440
41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. 44
B. 150
C. 105
D. 45

40. 7 different items are arranged in a row such that 2 particular items must be next to each other. Find the number of different arrangements.
- A. 10080 B. 720
C. 5040 D. 1440
41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. 44 B. 150
C. 105 D. 45

41. The n th term of a sequence is $T(n) = 4n^2 + n$. Find $T(6) - T(5)$.
- A. 44 B. 150
C. 105 D. 45

