

# Chapter 1

## Logic and sets

### 1.1 Logical formulas

1. Prove that the following sentences are true:

1.  $\neg(\neg p) \Leftrightarrow p$

2.  $p \vee \neg p$

3.  $\neg(p \wedge q) = (\neg p) \vee (\neg q)$

4.  $\neg(p \vee q) = (\neg p) \wedge (\neg q)$

5.  $(p \Rightarrow q) \Leftrightarrow (\neg p) \vee q$

6.  $0 \Rightarrow 1$

### 1.2 Basic set theory

#### 1.2.1 Rough ideas

2. Prove that  $\{1, 1, 2, 2, 2\} = \{1, 2\}$

3. Let  $X$  be a set built from all sets such that  $A \notin A$ . Prove that  $X$  does not exist. Hint: what if  $X \in X$ ? What if  $X \notin X$ ?

#### 1.2.2 A few ways of constructing new sets

4. Let  $A, B, C$  be sets. Prove that:

1.  $A \cup A = A$

2.  $A \cup B = B \cup A$

3.  $A \cup (B \cup C) = (A \cup B) \cup C$

4.  $A \cap A = A$

5.  $A \cap B = B \cap A$

6.  $A \cap (B \cap C) = (A \cap B) \cap C$
7.  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
8.  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

5. Prove that there is no set of all sets. Hint: assume there is one. Then you can select some sets to form a set that does not exist.

### 1.2.3 Subsets and complements

6. Prove that  $A = B$  iff  $A \subseteq B \wedge B \subseteq A$ .
7. Prove that for any set  $A$ ,  $\emptyset \subseteq A$ .
8. Let  $\emptyset$  and  $\emptyset'$  be empty sets. Prove that  $\emptyset = \emptyset'$ .

9. Prove the following set identities:

1. Let  $A \subseteq B$ . Prove that  $(A^c)^c = A$ .
2. Let  $A, B \subset U$ . Prove that  $(A \cup B)^c = A^c \cap B^c$
3. Let  $A, B \subset U$ . Prove that  $(A \cap B)^c = A^c \cup B^c$
4.  $\{a \in A : a \in B\} = \{b \in B : b \in A\}$

10.

1. Let  $A = \{1, 2, 3\}$ . Find  $2^A$ . What is the number of elements in  $2^A$ ? How is it related to the number of elements of  $A$ ?
2. Let  $A$  be a finite set with  $n$  elements. Using the approach in which you choose which elements belong to a subset, prove that  $2^A$  has  $2^n$  elements.

### 1.2.4 Infinite collections of sets

11. Prove that for finite  $I$  these definitions agree with the previous.

12. Let  $A_i \subseteq U$ ,  $i \in I$  and

$$\sigma = \bigcup_{i \in I} A_i, \pi = \bigcap_{i \in I} A_i$$

Prove that:

1. if  $k \in I$ , then  $A_k \cup \sigma = \sigma$
2.  $\pi \subseteq \sigma$
3.  $\sigma \cap \pi = \pi$

13. Find sum and intersection of family of subsets of  $\mathbb{R}$ :  $A_r = \{r, -r\}$  for  $r \geq 0$ .

14. Let  $A \subseteq X_i \subseteq U$  for  $i \in I$ . Prove that  $A \subseteq \bigcup_{i \in I} X_i$
15. For every point  $a \in A$  there is a set  $U_a \subseteq A$  such that  $a \in U_a$ . Prove that

$$A = \bigcup_{a \in A} U_a.$$

### 1.2.5 Cartesian product

16. Let  $A = \{\{a\}, \{a, b\}\}$ ,  $B = \{\{c\}, \{c, d\}\}$ . Prove that  $A = B$  iff  $a = c \wedge b = d$ . Such a set  $A$  we call **the ordered pair**  $(a, b)$  as it has the property  $(a, b) = (c, d)$  iff  $a = c$  and  $b = d$ . Now you can forget how it has been constructed, and just remember this property.
17. Prove that  $(a, (b, c)) = (d, (e, f))$  iff  $a = d \wedge b = e \wedge c = f$ .
18. Check that defining  $(a, b, c)$  as  $((a, b), c)$  also works (so two ordered tuples are the same if they have the same first element, the same second element, ...)
19. Check that, in terms of sets,  $(a, (b, c)) \neq ((a, b), c)$ , so formally we do need to stick to one convention. However as we are interested in the property of ordered tuple, we will not distinguish them and denote both of them just as  $(a, b, c)$ . Such notational problems appear in various places in mathematics, so we need to try to get used to them.
20. Do you remember the identification of  $(a, (b, c))$  and  $((a, b), c)$ ? Prove that  $A \times (B \times C) = (A \times B) \times C$ . Therefore we'll write it just as  $A \times B \times C$  without parentheses.

## 1.3 Natural numbers and mathematical induction

21. You can prove that  $2^n > n$  for every natural number  $n$ .
1. Prove that the formula works for  $n = 0$  (punch the first domino).
  2. Assume that for some  $n$  you proved on some way that  $2^n > n$ . Using this, prove that  $2^{n+1} > (n + 1)$  (if  $n$ -th domino falls, then  $n + 1$ -th domino also falls)
- 22.
1. Prove<sup>1</sup> that 6 divides  $n^3 - n$  for all natural  $n$ .
  2. Prove<sup>2</sup> that 6 divides  $n^3 - n$  for all integers  $n$ . You can use a slight modification mathematical induction principle proving the implication „if the theorem works for  $n$ , it works also for  $n - 1$ “.
23. (Bernoulli's inequality) Prove that for real  $x > -1$  and natural  $n \geq 1$ , the following inequality holds:

$$(1 + x)^n \geq 1 + nx.$$

---

<sup>1</sup>Another method is to notice that  $n^3 - n = (n - 1) \cdot n \cdot (n + 1)$ . Why 2 does divide it? Why 3?

<sup>2</sup>How  $n^3 - n$  and  $(-n)^3 - (-n)$  are related? Does it simplify the proof?

- 24.** In Mathsland there are  $n \geq 2$  cities. Between each pair of them there is a one-way road.
1. Prove that there is a city from which you can drive to all the other cities. Hint: assume that the hypothesis works for some  $n$  and any country with  $n$  cities. Now consider an arbitrary  $n + 1$ -city country. Hide one city and use your assumption.
  2. Prove that there is a city<sup>3</sup> to which you can drive from all the others.
- 25.** Let  $S \subseteq \mathbb{R}$ . We say that  $S$  is **well-ordered** iff any non-empty subset  $X \subset S$  has the smallest element.
1. Prove that reals and integers are not well-ordered.
  2. Assume that  $X \subseteq \mathbb{N}$  doesn't have the smallest element. Define  $A = \{n \in \mathbb{N} : \{0, 1, \dots, n\} \cap X = \emptyset\}$  and use mathematical induction to prove that  $X$  is empty.
  3. Why are natural numbers well-ordered?

## 1.4 Functions

### 1.4.1 Basics

- 26.** (Thanks to Antoni Hanke) How many are there functions from the empty set to  $\{1, 2, 3, 4\}$ ?
- 27.** Consider two functions:  $f : \{0, 1\} \rightarrow \{0, 1\}$  given by  $f(x) = 0$  and  $g : \{0, 1\} \rightarrow \{0\}$ . Prove that  $f = g$ .<sup>4</sup>
- 28.** Let  $f : A \rightarrow B$  and  $g : C \rightarrow B$ , where  $A \neq C$ . Is it possible that  $f = g$ ?
- 29.** Let  $f : A \rightarrow B$  and  $C \subseteq D \subseteq A$ . We define:  $f[C] = \{b \in B : b = f(c) \text{ for some } c \in C\}$  and analogously  $f[D]$ . Prove that  $f(C) \subseteq f(D)$ .

### 1.4.2 Injectivity, surjectivity and bijectivity

- 30.** As we remember,  $\mathbb{R}$  stands for well-known real numbers. Are the following functions surjective?
1.  $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^3$
  2.  $g : \mathbb{R} \rightarrow \mathbb{R}, g(x) = x^2$
  3.  $h : \mathbb{R} \rightarrow \{5\}$
- 31.** As we remember,  $\mathbb{R}$  stands for well-known real numbers. Are the following functions injective?
1.  $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$
  2.  $h : \{0, 1, 2, 3\} \rightarrow \mathbb{R}, h(x) = x$

---

<sup>3</sup>Nice trick: what does happen if you reverse each way? Can you use the former result?

<sup>4</sup>Some mathematicians, as Bourbaki use an alternative definition of function - for them a function is the triple  $(A, B, f)$ , where  $f$  is defined as in the our case. We see that this definition is incompatible with ours. Fortunately, as in the case with different definitions of ordered tuples, this problem will never occur explicitly in the further chapters.

**32.** Construct function that is:

1. surjective, but not injective
2. injective, but not surjective
3. neither injective nor surjective
4. bijective

**33.** Prove that, if exists,  $g$  is unique.

**34.** Assume that  $f^{-1}$  exists. Prove that  $(f^{-1})^{-1}$  exists and is equal to  $f$ .

### 1.4.3 Function composition

**35.** Find functions  $f, g$  such that:

1.  $g \circ f$  exists, but  $f \circ g$  is not defined
2. both  $f \circ g$  and  $g \circ f$  exist, but  $f \circ g \neq g \circ f$

**36.** Let  $f : A \rightarrow B, g : B \rightarrow C, h : C \rightarrow D$ . Prove that

$$h \circ (g \circ f) = (h \circ g) \circ f.$$

**37.**

1. Prove that composition of two surjections is surjective.
2. Prove that composition of two injections is injective.
3. Prove that composition of two bijections is bijective.

**38.** We will rephrase the definition of the inverse function as follows:

1. If  $X$  is a set, we define **the identity function**

$$\text{Id}_X = \{(x, x) \in X^2 : x \in X\}.$$

Prove that it is indeed a function. What is its domain?

2. Let  $f : A \rightarrow B, g : B \rightarrow A$ . Prove that  $f = g^{-1}$  iff

$$g \circ f = \text{Id}_A \text{ and } f \circ g = \text{Id}_B$$

## 1.5 Countability

### 1.5.1 Finite sets

**39.** What is the cardinality of  $\{a, a+1, a+2, \dots, a+n\}$ ?

**40.** Let  $A$ ,  $B$  and  $C$  be finite sets. Prove that:

1.  $|2^A| = 2^{|A|}$
2.  $|A \cup B| = |A| + |B|$  iff  $A$  and  $B$  are disjoint.
3.  $|A \setminus B| = |A| - |B|$  if  $B \subseteq A$ .
4.  $|A| \geq |B|$  if  $B \subseteq A$ . When does the equality hold?
5.  $|A \cup B| = |A| + |B| - |A \cap B|$
6.  $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |B \cap C| - |C \cap A| + |A \cap B \cap C|$

**41.** Assume that  $A$  and  $B$  are finite sets. Prove that  $|A| = |B|$  iff there is a bijection between  $A$  and  $B$ .

**42.** Above we find the way of saying that two cardinalities are equal using existence of a bijection. Let's find a way to compare which is less using another kind of function.

1. Let  $O_n = \{1, 2, \dots, n\}$ . Prove that there is no injection from  $O_{n+1}$  into  $O_n$ . Hint: use mathematical induction.
2. Let  $A$  and  $B$  be finite. Prove that there is an injection from  $A$  to  $B$  iff  $|A| \leq |B|$ .

**43.** Using the above results, prove in one line<sup>5</sup> that if there is an injection from  $A$  onto  $B$  and an injection from  $B$  into  $A$ , then there exists a bijection from  $A$  onto  $B$ .

### 1.5.2 Infinite sets

**44.** Let  $A$ ,  $B$  and  $C$  be sets. Prove that if  $|A| = |B|$  and  $|B| = |C|$ , then  $|A| = |C|$ . Hint: find the bijection between  $A$  and  $C$ .

**45.** Prove that:

1.  $|\mathbb{N}| = |\mathbb{Z}|$ .
2.  $|\mathbb{N}| = |\mathbb{N} \times \mathbb{N}|$ .
3.  $|\mathbb{N}| = |\mathbb{Q}|$ .

**46.** Prove that if  $A \subseteq B$ , then  $|A| \leq |B|$ .

**47.** Let  $A$ ,  $B$  and  $C$  be sets. Prove that if  $|A| \leq |B|$  and  $|B| \leq |C|$ , then  $|A| \leq |C|$ .

---

<sup>5</sup>The main step is  $|A| \leq B$  and  $|B| \leq |A|$ , so  $|A| = |B|$ .

**48.** Here you can prove that there are more real numbers than naturals or rationals. We define  $X = \{x \in \mathbb{R} : 0 \leq x \leq 1\}$  and choose one convention of writing reals (e.g.  $0.999\dots = 1.000\dots$ , so we can choose to use nines)

1. Assume that you have written all the elements of  $X$  in a single column. Can you find a real number that does not occur in the list?
2. Using the above, prove that  $|\mathbb{N}| < |X|$
3. Prove that  $|\mathbb{Q}| < |\mathbb{R}|$ .

**49.** We know that  $|\mathbb{R}| > |\mathbb{N}|$ . Using binary system prove that  $\mathbb{R} = 2^{\mathbb{N}}$ . Do you see similarity between the previous result and  $2^n > n$  for natural  $n$ ?

**50. Cantor's theorem** You will prove that  $|A| < |2^A|$  for any set  $A$ . Let  $A$  be a set and  $f : A \rightarrow 2^A$ .

1. Consider  $X = \{a \in A : a \notin f(a)\} \in 2^A$ . Is there  $x \in A$  for which  $f(x) = X$ ?
2. Is  $f$  surjective?
3. Find an injective function  $g : A \rightarrow 2^A$ .
4. Prove that  $|A| < |2^A|$  for any set  $A$ .
5. Use Cantor's theorem to prove that there is no set of all sets.

**51. Cantor-Schroeder-Bernstein theorem** Let's prove that if  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$  for any sets.

1. (Knaster-Tarski) Now assume that  $F$  has *monotonicity* property:  $F(X) \subseteq F(Y)$  if  $X \subseteq Y$ . Prove that  $F(S) = S$ , where:

$$S = \bigcup_{X \in U} X, \text{ where } U = \{Y \in 2^A : Y \subseteq f(Y)\}.$$

2. (Banach) Let  $f : A \rightarrow B$  and  $g : B \rightarrow A$  be injections. We introduce new symbol:  $f[X] = \{b \in B : b = f(x) \text{ for some } x \in X\}$ . Prove that function

$$F : 2^A \rightarrow 2^A, F(X) = A - g[B - f[X]]$$

has the monotonicity property.

3. Using the above statements we know that there is  $S$  defined above in the our case. Prove that function

$$h(x) = \begin{cases} f(x), & x \in S \\ g^{-1}(x), & x \notin S \end{cases}$$

is a bijection. You will need to prove that  $X \setminus S \subseteq \text{Im } g$ .

### 1.5.3 Pre-image of a function

## Chapter 2

# General topology

### 2.1 Basics

#### 2.1.1 Topology and open sets

- 52. Using mathematical induction prove that the intersection of finitely many open sets is open.
- 53. **Trivial topology** Prove that for any  $X$ , set  $\{\emptyset, X\}$  is a topology.
- 54. **Discrete topology** Prove that for any  $X$ , it's power set  $2^X$  is a topology.

#### 2.1.2 Definition of intervals