

## Bivariate Probability Distribution

Example - Two discrete rv's  $X$  and  $Y$

Bivariate pdf					
		Y			
		%	0	1	Pr(X)
X	0		1/8	0	1/8
	1		2/8	1/8	3/8
	2		1/8	2/8	3/8
	3		0	1/8	1/8
		Pr(Y)	4/8	4/8	1

$p(x, y) = \Pr(X = x, Y = y) = \text{values in table}$   
*e.g.*,  $p(0, 0) = \Pr(X = 0, Y = 0) = 1/8$

Properties of joint pdf  $p(x, y)$

$$S_{XY} = \{(0, 0), (0, 1), (1, 0), (1, 1), (2, 0), (2, 1), (3, 0), (3, 1)\}$$

$$p(x, y) \geq 0 \text{ for } x, y \in S_{XY}$$

$$\sum_{x, y \in S_{XY}} p(x, y) = 1$$

Marginal pdfs

$$p(x) = \Pr(X = x) = \sum_{y \in S_Y} p(x, y)$$

= sum over columns in joint table

$$p(y) = \Pr(Y = y) = \sum_{x \in S_X} p(x, y)$$

= sum over rows in joint table

## Conditional Probability

Suppose we know  $Y = 0$ . How does this knowledge affect the probability that  $X = 0, 1, 2$ , or  $3$ ? The answer involves conditional probability.

Example

$$\begin{aligned} \Pr(X = 0|Y = 0) &= \frac{\Pr(X = 0, Y = 0)}{\Pr(Y = 0)} \\ &= \frac{\text{joint probability}}{\text{marginal probability}} = \frac{1/8}{4/8} = 1/4 \end{aligned}$$

Remark

$$\begin{aligned} \Pr(X = 0|Y = 0) &= 1/4 \neq \Pr(X = 0) = 1/8 \\ \implies X &\text{ depends on } Y \end{aligned}$$

The marginal probability,  $\Pr(X = 0)$ , ignores information about  $Y$ .

### Definition - Conditional Probability

- The conditional pdf of  $X$  given  $Y = y$  is, for all  $x \in S_X$ ,

$$p(x|y) = \Pr(X = x|Y = y) = \frac{\Pr(X = x, Y = y)}{\Pr(Y = y)}$$

- The conditional pdf of  $Y$  given  $X = x$  is, for all values of  $y \in S_Y$

$$p(y|x) = \Pr(Y = y|X = x) = \frac{\Pr(X = x, Y = y)}{\Pr(X = x)}$$

### Conditional Mean and Variance

$$\begin{aligned}\mu_{X|Y=y} &= E[X|Y = y] = \sum_{x \in S_X} x \cdot \Pr(X = x|Y = y), \\ \mu_{Y|X=x} &= E[Y|X = x] = \sum_{y \in S_Y} y \cdot \Pr(Y = y|X = x).\end{aligned}$$

$$\begin{aligned}\sigma_{X|Y=y}^2 &= \text{var}(X|Y = y) = \sum_{x \in S_X} (x - \mu_{X|Y=y})^2 \cdot \Pr(X = x|Y = y), \\ \sigma_{Y|X=x}^2 &= \text{var}(Y|X = x) = \sum_{y \in S_Y} (y - \mu_{Y|X=x})^2 \cdot \Pr(Y = y|X = x).\end{aligned}$$

Example:

$$\begin{aligned}E[X] &= 0 \cdot 1/8 + 1 \cdot 3/8 + 2 \cdot 3/8 + 3 \cdot 1/8 = 3/2 \\E[X|Y = 0] &= 0 \cdot 1/4 + 1 \cdot 1/2 + 2 \cdot 1/4 + 3 \cdot 0 = 1, \\E[X|Y = 1] &= 0 \cdot 0 + 1 \cdot 1/4 + 2 \cdot 1/2 + 3 \cdot 1/4 = 2,\end{aligned}$$

$$\begin{aligned}\text{var}(X) &= \text{var}(X|Y = 0) = (0 - 3/2)^2 \cdot 1/8 + (1 - 3/2)^2 \cdot 3/8 \\&\quad + (2 - 3/2)^2 \cdot 3/8 + (3 - 3/2)^2 \cdot 1/8 = 3/4, \\ \text{var}(X|Y = 0) &= (0 - 1)^2 \cdot 1/4 + (1 - 1)^2 \cdot 1/2 \\&\quad + (2 - 1)^2 \cdot 1/2 + (3 - 1)^2 \cdot 0 = 1/2, \\ \text{var}(X|Y = 1) &= (0 - 2)^2 \cdot 0 + (1 - 2)^2 \cdot 1/4 \\&\quad + (2 - 2)^2 \cdot 1/2 + (3 - 2)^2 \cdot 1/4 = 1/2.\end{aligned}$$

## Independence

Let  $X$  and  $Y$  be discrete rvs with pdfs  $p(x)$ ,  $p(y)$ , sample spaces  $S_X, S_Y$  and joint pdf  $p(x, y)$ . Then  $X$  and  $Y$  are independent rv's if and only if

$$\begin{aligned}p(x, y) &= p(x) \cdot p(y) \\&\text{for all values of } x \in S_X \text{ and } y \in S_Y\end{aligned}$$

Result: If  $X$  and  $Y$  are independent rv's, then

$$\begin{aligned}p(x|y) &= p(x) \text{ for all } x \in S_X, y \in S_Y \\p(y|x) &= p(y) \text{ for all } x \in S_X, y \in S_Y\end{aligned}$$

Intuition

Knowledge of  $X$  does not influence probabilities associated with  $Y$

Knowledge of  $Y$  does not influence probabilities associated with  $X$

## Bivariate Distributions - Continuous rv's

The joint pdf of  $X$  and  $Y$  is a non-negative function  $f(x, y)$  such that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$$

Let  $[x_1, x_2]$  and  $[y_1, y_2]$  be intervals on the real line. Then

$$\begin{aligned} \Pr(x_1 \leq X \leq x_2, y_1 \leq Y \leq y_2) \\ &= \int_{x_1}^{x_2} \int_{y_1}^{y_2} f(x, y) dx dy \\ &= \text{volume under probability surface} \\ &\quad \text{over the intersection of the intervals} \\ &\quad [x_1, x_2] \text{ and } [y_1, y_2] \end{aligned}$$

## Marginal and Conditional Distributions

The marginal pdf of  $X$  is found by integrating  $y$  out of the joint pdf  $f(x, y)$  and the marginal pdf of  $Y$  is found by integrating  $x$  out of the joint pdf:

$$\begin{aligned} f(x) &= \int_{-\infty}^{\infty} f(x, y) dy, \\ f(y) &= \int_{-\infty}^{\infty} f(x, y) dx. \end{aligned}$$

The conditional pdf of  $X$  given that  $Y = y$ , denoted  $f(x|y)$ , is computed as

$$f(x|y) = \frac{f(x, y)}{f(y)},$$

and the conditional pdf of  $Y$  given that  $X = x$  is computed as

$$f(y|x) = \frac{f(x, y)}{f(x)}.$$

The conditional means are computed as

$$\mu_{X|Y=y} = E[X|Y = y] = \int x \cdot p(x|y)dx,$$
$$\mu_{Y|X=x} = E[Y|X = x] = \int y \cdot p(y|x)dy$$

and the conditional variances are computed as

$$\sigma_{X|Y=y}^2 = \text{var}(X|Y = y) = \int (x - \mu_{X|Y=y})^2 p(x|y)dx,$$
$$\sigma_{Y|X=x}^2 = \text{var}(Y|X = x) = \int (y - \mu_{Y|X=x})^2 p(y|x)dy.$$

### **Independence.**

Let  $X$  and  $Y$  be continuous random variables.  $X$  and  $Y$  are independent iff

$$f(x|y) = f(x), \text{ for } -\infty < x, y < \infty,$$
$$f(y|x) = f(y), \text{ for } -\infty < x, y < \infty.$$

Result: Let  $X$  and  $Y$  be continuous random variables .  $X$  and  $Y$  are independent iff

$$f(x, y) = f(x)f(y)$$

The result in the above proposition is extremely useful in practice because it gives us an easy way to compute the joint pdf for two independent random variables: we simply compute the product of the marginal distributions.

Example: Bivariate standard normal distribution

Let  $X \sim N(0, 1)$ ,  $Y \sim N(0, 1)$  and let  $X$  and  $Y$  be independent. Then

$$\begin{aligned} f(x, y) &= f(x)f(y) = \frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}x^2} \frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}y^2} \\ &= \frac{1}{2\pi}e^{-\frac{1}{2}(x^2+y^2)}. \end{aligned}$$

To find  $\Pr(-1 < X < 1, -1 < Y < 1)$  we must solve

$$\int_{-1}^1 \int_{-1}^1 \frac{1}{2\pi} e^{-\frac{1}{2}(x^2+y^2)} dx dy$$

which, unfortunately, does not have an analytical solution. Numerical approximation methods are required to evaluate the above integral. See R package mvtnorm.

## Covariance and Correlation - Measuring linear dependence between two rv's

Covariance: Measures direction but not strength of linear relationship between 2 rv's

$$\begin{aligned} \sigma_{XY} &= E[(X - \mu_X)(Y - \mu_Y)] \\ &= \sum_{x,y \in S_{XY}} (x - \mu_X)(y - \mu_Y) \cdot p(x, y) \quad (\text{discrete}) \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - \mu_X)(y - \mu_Y) f(x, y) dx dy \quad (\text{cts}) \end{aligned}$$

Example: For the data in Table 2, we have

$$\begin{aligned}\sigma_{XY} = \text{Cov}(X, Y) &= (0 - 3/2)(0 - 1/2) \cdot 1/8 \\ &\quad + (0 - 3/2)(1 - 1/2) \cdot 0 + \dots \\ &\quad + (3 - 3/2)(1 - 1/2) \cdot 1/8 = 1/4\end{aligned}$$

### Properties of Covariance

$$\text{Cov}(X, Y) = \text{Cov}(Y, X)$$

$$\text{Cov}(aX, bY) = a \cdot b \cdot \text{Cov}(X, Y) = a \cdot b \cdot \sigma_{XY}$$

$$\text{Cov}(X, X) = \text{Var}(X)$$

$$X, Y \text{ independent} \implies \text{Cov}(X, Y) = 0$$

$$\text{Cov}(X, Y) = 0 \not\Rightarrow X \text{ and } Y \text{ are independent}$$

$$\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$$



**Correlation:** Measures direction and strength of linear relationship between 2 rv's

$$\begin{aligned}\rho_{XY} &= \text{Cor}(X, Y) = \frac{\text{Cov}(X, Y)}{\text{SD}(X) \cdot \text{SD}(Y)} \\ &= \frac{\sigma_{XY}}{\sigma_X \cdot \sigma_Y} = \text{scaled covariance}\end{aligned}$$

Example: For the Data in Table 2

$$\rho_{XY} = \text{Cor}(X, Y) = \frac{1/4}{\sqrt{(3/4) \cdot (1/2)}} = 0.577$$

## Properties of Correlation

$$-1 \leq \rho_{XY} \leq 1$$

$$\rho_{XY} = 1 \text{ if } Y = aX + b \text{ and } a > 0$$

$$\rho_{XY} = -1 \text{ if } Y = aX + b \text{ and } a < 0$$

$$\rho_{XY} = 0 \text{ if and only if } \sigma_{XY} = 0$$

$$\rho_{XY} = 0 \not\Rightarrow X \text{ and } Y \text{ are independent in general}$$

$$\rho_{XY} = 0 \implies \text{independence if } X \text{ and } Y \text{ are normal}$$

## Bivariate normal distribution

Let  $X$  and  $Y$  be distributed bivariate normal. The joint pdf is given by

$$f(x, y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \times \exp \left\{ -\frac{1}{2(1-\rho^2)} \left[ \left( \frac{x-\mu_X}{\sigma_X} \right)^2 + \left( \frac{y-\mu_Y}{\sigma_Y} \right)^2 - \frac{2\rho(x-\mu_X)(y-\mu_Y)}{\sigma_X\sigma_Y} \right] \right\}$$

where  $E[X] = \mu_X$ ,  $E[Y] = \mu_Y$ ,  $SD(X) = \sigma_X$ ,  $SD(Y) = \sigma_Y$ , and  $\rho = \text{cor}(X, Y)$ .

## Linear Combination of 2 rv's

Let  $X$  and  $Y$  be rv's. Define a new rv  $Z$  that is a linear combination of  $X$  and  $Y$  :

$$Z = aX + bY$$

where  $a$  and  $b$  are constants. Then

$$\begin{aligned}\mu_Z &= E[Z] = E[aX + bY] \\ &= aE[X] + bE[Y] \\ &= a \cdot \mu_X + b \cdot \mu_Y\end{aligned}$$

and

$$\begin{aligned}\sigma_Z^2 &= \text{Var}(Z) = \text{Var}(a \cdot X + b \cdot Y) \\ &= a^2 \text{Var}(X) + b^2 \text{Var}(Y) + 2a \cdot b \cdot \text{Cov}(X, Y) \\ &= a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2a \cdot b \cdot \sigma_{XY}\end{aligned}$$

If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  then  $Z \sim N(\mu_Z, \sigma_Z^2)$

Example: Portfolio returns

$R_A$  = return on asset  $A$  with  $E[R_A] = \mu_A$  and  $\text{Var}(R_A) = \sigma_A^2$

$R_B$  = return on asset  $B$  with  $E[R_B] = \mu_B$  and  $\text{Var}(R_B) = \sigma_B^2$

$\text{Cov}(R_A, R_B) = \sigma_{AB}$  and  $\text{Cor}(R_A, R_B) = \rho_{AB} = \frac{\sigma_{AB}}{\sigma_A \sigma_B}$

Portfolio

$x_A$  = share of wealth invested in asset  $A$ ,  $x_B$  = share of wealth invested in asset  $B$

$x_A + x_B = 1$  (exhaust all wealth in 2 assets)

$R_P = x_A \cdot R_A + x_B \cdot R_B$  = portfolio return

Portfolio Problem: How much wealth should be invested in assets  $A$  and  $B$ ?

Portfolio expected return (gain from investing)

$$\begin{aligned}E[R_P] &= \mu_P = E[x_A \cdot R_A + x_B \cdot R_B] \\&= x_A E[R_A] + x_B E[R_B] \\&= x_A \mu_A + x_B \mu_B\end{aligned}$$

Portfolio variance (risk from investing)

$$\begin{aligned}\text{Var}(R_P) &= \sigma_P^2 = \text{Var}(x_A R_A + x_B R_B) \\&= x_A^2 \text{Var}(R_A) + x_B^2 \text{Var}(R_B) + \\&\quad 2 \cdot x_A \cdot x_B \cdot \text{Cov}(R_A, R_B) \\&= x_A^2 \sigma_A^2 + x_B^2 \sigma_B^2 + 2x_A x_B \sigma_{AB} \\SD(R_P) &= \sqrt{\text{Var}(R_P)} = \sigma_P \\&= \left( x_A^2 \sigma_A^2 + x_B^2 \sigma_B^2 + 2x_A x_B \sigma_{AB} \right)^{1/2}\end{aligned}$$

Linear Combination of  $N$  rv's.

Let  $X_1, X_2, \dots, X_N$  be rvs and let  $a_1, a_2, \dots, a_N$  be constants. Define

$$Z = a_1 X_1 + a_2 X_2 + \dots + a_N X_N = \sum_{i=1}^N a_i X_i$$

Then

$$\begin{aligned}\mu_Z &= E[Z] = a_1 E[X_1] + a_2 E[X_2] + \dots + a_N E[X_N] \\&= \sum_{i=1}^N a_i E[X_i] = \sum_{i=1}^N a_i \mu_i\end{aligned}$$

For the variance,

$$\begin{aligned}\sigma_Z^2 = \text{Var}(Z) &= a_1^2 \text{Var}(X_1) + \cdots + a_N^2 \text{Var}(X_N) \\ &+ 2a_1a_2 \text{Cov}(X_1, X_2) + 2a_1a_3 \text{Cov}(X_1, X_3) + \cdots \\ &+ 2a_2a_3 \text{Cov}(X_2, X_3) + 2a_2a_4 \text{Cov}(X_2, X_4) + \cdots \\ &+ 2a_{N-1}a_N \text{Cov}(X_{N-1}, X_N)\end{aligned}$$

Note:  $N$  variance terms and  $N(N - 1) = N^2 - N$  covariance terms. If  $N = 100$ , there are  $100 \times 99 = 9900$  covariance terms!

Result: If  $X_1, X_2, \dots, X_N$  are each normally distributed random variables then

$$Z = \sum_{i=1}^N a_i X_i \sim N(\mu_Z, \sigma_Z^2)$$

Example: Portfolio variance with three assets

$R_A, R_B, R_C$  are simple returns on assets A, B and C

$x_A, x_B, x_C$  are portfolio shares such that  $x_A + x_B + x_C = 1$

$$R_p = x_A R_A + x_B R_B + x_C R_C$$

Portfolio variance

$$\begin{aligned}\sigma_P^2 &= x_A^2 \sigma_A^2 + x_B^2 \sigma_B^2 + x_C^2 \sigma_C^2 \\ &+ 2x_A x_B \sigma_{AB} + 2x_A x_C \sigma_{AC} + 2x_B x_C \sigma_{BC}\end{aligned}$$

Note: Portfolio variance calculation may be simplified using matrix layout

$$\begin{array}{cccc}
 & x_A & x_B & x_C \\
 x_A & \sigma_A^2 & \sigma_{AB} & \sigma_{AC} \\
 x_B & \sigma_{AB} & \sigma_B^2 & \sigma_{BC} \\
 x_C & \sigma_{AC} & \sigma_{BC} & \sigma_C^2
 \end{array}$$

Example: Multi-period continuously compounded returns

$$r_t = \ln(1 + R_t) = \text{monthly cc return}$$

$$r_t \sim N(\mu, \sigma^2) \quad \text{for all } t$$

$$\text{Cov}(r_t, r_s) = 0 \quad \text{for all } t \neq s$$

Annual return

$$\begin{aligned}
 r_t(12) &= \sum_{j=0}^{11} r_{t-j} \\
 &= r_t + r_{t-1} + \cdots + r_{t-11}
 \end{aligned}$$

Then

$$\begin{aligned} E[r_t(12)] &= \sum_{j=0}^{11} E[r_{t-j}] \\ &= \sum_{j=0}^{11} \mu \quad (E[r_t] = \mu \text{ for all } t) \\ &= 12\mu \quad (\mu = \text{mean of monthly return}) \end{aligned}$$

$$\begin{aligned} \text{Var}(r_t(12)) &= \text{Var}\left(\sum_{j=0}^{11} r_{t-j}\right) \\ &= \sum_{j=0}^{11} \text{Var}(r_{t-j}) = \sum_{j=0}^{11} \sigma^2 \\ &= 12 \cdot \sigma^2 \quad (\sigma^2 = \text{monthly variance}) \\ \text{SD}(r_t(12)) &= \sqrt{12} \cdot \sigma \quad (\text{square root of time rule}) \end{aligned}$$

Then

$$r_t(12) \sim N(12\mu, 12\sigma^2)$$

For example, suppose

$$r_t \sim N(0.01, (0.10)^2)$$

Then

$$E[r_t(12)] = 12 \times (0.01) = 0.12$$

$$\text{Var}(r_t(12)) = 12 \times (0.10)^2 = 0.12$$

$$\text{SD}(r_t(12)) = \sqrt{0.12} = 0.346$$

$$r_t(12) \sim N(0.12, (0.346)^2)$$