

Exercise : Consider these three points: $\{(1, 1), (2.5, 8), (4, 5)\}$. Find the polynomial $P(x)$ of degree 2 which passes through these points. Do this three different ways, by using

(a) the Vandermonde matrix method,

```

1 >> x=[1,2.5,8]';
2 >> y=[1,8,5]';
3 >> v=vander(x)
4 v =
5
6      1.0000      1.0000      1.0000
7      6.2500      2.5000      1.0000
8     64.0000      8.0000      1.0000
9
10 >> c=v\y
11 c =
12
13     -0.74459
14      7.27273
15     -5.52814
16
17 >> P=@(x) c(1)*x.^2+c(2)*x+c(3)
18 P =
19
20 @ (x) c (1) * x .^ 2 + c (2) * x + c (3)
21
22 >> P(x)
23 ans =
24
25      1.0000
26      8.0000
27      5.0000
28
29 >> y
30 y =
31
32      1
33      8
34      5
35
36 >> diary off

```

(b) the Newton form and its triangular matrix method

```

1 >> A=[1,0,0;1,x(2)-x(1),0;1,x(3)-x(1),(x(3)-x(1))*(x(3)-x(2))]
2 A =
3
4      1.00000      0.00000      0.00000
5      1.00000      1.50000      0.00000

```

```

6      1.00000    7.00000    38.50000
7
8  >> a=A\y
9  a =
10
11      1.00000
12      4.66667
13     -0.74459
14
15  >> P=@(x) a(1)+a(2)*(x-x(1))+a(3).*(x-x(1)).*(x-x(2))
16  P =
17
18  @(x) a(1) + a(2) * (x - x(1)) + a(3) .* (x - x(1)) .* (x - x(2))
19
20  >> P(x)
21  ans =
22
23      1.0000
24      8.0000
25      5.0000
26
27  >> y
28  y =
29
30      1
31      8
32      5
33
34  >> diary off

```

In standard form:

```

1  >> c=[a(3),-x(1)*a(3)-x(2)*a(3)+a(2),x(1)*x(2)*a(3)-x(1)*a(2)+a(1)]
2  c =
3
4     -0.74459     7.27273    -5.52814
5
6  >> diary off

```

These are exactly the same coefficients as part a.

(c) the Lagrange form

```

1  >> t1=@(X) y(1)*((X-x(2)).*(X-x(3)))./((x(1)-x(2)).*(x(1)-x(3)))
2  t1 =
3
4  @(X) y(1) * ((X - x(2)) .* (X - x(3))) ./ ((x(1) - x(2)) .* (x(1) - x(3)))
5
6  >> t2=@(X) y(2)*((X-x(1)).*(X-x(3)))./((x(2)-x(1)).*(x(2)-x(3)))
7  t2 =

```

```

8
9 @ (X) y (2) * ((X - x (1)) .* (X - x (3))) ./ ((x (2) - x (1)) .* (x (2) - x (3)))
10
11 >> t3=@ (X) y (3) * ((X-x (1)) .* (X-x (2))) ./ ((x (3)-x (1)) .* (x (3)-x (2)))
12 t3 =
13
14 @ (X) y (3) * ((X - x (1)) .* (X - x (2))) ./ ((x (3) - x (1)) .* (x (3) - x (2)))
15
16 >> P=@ (X) t1 (X) +t2 (X) +t3 (X)
17 P =
18
19 @ (X) t1 (X) + t2 (X) + t3 (X)
20
21 >> P (x)
22 ans =
23
24     1
25     8
26     5
27
28 >> diary off

```

In standard form:

```

1 >> c (1)=y (1) / ((x (1)-x (2)) .* (x (1)-x (3)))
2 c =
3
4     0.095238     0.000000     0.000000
5
6 >> c (1) += y (2) / ((x (2)-x (1)) .* (x (2)-x (3)))
7 c =
8
9    -0.87446     0.00000     0.00000
10
11 >> c (1) += y (3) / ((x (3)-x (1)) .* (x (3)-x (2)))
12 c =
13
14    -0.74459     0.00000     0.00000
15
16 >> diary off
17 >> c (2)=y (1) * (-x (2)-x (3)) ./ ((x (1)-x (2)) .* (x (1)-x (3)))
18 c =
19
20    -0.74459    -1.00000     0.00000
21
22 >> c (2) += y (2) * (-x (1)-x (3)) ./ ((x (2)-x (1)) .* (x (2)-x (3)))
23 c =
24
25    -0.74459     7.72727     0.00000
26
27 >> c (2) += y (3) * (-x (1)-x (2)) ./ ((x (3)-x (1)) .* (x (3)-x (2)))

```

```

28 c =
29
30 -0.74459    7.27273    0.00000
31
32 >> diary off
33 >> c(3)=y(1)*(x(2)*x(3))./( (x(1)-x(2)).*(x(1)-x(3)) )
34 c =
35
36 -0.74459    7.27273    1.90476
37
38 >> c(3) += y(2)*(x(1)*x(3))./( (x(2)-x(1)).*(x(2)-x(3)) )
39 c =
40
41 -0.74459    7.27273   -5.85281
42
43 >> c(3) += y(3)*(x(1)*x(2))./( (x(3)-x(1)).*(x(3)-x(2)) )
44 c =
45
46 -0.74459    7.27273   -5.52814
47
48 >> diary off

```

These are exactly the same coefficients as part a.

(d) True as shown above.


Exercise : 8.1

(a) The estimated value at 120 is 459.60. I feel like this estimate is way too high, a better estimate would be around 300, following a best fit line, as the data appears to be linear.

```

1 year=[0:20:100]';
2 pop=[76.0,105.7,131.7,179.3,226.5,281.4]';
3 plot(year,pop,'o');
4 v=vander(year);
5 c=v\pop;
6 x=[0:.5:120];
7 hold on;
8 plot(x,polyval(c,x));
9 polyval(c,120)
10 hold off;

```



- (b) the newton form constants are independent of the number of additional points, so the zero order constant is the first constant of the first order and the first constant of the second order. Thus I can simply give the second order constants and the first two are the first order constants and the first one is the zero order constant.

```

1  x=[0:20:40]';
2  y=[76.0,105.7,131.7]';
3  t1=@(X) y(1)*((X-x(2)).*(X-x(3)))/((x(1)-x(2)).*(x(1)-x(3)));
4  t2=@(X) y(2)*((X-x(1)).*(X-x(3)))/((x(2)-x(1)).*(x(2)-x(3)));
5  t3=@(X) y(3)*((X-x(1)).*(X-x(2)))/((x(3)-x(1)).*(x(3)-x(2)));
6  L=@(X) t1(X)+t2(X)+t3(X);
7  #this is the Lagrange form
8
9  A=[1,0,0;1,x(2)-x(1),0;1,x(3)-x(1),(x(3)-x(1)).*(x(3)-x(2))];
10 a=A\y
11 N=@(X) a(1)+a(2)*(X-x(1))+a(3).*(X-x(1)).*(X-x(2));
12 #this is the newton form
13

```

```

14 #to quickly verifi that they are the same
15 r=(rand(20000,1)-.5)*400;
16 l=L(r);
17 n=N(r);
18 dif=abs(l-n);
19 errorval=sum(dif)

```

```

1 >> q81b
2 a =
3
4     7.6000e+01
5     1.4850e+00
6    -4.6250e-03
7
8 errorval =     6.0952e-09
9 >> diary off

```

Noting that the sum of the absolute error is extremely small, order 10^{-9} , for 20000 points we can say that the two are identical, within machine error.

Exercise : 8.2

The Lagrange form of the third order polynomial would be

$$P(x) = \sum_{i=k-3}^k f(i) * \prod_{j \in [k-3, k]; j \neq i} \frac{x - x_j}{x_i - x_j}$$

```

1 >> f=@(x) x.^3-2;
2 >> x=[0,1,2]';
3 >> y=f(x);
4 >> v=vander(x);
5 >> c=v\y;
6 >> si=[-1,1];
7 >> tx=(-c(2)+si*sqrt(c(2)^2-4*c(1)*c(3)))/(2*c(1))
8 tx =
9
10    -0.54858    1.21525
11
12 >> diary off

```

since 1.21525 is closer to 2, it is our next x value.

Exercise : 8.7

```

1 R=@(x) 1./(x.^2+1);
2 px=[-4:.01:4]';

```

```

3 plot(px,R(px));
4 x=linspace(-5,5,13);
5 fx=R(x);
6 y=interp1(x,fx,px);
7 hold on;
8 plot(px,y,'r-');
9 h=12/10
10 dxR=@(x) -2*x./(x.^2+1).^2;
11 fpx=dxR(x);
12 for i=[1:length(px)]
13     y(i)=herm(x,fx,fpx,px(i));
14 end
15 plot(px,y,'g-');
16 hold off;
17 figure(2);
18 plot(px,abs(y-R(px)));


```


```

1 function yout=herm(x,y,yp,xeval)
2 i=2;
3 while(true)
4     if(x(i-1)<=xeval && x(i)>xeval)
5         break;
6     end
7     i+=1;
8 end
9 h=x(i)-x(i-1);
10 alph=3/h^2*(yp(i-1)+yp(i))+6/h^3*(y(i-1)-y(i));
11 yout=-yp(i-1)/h*((xeval-x(i))^2/2-h^2/2)+yp(i)/h*(xeval-x(i-1))^2/2;
12 yout+=alph*(xeval-x(i-1))^2*((xeval-x(i-1))/3-h/2)+y(i-1);
13 endfunction

```

I will present my results as a graph, however the results for the Hermite polynomial are so close that one would need a zoom to make out the difference. Thus I also give a difference plot between the Hermite polynomial and the function.



**Exercise : 8.8**

```
1 x=linspace(0,pi,1000);  
2 y=sin(x);
```

```

3 r=rand(100,1)*pi;
4 ry=sin(r);
5 Rest=r;
6 for i=[1:length(r)]
7     Rest(i)=rest(x,y,r(i));
8 end
9 abserr=max(abs(Rest-ry))
10 relerr=max(abs((Rest-ry)./ry))

```

```

1 function yest=rest(x,y,r)
2 i=2;
3 while(true)
4     if(x(i-1)<=r && r<x(i))
5         break;
6     end
7     i+=1;
8 end
9 m=(y(i-1)-y(i))/(x(i-1)-x(i));
10 yest=m*(r-x(i))+y(i);
11 endfunction

```

```

1 >> q88
2 abserr = 1.2278e-06
3 relerr = 1.2361e-06
4 >> q88
5 abserr = 1.2091e-06
6 relerr = 1.2326e-06
7 >> q88
8 abserr = 1.2149e-06
9 relerr = 1.2361e-06
10 >> q88
11 abserr = 1.2095e-06
12 relerr = 1.2361e-06
13 >> q88
14 abserr = 1.2048e-06
15 relerr = 1.2362e-06
16 >> diary off

```

Exercise : 7.14

```

1 x=linspace(0,1,50)';
2 y=cos(4*x);
3 v=vander(x',10);
4 #note we want v*c=y, however this is imposible, as c does not have enough variables
5 c=v\y;
6 c2=(v'*v)\(v'*y);
7 format long e

```

```
8  c
9  c2
10 err=abs(c-c2)
```

```
1  >> q714
2  c =
3
4  -5.95484314049769e-01
5    2.07303158488186e+00
6    8.48610575052772e-03
7  -5.93544628891181e+00
8    1.82203937403852e-01
9    1.06028268880602e+01
10   1.17638892210016e-02
11  -8.00106220766967e+00
12   3.67105073956209e-05
13   9.99999815646296e-01
14
15  c2 =
16
17  -5.95531446839227e-01
18    2.07324609938819e+00
19    8.07783930992307e-03
20  -5.93502522532791e+00
21    1.81949130193354e-01
22    1.06029183399554e+01
23    1.17451885163276e-02
24  -8.00106024972344e+00
25    3.66287854458158e-05
26    9.99999816219448e-01
27
28  err =
29
30    4.71327894575602e-05
31    2.14514506335473e-04
32    4.08266440604644e-04
33    4.21063583897485e-04
34    2.54807210497449e-04
35    9.14518952388477e-05
36    1.87007046739979e-05
37    1.95794623181200e-06
38    8.17219498050814e-08
39    5.73151748284317e-10
40
41  >> diary off
```

The errors for the low order constants have more error than the high order ones.