



APPLICATION GUIDELINES FOR ARABLEONE™

Introduction



ArableONE™ is a revolutionary soil conditioning formula designed to transform your soil into a thriving, productive growing environment. This unique product works by enhancing soil biology, improving nutrient availability, and creating optimal conditions for healthy plant growth.

What makes ArableONE™ special? Unlike traditional fertilizers that simply add nutrients, ArableONE™ works with nature to unlock your soil's existing potential. It stimulates beneficial microorganisms, improves soil structure, and helps plants access nutrients more efficiently.



How ArableONE™ Works

ArableONE™ addresses multiple soil challenges simultaneously:

- **Enhances Soil Biology:** Provides nutrients and conditions that beneficial soil microorganisms need to thrive
- **Improves Soil Structure:** Contains flocculating agents that increase soil pore space, improving water penetration and root development
- **Increases Nutrient Availability:** Contains organic complexing agents that make nutrients more available to plants
- **Manages Salt Problems:** Helps buffer harmful salts, especially important in coastal and arid regions
- **Reduces Compaction Effects:** Improves soil aeration and water movement

The Science Behind It: Most plant nutrients come from biological sources in the soil. ArableONE™ enhances this natural process by creating conditions where soil microorganisms can flourish and make nutrients available to plants when they need them.

Key Benefits for Your Farm

- Works on any soil type and with any crop
- Can be applied at any time during the growing season
- Compatible with most fertilizers and agricultural chemicals
- Helps reduce overall fertilizer requirements over time
- Improves crop response and soil health simultaneously

Understanding Crop Categories

Permanent Crops

These are crops that grow on plants lasting many seasons without replanting. Examples include fruit trees, nut trees, and grape vines. These crops require ongoing soil management since the same root systems remain in place for years.

Row Crops (Temporary)

These crops are planted in rows and typically replanted each season. Examples include vegetables like lettuce, cabbage, and carrots. They usually have shorter growing seasons and are harvested within the same year.

Turf and Landscaping

This includes all grass areas such as lawns, sod, sports fields, and ornamental plants like flowers, shrubs, and decorative trees. These areas require consistent care throughout the growing season.

Application Methods and Equipment



Standard Application Rate: 2 Quarts Per Acre

Application Method	Rate Per Acre	Water Volume
Broadcast Spray	2 quarts	Mix with 5-50 gallons water
Band Spray/Injection/Trench	2 quarts	Mix with 5-50 gallons water
Sprinkler Systems	2 quarts	Through irrigation system
Flood/Furrow Irrigation	2 quarts	Direct application
Drip/Micro-sprinkler	2 quarts	Through irrigation system

ArableONE™ can be applied using various methods to fit your existing equipment and farming practices:

Essential Application Guidelines

- Always shake well before using Apply to
- moist soil for best results Do not exceed
- recommended rates
- Apply directly to soil, not plant foliage
- Distribute evenly across treatment area
- Can be tank-mixed with most fertilizers and chemicals
-

Application Schedules by Crop Type

Crop Type	Best Timing	Frequency	Rate per Application	Application Interval
Permanent Crops	Early season	4 times per year	1 quart per application	90 days between applications
Field & Row Crops	Early season	1 time per season	2 quarts per application	Single application
Turf & Grass	Any time	4-6 times per year	1 quart per application	30 days between first two applications

Cereal Crops Application Guide



Crop	Application Rate	Frequency	Irrigation Method	Best Timing
Wheat	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Maize (Corn)	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Rice	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Sorghum	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Barley	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Rye	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Oats	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage
Millet	2 quarts per acre	One application per season	Flood, Rain, Center Pivot	1 week before planting or at foliar stage

Important Notes for Cereal Crops

- Must be applied on moist soil
- Do not exceed suggested amounts per acre
- Apply 7-9 days before planting or shortly after plant breaks ground Disperse evenly
- over entire area

Row Crops (Vegetables) Application Guide

Leafy and Brassica Vegetables

Crop	Application Rate	Frequency	Irrigation Method	Best Timing
Artichokes	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Asparagus	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Cabbage	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Cauliflower	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Broccoli	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Lettuce	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Spinach	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Chicory	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage

Root, Bulb and Tuberous Vegetables

Crop	Application Rate	Frequency	Irrigation Method	Best Timing
Carrots	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Turnips	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Garlic	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Onions	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Cassava	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage

- Must be applied on moist soil
- Do not exceed suggested amounts per acre
- Apply 7-9 days before planting or shortly after plant breaks ground
Disperse evenly over entire area
- Apply directly to soil, do not spray plants



Fruit-Bearing Vegetables Application Guide



Crop	Application Rate	Frequency	Irrigation Method	Best Timing
Cucumbers	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Eggplants	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Tomatoes	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage

Watermelons	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Cantaloupes	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Pumpkins	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage
Squash	2 quarts per acre	One application per season	Flood, Drip, Pivot, Any	1 week before planting or at foliar stage

Fruits and Nuts (Permanent Crops)

Tree Fruits

Crop	Initial Application	Maintenance Applications	Irrigation Method	Best Timing
Avocados	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Bananas	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Dates	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Figs	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Mangoes	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Papayas	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Pineapples	2 quarts per acre	1 quart every 90 days	Any method	Early season is best

Citrus Fruits

Crop	Initial Application	Maintenance Applications	Irrigation Method	Best Timing
Grapefruit	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Lemons	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Limes	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Oranges	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Tangerines	2 quarts per acre	1 quart every 90 days	Any method	Early season is best

Mandarins	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
-----------	-------------------	-----------------------	------------	----------------------

Stone Fruits and Berries

Crop	Initial Application	Maintenance Applications	Irrigation Method	Best Timing
Apples	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Apricots	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Cherries	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Peaches	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Pears	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Plums	2 quarts per acre	1 quart every 90 days	Any method	Early season is best

Crop	Initial Application	Maintenance Applications	Irrigation Method	Best Timing
Grapes	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Strawberries	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Blueberries	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Raspberries	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Blackberries	2 quarts per acre	1 quart every 90 days	Any method	Early season is best

Kiwi	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
------	-------------------	-----------------------	------------	----------------------

Nuts Application Guide

Crop	Initial Application	Maintenance Applications	Irrigation Method	Best Timing
Almonds	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Cashews	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Chestnuts	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Hazelnuts	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Pistachios	2 quarts per acre	1 quart every 90 days	Any method	Early season is best
Walnuts	2 quarts per acre	1 quart every 90 days	Any method	Early season is best

Important Notes for Permanent Crops

- Must be applied on moist soil
- Do not exceed suggested amounts per acre
- Apply early season for best results
Disperse evenly over entire root zone area
Apply directly to soil, do not spray plants
- Maintain consistent 90-day intervals between applications

Key Success Factors

Timing: Early season applications provide the best results

Soil Conditions: Always apply to moist soil for optimal activation

Application Rate: Follow recommended rates - more is not better

Distribution: Ensure even coverage across the treatment area

Soil Contact: Apply directly to soil, not plant foliage

Consistency: Maintain regular application schedules for permanent crops

Remember: ArableONE™ is designed to work with your existing fertilization program, not replace it. Over time, as soil health improves, you may find that you can reduce traditional fertilizer inputs while maintaining or improving crop performance.

For technical support or questions about specific applications, consult with your agricultural advisor or contact our technical support team.

