

1 SEARCH FOR $t\bar{t}Z' \rightarrow t\bar{t}t\bar{t}$ PRODUCTION IN THE MULTILEPTON FINAL STATE IN
2 pp COLLISIONS AT $\sqrt{s} = 13$ TEV WITH THE ATLAS DETECTOR

3 By

4 Hieu Le

5 A DISSERTATION

6 Submitted to
7 Michigan State University
8 in partial fulfillment of the requirements
9 for the degree of

10 Physics — Doctor of Philosophy

11 2025

ABSTRACT

13 Lorem ipsum dolor sit amet, consectetur adipiscing elit, sed do eiusmod tempor incididunt ut
14 labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco
15 laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit in
16 voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat
17 non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

ACKNOWLEDGMENTS

- 19 Advisor: Reinhard Schwienhorst
- 20 Postdoc: Binbin Dong
- 21 Committee
- 22 MSU group
- 23 ATLAS analysis group
- 24 Friend: Daniel, Grayson, Bella, Eric, Jordan
- 25 Other friends: Jasper, Adam, Brittany
- 26 Parents
- 27 Spouse: Allen Sechrist
- 28 ATLAS in general & funding agencies

PREFACE

30 This is my preface. remarks remarks remarks

TABLE OF CONTENTS

31	List of Tables	vii
32	List of Figures	viii
33	KEY TO ABBREVIATIONS	x
34	Roadmap	1
35	Chapter 1. Introduction	2
36	Chapter 2. Theoretical Overview	3
37	2.1 The Standard Model	3
38	2.1.1 Elementary particles	3
39	2.1.2 Mathematical formalism	7
40	2.1.2.1 Quantum chromodynamics	8
41	2.1.2.2 Electroweak theory	9
42	2.1.2.3 Higgs mechanism	12
43	2.2 Beyond the Standard Model	15
44	2.2.1 Top-philic vector resonance	15
45	2.2.2 BSM four-top quark production	18
46	Chapter 3. LHC & ATLAS Experiment	20
47	3.1 The Large Hadron Collider	20
48	3.1.1 Overview	20
49	3.1.2 LHC operations	20
50	3.2 The ATLAS detector	21
51	3.2.1 Inner detector	22
52	3.2.2 Calorimeter systems	24
53	3.2.3 Muon spectrometer	25
54	3.2.4 Forward detectors	27
55	3.2.5 Magnetic systems	28
56	3.2.6 Trigger & data acquisition	28
57	Chapter 4. Particle Reconstruction & Identification	30
58	4.1 Primary reconstruction	30
59	4.1.1 Tracks	30
60	4.1.2 Vertices	31
61	4.1.3 Topological clusters	32
62	4.2 Jets	33
63	4.2.1 Jet reconstruction	34
64	4.2.2 Flavor tagging	35
65	4.3 Leptons	39

66	4.3.1	Electrons	39
67	4.3.2	Muons	42
68	4.4	Missing transverse momentum	44
69	4.5	Overlap removal	45
70	Chapter 5. Data & Simulated Samples		46
71	5.1	Data samples	46
72	5.2	Monte Carlo samples	46
73	5.2.1	$t\bar{t}Z'$ signal samples	47
74	5.2.2	Background samples	49
75	Chapter 6. Analysis Strategy		51
76	6.1	Event selection	51
77	6.1.1	Object definition	52
78	6.1.2	Event categorization	52
79	6.2	Analysis regions	53
80	6.2.1	Signal regions	55
81	6.2.2	Control regions	55
82	6.3	Background estimation	58
83	6.3.1	Template fitting for fake/non-prompt estimation	59
84	6.3.2	Charge misidentification data-driven estimation	59
85	6.3.3	$t\bar{t}W$ background data-driven estimation	61
86	Chapter 7. Systematic Uncertainties		64
87	7.1	Experimental uncertainties	64
88	7.1.1	Luminosity & pile-up reweighting	64
89	7.1.2	Leptons	64
90	7.1.3	Jets	66
91	7.1.4	Missing transverse energy	68
92	7.2	Modeling uncertainties	69
93	7.2.1	Signal and irreducible background uncertainties	69
94	7.2.2	Reducible background uncertainties	71
95	Chapter 8. Results		73
96	8.1	Statistical analysis	73
97	8.1.1	Profile likelihood fit	73
98	8.1.2	Exclusion limits	75
99	8.2	Fit results	75
100	Chapter 9. Summary		76
101	References		77

List of Tables

102		
103	Table 4.1:	Overlap removal process for this analysis, applied sequentially from top to
104		bottom. 45
105	Table 5.1:	Summary of all HLT triggers used in this analysis. Events are required to
106		pass at least one trigger. 47
107	Table 5.2:	Summary of all Monte-Carlo samples used in this analysis. V refers to
108		an EW ($W^\pm/Z/\gamma^*$) or Higgs boson. Matrix element (ME) order refers
109		to the order in QCD of the perturbative calculation. Tune refers to the
110		underlying-event tune of the parton shower (PS) generator. 48
111	Table 6.1:	Summary of object selection criteria used in this analysis. 52
112	Table 6.2:	Definitions of signal, control and validation regions (VR) used in this anal-
113		ysis. N_{jets} and N_b refers to the number of jets and number of b -tagged
114		jets respectively. ℓ_1 refers to the leading lepton, ℓ_2 refers to the subleading
115		lepton and so on. H_T refers to the p_T scalar sum of all leptons and jets
116		in the event. $m_{\ell\ell}$ refers to the dilepton invariant mass, which must not
117		coincide with the Z -boson mass range of 81-101 GeV for SS2L+3L events. 54
118	Table 6.3:	Definitions of SR sub-regions. Events are sorted into different sub-regions
119		based on the number of b -tagged jets and leptons present. 55
120	Table 6.4:	List of possible assigned values for DFCAA. 57
121	Table 7.1:	Summary of the experimental systematic uncertainties considered in this
122		analysis. 65

List of Figures

123		
124	Figure 2.1: Particles within the SM and their properties.	4
125	Figure 2.2: Feynman diagram for $t\bar{t}$ production and subsequent decay processes. Top	
126	quark decays into a W -boson and b -quark, and W -boson can decay to a	
127	$q\bar{q}$ or a $\ell\nu_\ell$ pair.	6
128	Figure 2.3: Illustration of a common representation of the Higgs potential. Before	
129	SSB, the ground state $\phi(0)$ is located at A which is symmetric with respect	
130	to the potential. A perturbation to this state fixes the ground state energy	
131	$ \phi(0) ^2$ to a particular value at B, "spontaneously" breaking the symmetry	
132	and degeneracy in $ \phi(0) ^2$	13
133	Figure 2.4: Feynman diagrams for tree level Z' production in association with (a) $t\bar{t}$,	
134	(b) tj (light quark) and (c) tW , decaying to final states containing (a) $t\bar{t}t\bar{t}$	
135	or (b)(c) $t\bar{t}t$	17
136	Figure 2.5: Branching ratios for $t\bar{t}t\bar{t}$ decay. The same-sign dilepton and multilepton	
137	channels together forms the SSML channel.	19
138	Figure 3.1: Caption	22
139	Figure 3.2: Caption	24
140	Figure 4.1: Stages of topo-cluster formation corresponding to each threshold. In (a),	
141	proto-clusters are seeded from cells with adequate signal significance $\zeta_{\text{cell}}^{\text{EM}}$.	
142	The clusters are further merged and split in (b) according to a predefined	
143	cluster growth threshold. The process stops in (c) when all sufficiently	
144	significant signal hits have been matched to a cluster.	33
145	Figure 4.2: Jet energy scale calibration sequence for EM-scale jets.	35

146	Figure 4.3:	Overview of the GN2 architecture. The number of jet and track features	
147		are represented by n_{jf} and n_{tf} respectively. The global jet representation	
148		and track embeddings output by the Transformer encoder are used as	
149		inputs for three task-specific networks.	36
150	Figure 4.4:	The c -, light- and τ -jet rejection rate as a function of b -tagging efficiency	
151		for GN2 and DL1d using (a) jets in the $t\bar{t}$ sample, and (b) jets in the Z'	
152		sample. The performance ratios of GN2 to DL1d are shown in the bottom	
153		panels.	38
154	Figure 6.1:	Charge flip rate calculated for SR and CR $t\bar{t}W$ in bins of $ \eta $ and p_{T} . . .	61
155	Figure 7.1:	Combined QmisID uncertainty rate for SR in bins of $ \eta $ and p_{T}	72

KEY TO ABBREVIATIONS

Physical & mathematical quantities

χ^2 chi-squared

ΔR angular distance

η pseudorapidity

E_T transverse energy

E_T^{miss} missing transverse momentum

γ_μ Dirac matrices

I weak isospin

L instantaneous luminosity

$m_{\ell\ell}$ dilepton invariant mass

μ signal strength

p_T transverse momentum

Particles

b bottom quark

pp proton-proton

$t\bar{t}$ top/anti-top quark

$t\bar{t}t\bar{t}$ four-top-quark

tW single-top

Acronyms

1LOS one lepton, or two leptons of opposite charges

AF3 AtlFast3 fast simulation

ATLAS A Toroidal LHC ApparatuS

BDT boosted decision tree

BSM Beyond the Standard Model

181 **CERN** European Organization for Nuclear Research
 182 **CKM** CabibboKobayashiMaskawa matrix
 183 **CMS** Compact Muon Solenoid
 184 **CR** control region
 185 **CSC** Cathode Strip Chambers
 186 **ECIDS** Electron Charge ID Selector
 187 **EM** electromagnetic
 188 **EW** electroweak
 189 **FS** full detector simulation
 190 **GNN** graph neural network
 191 **GRL** Good Run List
 192 **GSF** Gaussian-sum filter
 193 **GUT** Grand Unified Theory
 194 **HF** heavy-flavor
 195 **HLT** High-Level Trigger
 196 **ID** inner detector
 197 **JER** jet energy resolution
 198 **JES** jet energy scale
 199 **JVT** Jet Vertex Tagger
 200 **L1** Level 1
 201 **LH** likelihood
 202 **LLH** log-likelihood
 203 **LO** leading order
 204 **LAr** liquid argon
 205 **LHC** Large Hadron Collider
 206 **MC** Monte Carlo simulation
 207 **ME** matrix element

208 **ML** multilepton
 209 **MS** muon spectrometer
 210 **MDT** Monitored Drift Tubes
 211 **MET** missing transverse energy
 212 **NF** normalization factor
 213 **NNJvt** Neural Network-based Jet Vertex Tagger
 214 **NLO** next-to-leading order
 215 **NNLO** next-to-next-to-leading order
 216 **NP** nuisance parameter
 217 **OP** operating point (also working point
 218 **OS** opposite-sign
 219 **PCBT** pseudo-continuous b -tagging
 220 **PDF** parton distribution function
 221 **POI** parameter of interest
 222 **PS** parton shower
 223 **PV** primary vertex
 224 **QCD** quantum chromodynamics
 225 **QED** quantum electrodynamics
 226 **QFT** quantum field theory
 227 **QmisID** charge mis-identification
 228 **SCT** Semiconductor Tracker
 229 **SF** scale factor
 230 **SM** Standard Model
 231 **SR** signal region
 232 **SS** same-sign
 233 **SSB** spontaneous symmetry breaking
 234 **SS2L** same-sign dilepton

²³⁵ **SSML** same-sign dilepton, or more than two leptons of any charges
²³⁶ **TDAQ** Trigger and Data Acquisition
²³⁷ **TRT** Transition Radiation Tracker
²³⁸ **VEV** vacuum expectation value
²³⁹ **VR** validation region

240 **Roadmap**

241 1. Finish adding bullets for all sections 06/04

242 Remaining

243 • introduction

244 2. Fill in details 06/13

245 • Add missing figures

246 • Add missing bib

247 3. Finalize analysis

248 4. String everything together

249 5. Miscellaneous/logistics (proofreading, review, ATLAS approval, etc.)

250 6. Submission to the graduate school 07/01

251 7. Defense 07/15

Chapter 1. Introduction

[1]

1. background and context

2. problem to be solved in thesis

3. aim of analysis: Z' consequences of many BSM theories, searching for Z'

4. hypothesis/research question: searching for Z' in $t\bar{t}t\bar{t}$ SSML channel

5. methodology: data collection - \sqrt{s} analysis regions - \sqrt{s} binned likelihood fit

6. thesis structure:

- ch2: SM/BSM theoretical background
- ch3: LHC/ATLAS experiment
- ch4: samples used in the analysis
- ch5: ATLAS particle reconstruction and identification techniques, and object definitions for the analysis
- ch6: analysis strategy
- ch7: systematic uncertainties affecting the analysis
- ch8: final results
- ch9: summary

Chapter 2. Theoretical Overview

2.1 The Standard Model

The Standard Model of Physics (SM) [2] is currently the most successful formalism to describe the physical world at a microscopic scale by providing descriptions for all currently known elementary particles, along with three out of four fundamental forces (electromagnetism, weak force, strong force) with the exception of gravity. The SM is however not perfect, and there remains unanswered questions that require development and discovery of new physics beyond the Standard Model (BSM). This chapter describes an overview of important components within the SM and relevant BSM aspects for this analysis.

2.1.1 Elementary particles

Elementary particles in the SM can be classified into two groups: bosons consisting of particles following Bose-Einstein statistics with integer spin, and fermions consisting of particles following Fermi-Dirac statistics with half-integer spin. Fermions are the building blocks of composite particles and consequently all known matter, and can be further classified into quarks & leptons. Bosons act as force mediators for all fundamental forces described by the SM, and can either be a scalar boson with spin 0 or vector gauge bosons with spin 1. For each elementary particle, there also exists a corresponding antiparticle with identical mass and opposite charge (electric or color). Figure 2.1 shows all known elementary particles in the SM.




Figure 2.1: Particles within the SM and their properties [3].

Fermions

Fermions consist of quarks and leptons with six flavors each, grouped into three generations of doublets. The six quark flavors are up (u), down (d), charm (c), strange (s), bottom (b) and top (t), arranged in increasing order of mass. The quark flavors form three doublets (u, d), (c, s) and (t, b), with each doublet containing one quark with electric charge of $+2/3$ (u, s, t), and the other with charge of $-1/3$ (d, c, b). Each quark also possesses a property known as color charge, with possible values of red (R), green (G), blue (B) or their corresponding anticolor (\bar{R} , \bar{G} , \bar{B}). Color charge follows color confinement rules, which allows only configurations of quarks with total neutral color charge to exist in isolation. Neutral

charge configurations can be formed from either a set of three colors (R, G, B), a set of a color and its anticolor (q, \bar{q}), or any combination of the two. Consequently, quarks can only exist in bound states called hadrons and no isolated quark can be found in a vacuum. Quarks are the only elementary particles in the SM that can interact with all four fundamental forces.

The three leptons doublets consist of three charged leptons: electron (e), muon (μ), tau (τ), and their respective neutrino flavors: electron neutrino (ν_e), muon neutrino (ν_μ), tau neutrino (ν_τ). Charged leptons carry an electric charge of -1 , while their antiparticles carry the opposite charge ($+1$) and their corresponding neutrino flavors carry no charge. Charged leptons interact with all fundamental forces except the strong force, while neutrinos only interact with the weak force and gravity.

Bosons

The SM classifies bosons into two types: one scalar boson with spin 0 known as the Higgs (H) boson, and vector gauge bosons with spin 1 known as gluons (g), photon (γ), W^\pm and Z bosons. Gluons and photon are massless, while the W^\pm , Z and H bosons are massive. Each vector gauge boson serves as the mediator for a fundamental force described by the SM. Gluons are massless particles mediating the strong interaction by carrying color charges between quarks following quantum chromodynamics (QCD). Each gluon carries a non-neutral color charge out of eight linearly independent color states in the gluon color octet [4]. Photon is the massless and charge-neutral mediator particle for the electromagnetic interaction following quantum electrodynamics (QED). The W^\pm and Z bosons are massive mediator particles for the weak interaction, with the W^\pm boson carrying an electric charge of ± 1 while the Z boson is charge neutral.

Other than the vector gauge boson, the only scalar boson in the SM is the massive and

charge neutral Higgs boson. The Higgs boson does not mediate any fundamental force like vector bosons, but serve to provide the rest mass for all massive elementary particles in the SM through the Higgs mechanism described in section 2.1.2.3.

Top quark

As of now, the top quark (t) is the heaviest particle in the SM with mass of about 173 GeV [5]. For comparison, the heaviest boson, the Higgs boson, possesses mass of 125 GeV and the second most massive fermion, the b -quark has mass of about 4.2 GeV. This gives the top quark the strongest Yukawa coupling to the Higgs boson ($y_t \approx 1$) [6] and exotic resonances in many proposed BSM models [7–10], making the top quark and its processes attractive vehicles with which to probe new physics.

Due to its mass, the top quark has a very short lifetime of 10^{-24} s [5] and decays before it can hadronize following color confinement. The top quark decays to a W boson and a b -quark with a branching ratio of almost 100%. The W boson can subsequently decay hadronically or leptonically (Figure 2.2), with branching ratios of approximately 68% and 32% respectively. All lepton flavors have similar branching ratios during a leptonic W decay, assuming lepton universality.




Figure 2.2: Feynman diagram for $t\bar{t}$ production and subsequent decay processes [11]. Top quark decays into a W -boson and b -quark, and W -boson can decay to a $q\bar{q}$ or a $\ell\nu_\ell$ pair.

additional section on 4top production?

2.1.2 Mathematical formalism

The SM can be described within the formalism of quantum field theory (QFT) with the Lagrangian

$$\mathcal{L}_{\text{SM}} = \mathcal{L}_{\text{QCD}} + \underbrace{(\mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{fermion}} + \mathcal{L}_{\text{Higgs}} + \mathcal{L}_{\text{Yukawa}})}_{\mathcal{L}_{\text{EW}}} \quad (2.1)$$

where \mathcal{L}_{QCD} is the QCD term and \mathcal{L}_{EW} is the electroweak (EW) term of the Lagrangian. Formalism of QFT within the SM treats particles as excitations [12] of their corresponding quantum fields i.e. fermion field ψ , electroweak boson fields $W_{1,2,3}$ & B , gluon fields G_α and Higgs field ϕ .

The foundation of modern QFT involves gauge theory. A quantum field has gauge symmetry if there exists a continuous gauge transformation that when applied to every point in a field (local gauge transformation) leaves the field Lagrangian unchanged. The set of gauge transformations of a gauge symmetry is the symmetry group of the field which comes with a set of generators, each with a corresponding gauge field. Under QFT, the quanta of these gauge fields are called gauge bosons.

The SM Lagrangian is gauge invariant under global Poincaré symmetry and local $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge symmetry, with the $SU(3)_C$ symmetry group corresponding to the strong interaction and $SU(2)_L \times U(1)_Y$ to the EW interaction. Global Poincaré symmetry ensures that \mathcal{L}_{SM} satisfies translational symmetry, rotational symmetry and Lorentz boost frame invariance [13]. These symmetries give rise to corresponding conservation laws, which lead to conservation of momentum, angular momentum and energy in the SM as a result of Noether's theorem.

2.1.2.1 Quantum chromodynamics

Quantum chromodynamics is a non-Abelian gauge theory i.e. Yang-Mills theory [14, 15] describing the strong interaction between quarks in the SM with the gauge group $SU(3)_C$, where C represents conservation of color charge under $SU(3)_C$ symmetry. According to QFT, quarks can be treated as excitations of the corresponding quark fields ψ . The free Dirac Lagrangian for the quark fields

$$\mathcal{L}_0 = \bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi \quad (2.2)$$

is invariant under global $SU(3)$ symmetry, but not under local $SU(3)_C$ symmetry. To establish invariance under local $SU(3)_C$ symmetry, the gauge covariant derivative D_μ is defined so that

$$D_\mu\psi = (\partial_\mu - ig_s G_\mu^a T_a)\psi, \quad (2.3)$$

where $g_s = \sqrt{4\pi\alpha_s}$ is the QCD coupling constant, $G_\mu^a(x)$ are the eight gluon fields, and T_a are generators of $SU(3)_C$, represented as $T_a = \lambda_a/2$ with λ_a being the eight Gell-Mann matrices [4]. Let the gluon field strength tensors $G_{\mu\nu}^a$ be

$$G_{\mu\nu}^a \equiv \partial_\mu G_\nu^a - \partial_\nu G_\mu^a - g_s f^{abc} G_\mu^b G_\nu^c, \quad (2.4)$$

where f^{abc} are the structure constants of $SU(3)_C$. The gauge invariant QCD Lagrangian can then be written as

$$\begin{aligned}\mathcal{L}_{\text{QCD}} &= \bar{\psi}(i\gamma^\mu D_\mu - m)\psi - \frac{1}{4}G_{\mu\nu}^a G_a^{\mu\nu} \\ &= \underbrace{-\frac{1}{4}G_{\mu\nu}^a G_a^{\mu\nu}}_{\text{gluon kinematics \& self-interaction}} + \underbrace{\bar{\psi}(i\gamma^\mu \partial_\mu - m)\psi}_{\text{quark kinematics \& masses}} + \underbrace{\bar{\psi}^i (g_s \gamma^\mu (T_a)_{ij} G_\mu^a) \psi^j}_{\text{quark-gluon interaction}},\end{aligned}\quad (2.5)$$

where i, j are color indices with integer values from 1 to 3. Gluons are forced to be massless from the lack of a gluon mass term to maintain gauge invariance for the Lagrangian.

2.1.2.2 Electroweak theory

The electroweak interaction is the unified description of the weak interaction and electromagnetism under the $SU(2)_L \times U(1)_Y$ symmetry group, where L represents the left-handed chirality of the weak interaction and Y represents the weak hypercharge quantum number. Fermions can have either left-handed or right-handed chirality with the exception of neutrinos which can only have left-handed chirality in the SM, and can be divided into left-handed doublets and right-handed singlets

$$\begin{aligned}\psi_L &= \begin{pmatrix} \nu_e \\ e_L \end{pmatrix}, \begin{pmatrix} \nu_\mu \\ \mu_L \end{pmatrix}, \begin{pmatrix} \nu_\tau \\ \tau_L \end{pmatrix}, \begin{pmatrix} u_L \\ d_L \end{pmatrix}, \begin{pmatrix} c_L \\ s_L \end{pmatrix}, \begin{pmatrix} t_L \\ b_L \end{pmatrix} \\ \psi_R &= e_R, \mu_R, \tau_R, u_R, d_R, c_R, s_R, t_R, b_R.\end{aligned}\quad (2.6)$$

Similar to QCD, to establish invariance under local $U(1)_Y$ symmetry, the $U(1)_Y$ gauge covariant derivative D_μ is defined as

$$D_\mu \phi = \left(\partial_\mu - ig' \frac{Y}{2} B_\mu \right) \psi \quad (2.7)$$

388 where g' is the B_μ coupling constant and $B_\mu(x)$ is a vector gauge field that transforms under
 389 $U(1)_Y$ as

$$B_\mu \rightarrow B_\mu + \frac{1}{g'} \partial_\mu \theta(x). \quad (2.8)$$

390 Right-handed fermion singlets are not affected by $SU(2)_L$ transformation, so the fermion
 391 fields ψ transform under $SU(2)_L$ as

$$\begin{aligned} \psi_L &\rightarrow e^{iI_3 \vec{\theta}(x) \cdot \vec{\sigma}/2} \psi_L \\ \psi_R &\rightarrow \psi_R. \end{aligned} \quad (2.9)$$

392 where $\vec{\sigma}/2$ are generators of $SU(2)_L$ with $\vec{\sigma}$ being the Pauli matrices. In order to preserve
 393 local symmetry, let the gauge covariant derivative for $SU(2)_L$ be

$$D_\mu \psi_L = \left(\partial_\mu - ig \frac{\sigma_i}{2} W_\mu^i \right) \psi_L \quad (2.10)$$

394 where $W_\mu^i(x)$ ($i = 1, 2, 3$) are three boson gauge fields that transform under $SU(2)_L$ as

$$W_\mu^i \rightarrow e^{i \frac{\sigma_i}{2} \theta_i(x)} \left(W_\mu^i + \frac{i}{g} \partial_\mu \right) e^{-i \frac{\sigma_i}{2} \theta_i(x)} = W_\mu^i + \frac{2}{g} \partial_\mu \theta_a(x) + \epsilon^{ijk} \theta_j(x) W_\mu^k, \quad (2.11)$$

395 with g as the W_μ^i gauge coupling constant, and ϵ^{ijk} as the $SU(2)_L$ structure constant. The
 396 gauge covariant derivative for $SU(2)_L \times U(1)_Y$ can then be written as

$$\begin{aligned} D_\mu \psi_L &= \left(\partial_\mu - ig' \frac{Y_L}{2} B_\mu - ig \frac{\sigma_i}{2} W_\mu^i \right) \psi_L \\ D_\mu \psi_R &= \left(\partial_\mu - ig' \frac{Y_R}{2} B_\mu \right) \psi_R. \end{aligned} \quad (2.12)$$

397 Similar to QCD, the kinetic term is added by defining field strengths for the four gauge

398 fields

$$B_{\mu\nu} \equiv \partial_\mu B_\nu - \partial_\nu B_\mu \quad (2.13)$$

$$W_{\mu\nu}^i \equiv \partial_\mu W_\nu^i - \partial_\nu W_\mu^i - g e^{ijk} W_\mu^j W_\nu^k.$$

399 The local $SU(2)_L \times U(1)_Y$ invariant EW Lagrangian is then [16]

$$\begin{aligned} \mathcal{L}_{\text{EW}} &= i\bar{\psi}(\gamma^\mu D_\mu)\psi - \frac{1}{4}W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu} \\ &= \underbrace{i\bar{\psi}(\gamma^\mu \partial_\mu)\psi}_{\text{fermion kinematics}} - \underbrace{\bar{\psi}\left(\gamma^\mu g' \frac{Y}{2} B_\mu\right)\psi - \bar{\psi}_L\left(\gamma^\mu g \frac{\sigma_i}{2} W_\mu^i\right)\psi_L}_{\text{fermion-gauge boson interaction}} - \underbrace{\frac{1}{4}W_{\mu\nu}^i W_i^{\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu}}_{\text{boson kinematics \& self-interaction}}. \end{aligned} \quad (2.14)$$

400 Under ≈ 159.5 GeV, the EW symmetry $SU(2)_L \times U(1)_Y$ undergoes spontaneous symmetry
 401 breaking [17] into $U(1)_{\text{QED}}$ symmetry, which corresponds to a separation of the weak and
 402 electrodynamic forces. Electroweak spontaneous symmetry breaking replaces the four mass-
 403 less and similarly-behaved EW gauge bosons B_μ and W_μ^i with the EM boson γ and the weak
 404 bosons Z/W^\pm , as well as giving the Z and W^\pm bosons masses via the Higgs mechanism.
 405 This is due to a specific choice of gauge for the Higgs field leading to the reparameterization
 406 of the EW bosons B_μ and W_μ^i to $W^\pm/Z/\gamma$ using the relations

$$\begin{aligned} W_\mu^\pm &\equiv \frac{1}{\sqrt{2}}(W_\mu^1 \mp iW_\mu^2) \\ \begin{pmatrix} A_\mu \\ Z_\mu \end{pmatrix} &\equiv \begin{pmatrix} \cos \theta_W & \sin \theta_W \\ -\sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^3 \end{pmatrix} \end{aligned} \quad (2.15)$$

407 where $\theta_W \equiv \cos^{-1}\left(g/\sqrt{g^2 + g'^2}\right)$ is the weak mixing angle. The boson kinetic term can also
 408 be refactorized to extract cubic (three vertices) and quartic (four vertices) self-interactions

among the gauge bosons [16]. The Lagrangian can then be rewritten as

$$\begin{aligned}
\mathcal{L} = & \underbrace{e A_\mu \bar{\psi} (\gamma^\mu Q) \psi}_{\text{electromagnetism}} + \underbrace{\frac{e}{2 \sin \theta_W \cos \theta_W} \bar{\psi} \gamma^\mu (v_f - a_f \gamma_5) \psi Z_\mu}_{\text{neutral current interaction}} \\
& + \underbrace{\frac{g}{2\sqrt{2}} \sum_{\psi_L} [\bar{f}_2 \gamma^\mu (1 - \gamma_5) f_1 W_\mu^+ + \bar{f}_1 \gamma^\mu (1 - \gamma_5) f_2 W_\mu^-]}_{\text{charged current interaction}} \\
& + \mathcal{L}_{\text{kinetic}} + \underbrace{\mathcal{L}_{\text{cubic}} + \mathcal{L}_{\text{quartic}}}_{\text{boson self-interaction}}
\end{aligned} \tag{2.16}$$

where $\gamma_5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ is the chirality projection operator, $a_f = I_3$, $v_f = I_3(1 - 4|Q| \sin^2 \theta_W)$ and f_1, f_2 are up and down type fermions of a left-handed doublet.

2.1.2.3 Higgs mechanism

So far, the EW bosons are massless since the mass terms $-m\bar{\psi}\psi$ for fermions and $-mA^\mu A_\mu$ for bosons are not invariant under the EW Lagrangian symmetries. The particles must then acquire mass under another mechanism. The Brout-Engler-Higgs mechanism [18–20] was introduced in 1964 to rectify this issue and verified in 2012 with the discovery of the Higgs boson [21, 22].

The Higgs potential is expressed as

$$V(\phi^\dagger \phi) = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 \tag{2.17}$$

where μ^2 and $\lambda > 0$ are arbitrary parameters, and the $SU(2)_L$ doublet $\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$ is the Higgs field with complex scalar fields ϕ^+ and ϕ^0 carrying +1 and 0 electric charge respectively.

421 The Lagrangian for the scalar Higgs field is

$$\mathcal{L}_H = (\partial_\mu \phi)^\dagger (\partial^\mu \phi) - V(\phi^\dagger \phi). \quad (2.18)$$

422 Since the potential $V(\phi^\dagger \phi)$ is constrained by $\lambda > 0$, the ground state is solely controlled by
 423 μ . If $\mu^2 > 0$, the ground state energy is $\phi = 0$, and the EW bosons would remain massless.

424 If $\mu^2 < 0$, the ground state is

$$|\phi|^2 = -\frac{\mu^2}{2\lambda} \equiv \frac{v^2}{\sqrt{2}}, \quad (2.19)$$

425 where v is defined as the vacuum expectation value (VEV). The standard ground state for
 426 the Higgs potential without loss of generality can be chosen as $\phi(0) = 1/\sqrt{2} \begin{pmatrix} 0 \\ v \end{pmatrix}$.




Figure 2.3: Illustration of a common representation of the Higgs potential [23]. Before SSB, the ground state $\phi(0)$ is located at A which is symmetric with respect to the potential. A perturbation to this state fixes the ground state energy $|\phi(0)|^2$ to a particular value at B, "spontaneously" breaking the symmetry and degeneracy in $|\phi(0)|^2$.

427 Having $U(1)$ symmetry allows any $-e^{i\theta} \sqrt{\mu^2/\lambda}$ to be a ground state energy for the Higgs
 428 Lagrangian. This degeneracy results in spontaneous symmetry breaking of the $SU(2)_L \times$
 429 $U(1)_Y$ symmetry into $U(1)_{\text{EM}}$ symmetry when the Higgs field settles on a specific vacuum
 430 state as a result of a perturbation or excitation (Figure 2.3). The spontaneous symmetry

431 breaking introduces three massless (Nambu-Goldstone [24]) vector gauge boson ξ and a
 432 massive scalar boson η , each corresponds to a generator of the gauge group. The vector field
 433 for ξ and η are real fields parameterized as $\xi \equiv \phi^+ \sqrt{2}$ and $\eta \equiv \phi^0 \sqrt{2} - v$ [25]. The Higgs
 434 field now becomes

$$\phi = \frac{v + \eta + i\xi}{\sqrt{2}} = \frac{1}{\sqrt{2}} e^{i\xi \cdot \frac{\sigma}{2v}} \begin{pmatrix} 0 \\ v + \eta \end{pmatrix}. \quad (2.20)$$

435 Due to $U(1)_{\text{EM}}$ invariance, a unitary gauge with the transformation $\phi \rightarrow \exp(-i\xi \cdot \frac{\sigma}{2v})$ can
 436 be chosen for the Higgs field to eliminate the massless bosons and incorporate them into the
 437 EM/weak bosons via Equation 2.15. This leaves the massive η which can now be observed as
 438 an excitation of the Higgs field from the standard ground state and must be the Higgs boson
 439 h . Using the EW covariant derivative from Equation 2.12, the Higgs Lagrangian around the
 440 vacuum state becomes

$$\begin{aligned} \mathcal{L}_H &= (D_\mu \phi)^\dagger (D^\mu \phi) - \mu^2 \left(\frac{v+h}{\sqrt{2}} \right)^2 - \lambda \left(\frac{v+h}{\sqrt{2}} \right)^4 \\ &= (D_\mu \phi)^\dagger (D^\mu \phi) - \frac{1}{2} \mu^2 h^2 - \lambda v h^3 - \frac{\lambda}{4} h^4 - \dots \end{aligned} \quad (2.21)$$

441 The Higgs mass can be extracted from the quadratic term as $m_H = \sqrt{-2\mu^2}$. The kinetic
 442 term in the Lagrangian can be written as

$$\begin{aligned} (D_\mu \phi)^\dagger (D^\mu \phi) &= \frac{1}{2} (\partial_\mu h)^2 + \frac{g^2}{8} (v+h)^2 |W_\mu^1 - iW_\mu^2|^2 + \frac{1}{8} (v+h)^2 (g'W_\mu - gB_\mu)^2 \\ &= \frac{1}{2} (\partial_\mu h)^2 + (v+h)^2 \left(\frac{g^2}{4} W_\mu^+ W^{-\mu} + \frac{1}{8} (g^2 + g'^2) Z_\mu^0 Z^{0\mu} \right). \end{aligned} \quad (2.22)$$

443 Masses for the EW bosons can be extracted from the quadratic terms

$$m_{W^\pm} = \frac{v}{2} g, \quad m_Z = \frac{v}{2} \sqrt{g^2 + g'^2}, \quad m_\gamma = 0. \quad (2.23)$$

444 However, the fermion mass term $-m\bar{\psi}\psi$ still breaks EW invariance after spontaneous sym-
 445 metry breaking. Instead, fermions acquire mass by replacing the mass term with a gauge
 446 invariant Yukawa term in the EW Lagrangian representing fermions' interactions with the
 447 Higgs field [25]

$$\begin{aligned}
 \mathcal{L}_{\text{Yukawa}} &= -c_f \frac{v+h}{\sqrt{2}} (\bar{\psi}_R \psi_L + \bar{\psi}_L \psi_R) \\
 &= - \underbrace{\frac{c_f}{\sqrt{2}} v (\bar{\psi}\psi)}_{\text{fermion mass}} - \underbrace{\frac{c_f}{\sqrt{2}} (h\bar{\psi}\psi)}_{\text{fermion-Higgs interaction}}, \tag{2.24}
 \end{aligned}$$

448 where c_f is the fermion-Higgs Yukawa coupling. The fermion mass is then $m_f = c_f v / \sqrt{2}$.

449 2.2 Beyond the Standard Model

450 2.2.1 Top-philic vector resonance

451 Many BSM models extend the SM by adding to the SM gauge group additional $U(1)'$
 452 gauge symmetries [26], each with an associated vector gauge boson nominally called Z' . In
 453 the case of a BSM global symmetry group with rank larger than the SM gauge group, the
 454 symmetry group can spontaneously break into $G_{\text{SM}} \times U(1)'^n$, where G_{SM} is the SM gauge
 455 group $SU(3)_C \times SU(2)_L \times U(1)_Y$ and $U(1)'^n$ is any $n \geq 1$ number of $U(1)'$ symmetries. The
 456 existence of additional vector bosons Z' would open up many avenues of new physics e.g.
 457 extended Higgs sectors from $U(1)'$ symmetry breaking [27, 28], existence of flavor-changing
 458 neutral current (FCNC) mediated by Z' [29], and possible exotic production from heavy Z'
 459 decays [30].

460 Due to the top quark having the largest mass out of all known elementary particles in
 461 the SM, many BSM models [31–34] predict 'top-philic' vector resonances that have much

stronger coupling to the top quark compared to other quarks. The analysis in this thesis attempts to reconstruct a top-philic Z' resonance directly to avoid dependency on model choice. Previous model-independent BSM $t\bar{t}t\bar{t}$ search for top-philic resonances [35] in the single-lepton final state and similar mass ranges showed upper limits on observed (expected) Z' production cross section between 21 (14) fb to 119 (86) fb depending on parameter choice. This analysis is also motivated by the recent observation of SM $t\bar{t}t\bar{t}$ production in the same-sign multilepton (SSML) channel by ATLAS [36] and CMS [37] at 6.1σ and 5.6σ discovery significance respectively.

In addition to the model-independent search, a simplified color-singlet vector particle model [38, 39] is employed to study model-dependent interpretations. The interaction Lagrangian assumes only Z' to top coupling and has the form

$$\begin{aligned}\mathcal{L}_{Z'} &= \bar{t}\gamma_\mu (c_L P_L + c_R P_R) t Z'^\mu \\ &= c_t \bar{t}\gamma_\mu (\cos\theta P_L + \sin\theta P_R) t Z'^\mu,\end{aligned}\tag{2.25}$$

where $c_t = \sqrt{c_L^2 + c_R^2}$ is the Z' -top coupling strength, $P_{L/R} = (1 \mp \gamma_5)/2$ are the chirality projection operators, and $\theta = \tan^{-1}(c_R/c_L)$ is the chirality mixing angle. Expanding the Lagrangian results in

$$\mathcal{L}_{Z'} = \frac{1}{\sqrt{2}} \bar{t}\gamma_\mu \left[\sin\left(\theta + \frac{\pi}{4}\right) - \left(\sqrt{2} \cos\left(\theta + \frac{\pi}{4}\right)\right) \gamma_5 \right] t Z'^\mu,\tag{2.26}$$

which bears striking resemblance to the EW Lagrangian neutral current interaction term in Equation 2.16, showing the similarity between the Z' and the Z boson that acquires mass from $SU(2)_L \times U(1)_Y$ spontaneous symmetry breaking. Assuming the Z' mass $m_{Z'}$ is much larger than the top mass ($m_t^2/m_{Z'}^2 \approx 0$), the Z' decay width at leading-order (LO) can be

approximated as

$$\Gamma(Z' \rightarrow t\bar{t}) \approx \frac{c_t^2 m_{Z'}}{8\pi}. \quad (2.27)$$

It can be observed that $\Gamma/m_{Z'} \approx c_t^2/8\pi \ll 1$ for $c_t \approx 1$, which suggests a very narrow and well-defined resonance peak. This validates the narrow-width approximation for the choice of $c_t = 1$ and supports efforts to directly reconstruct the resonance.

The main production channels for the aforementioned heavy top-philic color singlet Z' are at tree level and loop level, with the one-loop level being the dominant processes. Loop level processes are dependent on the chirality angle θ , where $\theta = \pi/4$ suppresses all but gluon-initiated box sub-processes [38]. To minimize model dependence, only the tree level production was considered for this analysis by choosing $\theta = \pi/4$. Figure 2.4 illustrates several tree level Z' production processes.




Figure 2.4: Feynman diagrams for tree level Z' production in association with (a) $t\bar{t}$, (b) tj (light quark) and (c) tW , decaying to final states containing (a) $t\bar{t}t\bar{t}$ or (b)(c) $t\bar{t}t$ [38].

The single-top-associated channels tjZ' and tWZ' are suppressed by three-body phase space [38], resulting in smaller cross sections by a factor of two compared to the $t\bar{t}$ -associated process $t\bar{t}Z'$. Unlike tjZ' and tWZ' which are produced via EW and mixed QCD-EW interactions respectively, $t\bar{t}t\bar{t}$ production is governed by the strong interaction which can overpower phase space suppression. Additionally, $t\bar{t}t\bar{t}$ production is independent of θ while

tjZ' and tWZ' are minimally suppressed under pure left-handed interactions ($\theta = 0$) and maximally suppressed under pure right-handed interactions ($\theta = \pi/2$).

2.2.2 BSM four-top quark production

This analysis uses the $t\bar{t}t\bar{t}$ final state signal signature to search for the existence of a heavy BSM resonance that couples strongly to the top quark. Cross section for $t\bar{t}t\bar{t}$ production can be enhanced by many possible BSM models, in particular two-Higgs-doublet-models [40–42] (2HDM) or possible production of a heavy neutral resonance boson $Z'(\rightarrow t\bar{t})$ in association with a $t\bar{t}$ pair in composite Higgs scenarios [31, 32]. The $t\bar{t}Z'$ production mode and consequently $t\bar{t}t\bar{t}$ signal signature can provide a more sensitive channel for searches by avoiding contamination from the large SM $gg \rightarrow t\bar{t}$ background in an inclusive $Z' \rightarrow t\bar{t}$ search.

Decay modes

The different W boson decay modes shown in Figure 2.2 result in many different final states for $t\bar{t}Z'/t\bar{t}t\bar{t}$ decay, which can each be classified into one of three channels: all hadronic decays; exactly one lepton or two opposite-sign leptons (1LOS); exactly two same-sign leptons or three or more leptons (SSML). The branching ratio for each channel is shown in Figure 2.5. The all hadronic and 1LOS channels have much larger branching ratios compared to SSML channel but suffer heavily from $gg \rightarrow t\bar{t}$ background contamination, giving the SSML channel better sensitivity at the cost of lower statistics. This is also the targeted channel for this analysis.




Figure 2.5: Branching ratios for $t\bar{t}t\bar{t}$ decay [43]. The same-sign dilepton and multilepton channels together forms the SSML channel.

Chapter 3. LHC & ATLAS Experiment

3.1 The Large Hadron Collider

theoretical predictions are tested with experimental data obtained from particle accelerators world's largest accelerator built by CERN situated on the border of Switzerland and France has been operating since xxxx lifetime divided into 3 runs, currently on Run 3 with planned upgrades on the horizon responsible for a number of discoveries aka Higgs, etc.

3.1.1 Overview

[Basic info: location, size, main working mechanism, main detectors, main physics done]
- 27 km circumference, reusing LEP tunnels 175 m below ground level
- 7-13-13.6 TeV center of mass energies for pp collisions
- other than pp, also collides pPb, PbPb at 4 points with 4 main detectors: ATLAS, CMS (general purpose detectors), ALICE (heavy ion physics, ion collisions), LHCb (*b*-physics)

3.1.2 LHC operations

- focuses mainly on pp collisions for this thesis - beams split into bunches of 1.1×10^{11} protons with instantaneous luminosity of up to $2 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- beam energies ramp up in other accelerators before injection, full ramp up to 6.5 GeV about 20 minutes
(insert full diagram of accelerator chain)

Linac 4: hydrogen atoms, accelerated up to 160 MeV

PSB: H atoms stripped of electrons before injection, accelerated to 2 GeV

PS: 26 GeV, SPS: 450 GeV

LHC: injection in opposite directions, 6.5 TeV per beam

Run 1: 2010-2012, Run 2: 2015-2018, Run 3: 2022-2025, HL-LHC: 2029-?

COM energies: 7 & 8 TeV, 13 TeV, 13.6 TeV, 13.6 & 14 TeV

inbetween periods: long shutdowns (LS1, LS2, LS3)

Physics at the LHC

3.2 The ATLAS detector

multipurpose particle detector with a symmetric cylindrical geometry and a solid angle coverage of almost 4π

44m long, 25m diameter

inner detector, solenoid/toroid magnet, EM & hadronic calorimeters, muon spectrometer

(insert figure)

right-handed cylindrical system, z-axis follows beamline, azimuthal and polar (0 in the beam direction) angles measured with respect to beam axis.

pseudorapidity $\eta = -\ln \tan(\theta/2)$, approaches $\pm \infty$ along and 0 orthogonal to the beamline

distance $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2}$

transverse energy $E_T = \sqrt{p_T^2 + m^2}$




Figure 3.1: Caption [44]

transverse momentum p_T component of momentum orthogonal to the beam axis $p_T =$

$$\sqrt{p_x^2 + p_y^2}$$

3.2.1 Inner detector

- measures tracks of charged particles with high momentum resolution ($\sigma_{p_T}/p_T = 0.05\% \pm 1\%$)

- covers particles with $p_T > 0.5 \text{ GeV}$, $|\eta| < 2.5$

pixel detector -; semiconductor tracker -; transition radiation tracker, innermost to

outermost

- pixel detector:

- innermost, 250 μm silicon pixel layers

- detects charged particles from electron-hole pair production in silicon

- measures impact parameter resolution & vertex identification for reconstruction of short-lived particles

- spatial resolution of 10 μm in the $R - \phi$ plane and 115 μm in the z-direction

- 80.4m readout channels

- sct:

- surrounds pixel detector, silicon microstrip layers with 80 μm strip pitch

- particle tracks cross 8 strip layers

- measures particle momentum, impact parameters, vertex position

- spatial resolution of 17 μm in the $R - \phi$ plane and 580 μm in the z-direction

- 6.3m readout channels.

- trt:

- outermost, layers of 4 mm diameter gaseous straw tubes with transition radiation material (70% Xe + 27% CO_2 + 3% O_2) & 30 μm gold-plated wire in the center

- tubes 144 cm length in barrel region ($|\eta| < 1$), 37 cm in the endcap region ($1 < |\eta| < 2$), arranged in wheels instead of parallel to beamline)

- gas mixture produces transition radiation when ionized for electron identification

- resolution/accuracy of 130 μm for each straw tube in the $R - \phi$ plane
- 351k readout channels

3.2.2 Calorimeter systems

surrounds the inner detector & solenoid magnet, covers $|\eta| < 4.9$ and full ϕ range. Alternates passive and active material layers. Incoming particles passing through calorimeter produce EM cascades or hadronic showers in passive layer. Energies deposited and convert to electric signals in active layers for readout.




Figure 3.2: Caption [45]

EM calorimeter:

- innermost, lead-LAr detector (passive & active)

- measures EM cascades (bremsstrahlung & pair production) produced by electrons/photons

- divided into barrel region ($|\eta| < 1.475$) & endcap regions ($1.375 < |\eta| < 3.2$) with transition region ($1.372 < |\eta| < 1.52$) containing extra cooling materials for inner detector
- end-cap divided into outer wheel ($1.372 < |\eta| < 2.5$) & inner wheel ($2.5 < |\eta| < 3.2$)
- higher granularity in ID ($|\eta| < 2.5$) range for electrons/photons & precision physics, coarser elsewhere for jet reconstruction & MET measurements
- hadronic calorimeter:
- outermost
- measures hadronic showers from inelastic QCD collisions
- thick enough to prevent most particles from reaching muon spectrometer

- split into tile calorimeter in barrel region ($|\eta| < 1.0$) & extended barrel region ($0.8 < |\eta| < 1.7$), LAr hadronic end-cap calorimeter (HEC) in end-cap regions ($1.5 < |\eta| < 3.2$) & LAr forward calorimeters (FCal) in $3.1 < |\eta| < 4.9$ range.

- tile calorimeters: steel-plastic scintillating tiles, readout via photomultiplier tubes
- hec: behind tile calorimeters, 2 wheels per end-cap. copper plates-LAr. overlap with other calorimeter systems to cover for gaps between subsystems
- fcal: 1 copper module & 2 tungsten modules-LAr. copper optimized for EM measurements, tungsten for hadronic.

3.2.3 Muon spectrometer

- ATLAS outermost layer. measures muon momenta & charge in range $|\eta| < 2.7$

- 637 • momentum measured by deflection in track from toroid magnets producing magnetic
638 field orthogonal to muon trajectory
- 639 – large barrel toroids in $|\eta| < 1.4$, strength 0.5 T
- 640 – 2 smaller end-cap toroids in $1.6 < |\eta| < 2.7$, strength 1 T
- 641 – transition region $1.4 < |\eta| < 1.6$, deflection provided by a combination of barrel
642 and end-cap magnets
- 643 • chambers installed in 3 cylindrical layers, around the beam axis in barrel region & in
644 planes perpendicular to beam axis in the transition and end-cap regions
- 645 • split into high-precision tracking chambers (monitored drift tubes & cathode strip
646 chambers) & trigger chambers (resistive plate chambers & thin gap chambers)
- 647 • trigger chambers provide fast muon multiplicity & approximate energy range informa-
648 tion with L1 trigger logic
- 649 – mdt: 658 * muons pass through tube, ion-
650 * range $|\eta| < 2.7$, innermost layer 659 izing gas and providing signals.
651 $|\eta| < 2.0$ 660 Combining signals from tubes
652 * precision momentum measure- 661 forms track
653 ment 662 * maximumn drift time from wall
654 * layers of 30 mm drift tubes filled 663 to wire 700 ns
655 with 93% *Ar* & 7% *CO*₂, with 664 * resolution: 35 μm per chamber,
656 a 50 μm gold-plated tungsten- 665 80 μm per tube
657 rhenium wire at the center 666 – csc:

667 * forward region $2.0 < |\eta| < 2.7$ – rpc:
 668 highest particle flux and density 683 * range $|\eta| < 1.05$
 669 region 684 * provide fast meas
 670 * multiwire proportional chambers 685 – tgc:
 671 with higher granularity, filled 686 * range $1.05 < |\eta| < 2.7$
 672 with 80% *Ar* & 20% *CO*₂
 673 * shorter drift time than MDT,
 674 plus other features making CSC
 675 suitable for high particle den-
 676 sities and consequently able to
 677 handle background conditions
 678 * resolution: 40 μm in bending η -
 679 plane, 5 mm in nonbending ϕ -
 680 plane due to coarser cathode seg-
 681 mentation, per CSC plane

687 3.2.4 Forward detectors

- 688 • LUCID (LUminosity measurement using Cherenkov Integrating Detector): ± 17 m from
 689 interaction point, measures luminosity using *pp* scattering in the forward region
- 690 • ALFA (Absolute Luminosity for ATLAS): ± 240 m, measures *pp* scattering at small
 691 angles
- 692 • ZDC (Zero-Degree Calorimeter): ± 140 m, measures centrality in heavy-ion collisions

3.2.5 Magnetic systems

superconducting solenoid & toroid magnets cooled to 4.5 K with liquid helium

solenoid: 2.56 m diameter, 5.8 m length, 2 T strength axial magnetic field, encloses inner detector

toroid = barrel + endcap toroid x2

barrel toroid: 9.2/20.1 m inner/outer diameter, 25.3 m length, 0.5 T strength

endcap toroid: 1.65/10.7 m inner/outer diameter, 5 m length, 1 T strength

(show magnet system diagram)

3.2.6 Trigger & data acquisition

LHC produces large amount of data (40 MHz with 25 ns bunch crossing), necessitates a way to filter out trash from interesting events

handles online processing, selecting and recording interesting events for further offline processing and more in-depth analyses

- Level-1 (L1) trigger: online, fast hardware-based trigger, reduces to 100 kHz
 - L1 calorimeter triggers (L1Calo): selects high energy objects & MET
 - L1 muon triggers (L1Muon): selects using hit information from RPC & TGC
 - L1 topological trigger (L1Topo): select based on topological selection synthesized using information from L1Calo & L1Muon
 - Central Trigger Processor (CTP): uses L1Calo/Muon/Topo for final L1 trigger decision within $2.5 \mu\text{s}$ latency. Also identify regions of interest in η and ϕ to be

714 processed directly by HLT

715 • L1 trigger information read out by Front-End (FE) detector electronics then sent to
716 ReadOut Drivers (ROD) for preprocessing and subsequently to ReadOut System (ROS)
717 to buffer

718 • High-Level Trigger (HLT): offline, software-based trigger, using dedicated algorithms
719 and L1 output as input, reduces to 1 kHz

720 • Send to storage for analyses after HLT

721 overall trigger process reduces original collision data rate by a factor of about 10000 after
722 HLT

723 (show TDAQ diagram)

Chapter 4. Particle Reconstruction & Identification

Activity within the ATLAS detector are recorded as raw electronic signals, which can be utilized by ATLAS reconstruction software to derive physics objects for analysis. This chapter describes the reconstruction and identification of basic objects (e.g. interaction vertices, tracks, topological clusters of energy deposits) and subsequently of complex physics objects i.e. particles and particle signatures.

4.1 Primary reconstruction

4.1.1 Tracks

Charged particles traveling through the ATLAS detector deposit energy in different layers of the ID and MS. The ID track reconstruction software consists of two algorithm chains: inside-out and outside-in track reconstruction [46–48].

The inside-out algorithm is primarily used for the reconstruction of primary particles i.e. particles directly produced from pp collisions or decay products of short-lived particles. The process starts by forming space points from seeded hits in the silicon detectors within the pixel & SCT detectors. Hits further away from the interaction vertex are added to the track candidate using a combinatorial Kalman filter [49] pattern recognition algorithm. Track candidates are then fitted with a χ^2 filter [50] and loosely matched to a fixed-sized EM cluster. Successfully matched track candidates are re-fitted with a Gaussian-sum filter (GSF) [51], followed by a track scoring strategy to resolve fake tracks & hit ambiguity

between different tracks [52]. The track candidate is then extended to the TRT to form final tracks satisfying $p_T > 400$ MeV. The outside-in algorithm handles secondary tracks mainly produced from long-lives particles or decays of primary particles by back-tracking from TRT segments, which are then extended inward to match silicon hits in the pixel and SCT detectors to form track reconstruction objects.

4.1.2 Vertices

Vertices represent the point of interaction or decay for particles within the ATLAS detector. Primary vertices (PVs) are defined as the point of collision for hard-scattering pp interactions, while secondary or displaced vertices result from particle decays occurring at a distance from its production point.

Reconstruction of PVs is crucial to accurately profile the kinematic information of an event and form a basis for subsequent reconstruction procedures. Primary vertex reconstruction occurs in two stages: vertex finding and vertex fitting [53]. The vertex finding algorithm uses the spatial coordinates of reconstructed tracks to form the seed for a vertex candidate. An adaptive vertex fitting algorithm [54] then iteratively evaluates track-vertex compatibility to estimate a new best vertex position. Less compatible tracks are down-weighted in each subsequent iteration, and incompatible tracks are removed and can be used for another vertex seed; the process is repeated until no further PV can be found. All reconstructed vertices without at least two matched tracks are considered invalid and discarded.

Secondary vertex reconstruction uses the Secondary Vertex Finder (SVF) algorithm [55] which is primarily designed to reconstruct b - and c -hadrons for flavor tagging purposes. The SVF aims to reconstruct one secondary vertex per jet and only considers tracks that

are matched to a two-track vertex and contained within a p_T -dependent cone around the jet axis. The tracks are then used to reconstruct a secondary vertex candidate using an iterative process similar to the PV vertex fitting procedure.

Pile-up

At high luminosities, multiple interactions can be associated with one bunch crossing, resulting in many PVs. The effect is called pile-up, and usually result from soft QCD interactions. Pile-up can be categorized into two types: in-time pile-up, stemming from additional pp collisions in the same bunch crossing that is not the hard-scatter process; out-of-time pile-up, resulting from leftover energy deposits in the calorimeters from other bunch crossings [56].

4.1.3 Topological clusters

Topological clusters (topo-clusters) [57] consist of clusters of spatially related calorimeter cell signals. Topo-clusters are primarily used to reconstruct hadron- and jet-related objects in an effort to extract signal while minimizing electronic effects and physical fluctuations, and also allow for recovery of energy lost through bremsstrahlung or photon conversions. Cells with signal-to-noise ratio $\varsigma_{\text{cell}}^{\text{EM}}$ passing a primary seed threshold are seeded into a dynamic topological cell clustering algorithm as part of a proto-cluster. Neighboring cells satisfying a cluster growth threshold are collected into the proto-cluster. If a cell is matched to two proto-clusters, the clusters are merged. Two or more local signal maxima in a cluster satisfying $E_{\text{cell}}^{\text{EM}} > 500 \text{ MeV}$ suggest the presence of multiple particles in close proximity, and the cluster is split accordingly to maintain good resolution of the energy flow. The process continues iteratively until all cells with $\varsigma_{\text{cell}}^{\text{EM}}$ above a principal cell filter level have been matched to a

789 cluster.




Figure 4.1: Stages of topo-cluster formation corresponding to each threshold. In (a), proto-clusters are seeded from cells with adequate signal significance $\zeta_{\text{cell}}^{\text{EM}}$. The clusters are further merged and split in (b) following a predefined cluster growth threshold. The process stops in (c) when all sufficiently significant signal hits have been matched to a cluster [57].

790 4.2 Jets

791 Quarks, gluons and other hadrons with non-neutral color charge cannot be observed
 792 individually due to QCD color confinement, which forces a non-color-neutral hadron to

almost immediately undergo hadronization, producing a collimated cone of color-neutral hadrons defined as a jet. Jet signals can be used to reconstruct and indirectly observe the quarks or gluons from which the jet originated in the original hard-scattering process.

4.2.1 Jet reconstruction

The ATLAS jet reconstruction pipeline is largely carried out using a particle flow (PFlow) algorithm combined with an anti- k_t jet clustering algorithm. The PFlow algorithm [58] utilizes topo-clusters along with information from both the calorimeter systems and the ID in order to make use of the tracker system’s advantages in low-energy momentum resolution and angular resolution. First, the energy from charged particles is removed from the calorimeter topo-clusters; then, it is replaced by particle objects created using the remaining energy in the calorimeter and tracks matched to topo-clusters. The ensemble of ”particle flow objects” and corresponding matched tracks are used as inputs for the iterative anti- k_t algorithm [59].

The main components of the anti- k_t algorithm involve the distance d_{ij} between two jet candidates i and j , and the distance d_{iB} between the harder jet candidate of the two (defined as i) and the beamline B . If $d_{ij} < d_{iB}$, then the two jet candidates are combined and returned to the pool of candidates; otherwise, jet candidate i is considered a jet and removed from the pool. The distance d_{ij} is inversely proportional to a predefined radius parameter ΔR in order to control reconstruction quality for small- R and large- R jets. This analysis uses $\Delta R = 0.4$ to better handle heavily collimated small- R jets resulting from parton showers.

The anti- k_t jets so far have only been reconstructed at the EM level and need to be calibrated to match the energy scale of jets reconstructed at particle level. This is done via a MC-based jet energy scale (JES) calibration sequence, along with further calibrations

to account for pile-up effects and energy leakage. The full JES calibration sequence is shown in Figure 4.2. All calibration except origin correction are applied to the jet’s four-momentum i.e. jet p_T , energy and mass. Afterwards, a jet energy resolution (JER) [60] calibration step is carried out in a similar manner to JES to match the resolution of jets in dijet events. To further suppress pile-up effects, a neural-network based jet vertex tagger (NNJvt) discriminant was developed based on the previous jet vertex tagger (JVT) algorithm [56] and applied to low- p_T reconstructed jets.




Figure 4.2: Jet energy scale calibration sequence for EM-scale jets [61].

4.2.2 Flavor tagging

Identifying and classifying hadronic jets are important tasks for ATLAS physics, for example analyses involving Higgs decays $H \rightarrow b\bar{b}$ or top quarks. Flavor tagging or b -tagging is the process of identifying jets containing b -hadrons, c -hadrons, light-hadrons (uds -hadrons) or jets from hadronically decaying τ leptons. Distinguishing b -jets is of particular interest due to their characteristically long lifetime ($\tau \approx 1.5$ ps), displaced secondary decay vertex and high decay multiplicity.

Usage of b -tagging in this analysis is done via five operating points (OPs), corresponding

831 to 65%, 70%, 77%, 85% and 90% b -jet tagging efficiency ε_b in simulated $t\bar{t}$ events, in order
832 from the loosest to tightest discriminant cut point. The OPs are defined by placing selections
833 on the tagger output to provide a predefined ε_b level; the selection cuts act as a variable
834 trade-off between b -tagging efficiency and b -jet purity i.e. c - or light-jet rejection. For this
835 thesis, a jet is considered b -tagged if it passes the 85% OP. The b -tagged jet is then assigned a
836 pseudo-continuous b -tagging (PCBT) score, which quantifies a jet's ability to satisfy different
837 OPs. The score can take integer values between 1 and 6, where a score of 6 is assigned to
838 jets passing all OP thresholds; a score of 2 for jets that pass only the tightest OP (90%);
839 and a score of 1 for jets that pass no OP. A value of -1 is also defined for any jet that does
840 not satisfy b -tagging criteria.

841 GN2 b -tagging algorithm




Figure 4.3: Overview of the GN2 architecture. The number of jet and track features are represented by n_{jf} and n_{tf} respectively. The global jet representation and track embeddings output by the Transformer encoder are used as inputs for three task-specific networks [62].

842 For this analysis, b -jets are identified and tagged with the GN2v01 b -tagger [62]. The

GN2 algorithm uses a Transformer-based model [63] modified to incorporate domain knowledge and additional auxiliary physics objectives: grouping tracks with a common vertex and predicting the underlying physics process for a track. The network structure is shown in Figure 4.3. The GN2 b -tagger form the input vector by concatenating 2 jet variables and 19 track reconstruction variables (for up to 40 tracks), normalized to zero mean and unit variance. The output consists of a track-pairing output layer of size 2, a track origin classification layer of 7 categories, and a jet classification layer of size 4 for the probability of each jet being a b -, c -, light- or τ -jet respectively. For b -tagging purpose, a discriminant is defined using these four outputs

$$D_b = \ln \left(\frac{p_b}{f_c p_c + f_\tau p_\tau + (1 - f_c - f_\tau) p_{\text{light}}} \right) \quad (4.1)$$

where p_x is the probability of the jet being an x -jet as predicted by GN2, and f_c , f_τ are tunable free parameters controlling balance between c - and light-jet rejection.

Simulated SM $t\bar{t}$ and BSM Z' events from pp collisions were used as training and evaluation samples. In order to minimize bias, both b - and light-jet samples are re-sampled to match c -jet distributions. Figure 4.4 shows the performance of GN2 compared to the previous convolutional neural network-based standard b -tagging algorithm DL1d, in terms of c -, light- and τ -jet rejection as a function of b -tagging efficiency. The network gives a factor of 1.5-4 improvement in experimental applications compared to DL1d [62], without dependence on the choice of MC event generator or inputs from low-level flavor tagging algorithm.




Figure 4.4: The c -, light- and τ -jet rejection rate as a function of b -tagging efficiency for GN2 and DL1d using (a) jets in the $t\bar{t}$ sample, and (b) jets in the Z' sample. The performance ratios of GN2 to DL1d are shown in the bottom panels [62].

Efficiency calibration

Due to imperfect description of detector response and physics modeling effects in simulation, the b -tagging efficiency predicted by MC simulation $\varepsilon_b^{\text{sim}}$ requires a correction factor to match the efficiency measured in collision data $\varepsilon_b^{\text{data}}$. The correction scale factors (SFs) are defined as $\text{SF} = \varepsilon_b^{\text{data}} / \varepsilon_b^{\text{sim}}$ and are determined by data-to-MC calibration using samples enriched in dileptonic $t\bar{t}$ decays [64]. The resulting SFs are applied to MC simulated jets individually.

4.3 Leptons

Lepton reconstruction in ATLAS involves electron and muon reconstruction since tau decays quickly, and depending on decay mode can be reconstructed using either jets or light leptons. Leptons can be classified into two categories: prompt leptons resulting from heavy particle decays and non-prompt leptons resulting from detector or reconstruction effects, or from heavy-flavor hadron decays.

4.3.1 Electrons

Electrons leave energy signature in the detector by interacting with the detector materials and losing energy in the form of bremsstrahlung photons. A bremsstrahlung photon can produce an electron-positron pair which can itself deposit signals in the detector, creating a cascade of particles that can leave multiple of either tracks in the ID or EM showers in the calorimeters, all of which are considered part of the same EM topo-cluster. Electron signal signature has three characteristic components: localized energy deposits in the calorimeters, multiple tracks in the ID and compatibility between the above tracks and energy clusters in the $\eta \times \phi$ plane [45]. Electron reconstruction in ATLAS follows these steps accordingly.

Seed-cluster reconstruction and track reconstruction are performed sequentially in accordance with the iterative topo-clustering algorithm and track reconstruction method described in section 4.1. The seed-cluster and GSF-refitted track candidate not associated with a conversion vertex are matched to form an electron candidate. The cluster energy is then calibrated using multivariate techniques on data and simulation to match the original electron energy.

Electron identification

Additional LH-based identification selections using ID and EM calorimeter information are implemented to further improve the purity of reconstructed electrons in the central region of the detector ($|\eta| < 2.47$) [45]. The electron LH function is built with the signal being prompt electrons and background being objects with similar signature to prompt electrons i.e. hadronic jet deposits, photon conversions or heavy-flavor hadron decays. Three identification OPs are defined for physics analyses: *Loose*, *Medium* and *Tight*, optimized for 9 bins in $|\eta|$ and 12 bins in E_T with each OP corresponding to a fixed efficiency requirement for each bin. For typical EW processes, the target efficiencies for *Loose*, *Medium* and *Tight* start at 93%, 88% and 80% respectively and increase with E_T . Similar to b -tagging OPs, the electron identification OPs represent a trade-off in signal efficiency and background rejection. The electron efficiency are estimated using tag-and-probe method on samples of $J/\Psi \rightarrow ee$ and $Z \rightarrow ee$ [45].

Electron isolation

A characteristic distinction between prompt electrons and electrons from background processes is the relative lack of activity in both the ID and calorimeters within an $\Delta\eta \times \Delta\phi$ area surrounding the reconstruction candidate. Calorimeter-based and track-based electron isolation variables [45] are defined to quantify the amount of activity around the electron candidate using topo-clusters and reconstructed tracks respectively.

Calorimeter-based isolation variables $E_T^{\text{cone}XX}$ are computed by first summing the energy of topo-clusters with barycenters falling within a cone of radius $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = XX/100$ around the direction of the electron candidate. The final isolation variables are

911 obtained by subtracting from the sum the energy belonging to the candidate electron at the
 912 core of the cone, then applying corrections for pile-up effects and energy leakage outside of
 913 the core. Similar to calorimeter-based variables, track-based isolation variables $p_{\text{T}}^{\text{varconeXX}}$
 914 are calculated by summing all track p_{T} within a cone of radius ΔR around the electron
 915 candidate, minus the candidate's contribution. The cone radius is variable as a function of
 916 p_{T} and is described as

$$\Delta R \equiv \min \left(\frac{10}{p_{\text{T}}}, \Delta R_{\text{max}} \right), \quad (4.2)$$

917 where p_{T} is expressed in GeV and ΔR_{max} is the maximum cone size, defined to account for
 918 closer proximity of decay products to the electron in high-momentum heavy particle decays.
 919 Four isolation operating points are implemented to satisfy specific needs by physics analyses:
 920 *Loose*, *Tight*, *HighPtCaloOnly* and *Gradient* [45].

921 **Electron charge misidentification**

922 Charge misidentification is a crucial irreducible background, particularly for analyses
 923 with electron charge selection criteria. Electron charge is determined by the curvature of
 924 the associated reconstructed track, and misidentification of charge can occur via either an
 925 incorrect curvature measurement or an incorrectly matched track. Inaccurate measurement
 926 is more likely for high energy electrons due to the small curvature in track trajectories
 927 at high p_{T} , while track matching error usually results from bremsstrahlung pair-production
 928 generating secondary tracks in close proximity [45]. Suppression of this background is assisted
 929 via a boosted decision tree discriminant named the Electron Charge ID Selector (ECIDS) [65].
 930 The addition of ECIDS removed 90% of electrons with incorrect charge while selecting 98%

of electrons with correct charge from electrons in $Z \rightarrow ee$ events satisfying *Medium/Tight* identification and *Tight* isolation criteria.

4.3.2 Muons

Muons act as minimum-ionizing particles, leaving tracks in the MS or characteristics energy deposits in the calorimeter and can be reconstructed globally using information from the ID, MS and calorimeters. Five reconstruction strategies corresponding to five muon types [66] are utilized in ATLAS:

- Combined (CB): the primary ATLAS muon reconstruction method. Combined muons are first reconstructed using MS tracks then extrapolated to include ID tracks (outside-in strategy). A global combined track fit is performed on both MS and ID tracks.
- Inside-out combined (IO): complementary to CB reconstruction. IO muon tracks are extrapolated from ID to MS, then fitted with MS hits and calorimeter energy loss in a combined track fit.
- MS extrapolated (ME): ME muons are defined as muons with a MS track that cannot be matched to an ID track using CB reconstruction. ME muons allow extension of muon reconstruction acceptance to regions not covered by the ID ($2.5 < |\eta| < 2.7$)
- Segment-tagged (ST): ST muons are defined as a successfully matched ID track that satisfies tight angular matching criteria to at least one reconstructed MDT or CSC segment when extrapolated to the MS. MS reconstruction is used primarily when muons only crossed one layer of MS chambers.
- Calorimeter-tagged (CT): CT muons are defined as an ID track that can be matched to

energy deposits consistent with those of a minimum-ionizing particle when extrapolated through the calorimeter. CT reconstruction extends acceptance range to regions in the MS with sparse instrumentation ($|\eta| < 0.1$) with a higher p_T threshold of 5 GeV, compared to the 2 GeV threshold used by other muon reconstruction algorithms due to large background contamination at the low p_T range of $15 < p_T < 100$ GeV [67].

Muon identification

Reconstructed muons are further filtered by identification criteria to select for high-quality prompt muons. Requirements include number of hits in the MS and ID, track fit properties and compatibility between measurements of the two systems. Three standard OPs (*Loose*, *Medium*, *Tight*) are defined to better match the needs of different physics analyses concerning prompt muon p_T resolution, identification efficiency and non-prompt muon rejection. The default identification OP for ATLAS physics is *Medium* which provides efficiency and purity suitable for a wide range of analyses while minimizing systematic uncertainties [66].

Muon isolation

Muons from heavy particle decays are often produced in an isolated manner compared to muons from semileptonic decays, and is therefore an important tool for background rejection in many physics analyses. Muon isolation strategies are similar to that of electron in section 4.3.1, with track-based and calorimeter-based isolation variables. Seven isolation OPs are defined using either or both types of isolation variables [66].

4.4 Missing transverse momentum

Collisions at the LHC happen along the z -axis of the ATLAS coordination system between two particle beam of equal center-of-mass energy. By conservation of momentum, the sum of transverse momenta of outgoing particles should be zero. A discrepancy between measured momentum and zero would then suggest the presence of undetectable particles, which would consist of either SM neutrinos or some unknown BSM particles, making missing transverse momentum (E_T^{miss}) an important observable to reconstruct. Reconstructing E_T^{miss} utilizes information from fully reconstructed leptons, photons, jets and other matched track-vertex objects not associated with a prompt object (soft signals), defined with respect to the $x(y)$ -axis as

$$E_{x(y)}^{\text{miss}} = - \sum_{i \in \{\text{hard objects}\}} p_{x(y),i} - \sum_{j \in \{\text{soft signals}\}} p_{x(y),j}, \quad (4.3)$$

where $p_{x(y)}$ is the $x(y)$ -component of p_T for each particle [68]. The following observables can then be defined:

$$\begin{aligned} \mathbf{E}_T^{\text{miss}} &= (E_x^{\text{miss}}, E_y^{\text{miss}}), \\ E_T^{\text{miss}} &= |\mathbf{E}_T^{\text{miss}}| = \sqrt{(E_x^{\text{miss}})^2 + (E_y^{\text{miss}})^2}, \\ \phi^{\text{miss}} &= \tan^{-1}(E_y^{\text{miss}}/E_x^{\text{miss}}), \end{aligned} \quad (4.4)$$

984 where E_T^{miss} represents the magnitude of the missing transverse energy vector $\mathbf{E}_T^{\text{miss}}$, and
 985 ϕ^{miss} its direction in the transverse plane. The vectorial sum $\mathbf{E}_T^{\text{miss}}$ can be broken down into

$$\mathbf{E}_T^{\text{miss}} = - \underbrace{\sum_{\text{selected electrons}} \mathbf{p}_T^e - \sum_{\text{selected muons}} \mathbf{p}_T^\mu - \sum_{\text{accepted photons}} \mathbf{p}_T^\gamma - \sum_{\text{accepted } \tau\text{-leptons}} \mathbf{p}_T^\tau - \sum_{\text{accepted jets}} \mathbf{p}_T^{\text{jet}}}_{\text{hard term}} - \underbrace{\sum_{\text{unused tracks}} \mathbf{p}_T^{\text{track}}}_{\text{soft term}}. \quad (4.5)$$

986 Two OPs are defined for E_T^{miss} , *Loose* and *Tight*, with selections on jet p_T and JVT criteria
 987 [69]. The *Tight* OP is used in this analysis; *Tight* reduces pile-up dependence of E_T^{miss}
 988 by removing the phase space region containing more pile-up than hard-scatter jets, at the
 989 expense of resolution and scale at low pile-up,

990 4.5 Overlap removal

991 Since different objects are reconstructed independently, it is possible for the same de-
 992 tector signals to be used to reconstruct multiple objects. An overlap removal strategy is
 993 implemented to resolve ambiguities; the overlap removal process for this analysis applies
 994 selections in Table 4.1 sequentially, from top to bottom.

Table 4.1: Overlap removal process for this analysis, applied sequentially from top to bottom.

Remove	Keep	Matching criteria
Electron	Electron	Shared ID track, $p_{T,1}^e < p_{T,2}^e$
Muon	Electron	Shared ID track, CT muon
Electron	Muon	Shared ID track
Jet	Electron	$\Delta R < 0.2$
Electron	Jet	$\Delta R < 0.4$
Jet	Muon	$(\Delta R < 0.2 \text{ or ghost-associated}) \ \& \ N_{\text{track}} < 3$
Muon	Jet	$\Delta R < \min(0.4, 0.04 + 10\text{GeV}/p_T^\mu)$

Chapter 5. Data & Simulated Samples

5.1 Data samples

Data samples used in this analysis were collected by the ATLAS detector during Run 2 data-taking campaign between 2015-2018. The samples contain pp collisions at center-of-mass energy of $\sqrt{s} = 13$ TeV with 25 ns bunch-spacing, which corresponds to an integrated luminosity of 140 fb^{-1} with an uncertainty of 0.83% [70]. The HLT trigger strategy is similar to that of previous $t\bar{t}t\bar{t}$ observation analysis [36] and include single lepton and dilepton triggers. Calibration for di-muon and electron-muon triggers were not ready for the samples used in this analysis, and are therefore not included. Events are also required to contain at least one lepton matched to the corresponding object firing the trigger. Triggers used are summarized in Table 5.1.

5.2 Monte Carlo samples

Monte Carlo simulated samples are used to estimate signal acceptance before unblinding, profile the physics background for the analysis and to study object optimizations. Simulated samples for this analysis use are generated from ATLAS generalized MC20a/d/e samples for Run 2, using full detector simulation (FS) and fast simulation (AF3) to simulate detector response. MC samples used and simulation processes are summarized in Table 5.2

Table 5.1: Summary of all HLT triggers used in this analysis. Events are required to pass at least one trigger.

Trigger	Data period			
	2015	2016	2017	2018
Single electron triggers				
HLT_e24_lhmedium_L1EM20VH	✓	-	-	-
HLT_e60_lhmedium	✓	-	-	-
HLT_e120_lhloose	✓	-	-	-
HLT_e26_lhtight_nod0_ivarloose	-	✓	✓	✓
HLT_e60_lhmedium_nod0	-	✓	✓	✓
HLT_e140_lhloose_nod0	-	✓	✓	✓
Di-electron triggers				
HLT_2e12_lhloose_L12EM10VH	✓	-	-	-
HLT_2e17_lhvloose_nod0	-	✓	-	-
HLT_2e24_lhvloose_nod0	-	-	✓	✓
HLT_2e17_lhvloose_nod0_L12EM15VHI	-	-	-	✓
Single muon trigger				
HLT_mu20_iloose_L1MU15	✓	-	-	-
HLT_mu40	✓	-	-	-
HLT_mu26_ivarmedium	-	✓	✓	✓
HLT_mu50	-	✓	✓	✓

5.2.1 $t\bar{t}Z'$ signal samples

Signal $t\bar{t}Z'$ samples were generated based on the simplified top-philic resonance model in section 2.2.1 where a color singlet vector resonance couples strongly to only top and antitop. Six Z' mass points were utilized for the generation of the signal sample: 1000, 1250, 1500, 2000, 2500 and 3000 GeV. The top- Z' coupling c_t is chosen to be 1 for a narrow resonance peak, and the chirality angle θ is chosen to be $\pi/4$ to suppress loop production of Z' . The samples were then generated with MADGRAPH5_AMC@NLO v.3.5.0 [71] at LO with the NNPDF3.1L0 [72] PDF set interfaced with PYTHIA8 [73] using A14 tune and NNPDF2.31o PDF set for parton showering and hadronization. The resonance width is calculated to be

Table 5.2: Summary of all Monte-Carlo samples used in this analysis. V refers to an EW ($W^\pm/Z/\gamma^*$) or Higgs boson. Matrix element (ME) order refers to the order in QCD of the perturbative calculation. Tune refers to the underlying-event tune of the parton shower (PS) generator.

Process	ME Generator	ME Order	ME PDF	PS	Tune	Sim.
Signals						
$t\bar{t}Z'$	MADGRAPH5_AMC@NLO	LO	NNPDF3.1LO	PYTHIA8	A14	FS
$t\bar{t}t\bar{t}$ and $t\bar{t}t$						
$t\bar{t}t\bar{t}$	MADGRAPH5_AMC@NLO	NLO	NNPDF3.0nlo	PYTHIA8	A14	AF3
	MADGRAPH5_AMC@NLO	NLO	MMHT2014 LO	HERWIG7	H7-UE-MMHT	AF3
	SHERPA	NLO	NNPDF3.0nnlo	HERWIG7	SHERPA	FS
	MADGRAPH5_AMC@NLO	LO	NNPDF2.3lo	PYTHIA8	A14	AF3
$t\bar{t}V$						
$t\bar{t}H$	POWHEGBOX v2	NLO	NNPDF3.0nlo	PYTHIA8	A14	FS
	POWHEGBOX v2	NLO	NNPDF3.0nlo	HERWIG7	H7.2-Default	FS
$t\bar{t}(Z/\gamma^*)$	MADGRAPH5_AMC@NLO	NLO	NNPDF3.0nlo	PYTHIA8	A14	FS
	SHERPA	NLO	NNPDF3.0nnlo	SHERPA	SHERPA	FS
$t\bar{t}W$	SHERPA	NLO	NNPDF3.0nnlo	SHERPA	SHERPA	FS
	SHERPA	LO	NNPDF3.0nnlo	SHERPA	SHERPA	FS
$t\bar{t}$ and Single-Top						
$t\bar{t}$	POWHEGBOX v2	NLO	NNPDF3.0nlo	PYTHIA8	A14	FS
$t\bar{t}W$	POWHEGBOX v2	NLO	NNPDF3.0nlo	PYTHIA8	A14	FS
$t(q)b$	POWHEGBOX v2	NLO	NNPDF3.0nlo (s)	PYTHIA8	A14	FS
			NNPDF3.0nlo 4f (t)			FS
tWZ	MADGRAPH5_AMC@NLO	NLO	NNPDF3.0nlo	PYTHIA8	A14	FS
tZ	MADGRAPH5_AMC@NLO	LO	NNPDF3.0nlo 4f	PYTHIA8	A14	FS
$t\bar{t}VV$						
$t\bar{t}WW$	MADGRAPH5_AMC@NLO	LO	NNPDF3.0nlo	PYTHIA8	A14	FS
$t\bar{t}WZ$	MADGRAPH	LO	NNPDF3.0nlo	PYTHIA8	A14	AF3
$t\bar{t}HH$	MADGRAPH	LO	NNPDF3.0nlo	PYTHIA8	A14	AF3
$t\bar{t}WH$	MADGRAPH	LO	NNPDF3.0nlo	PYTHIA8	A14	AF3
$t\bar{t}ZZ$	MADGRAPH	LO	NNPDF3.0nlo	PYTHIA8	A14	AF3
$V(VV)+\text{jets}$ and VH						
$V+\text{jets}$	SHERPA	NLO	NNPDF3.0nnlo	SHERPA	SHERPA	FS
$VV+\text{jets}$	SHERPA	NLO	NNPDF3.0nnlo	SHERPA	SHERPA	FS
		LO ($gg \rightarrow VV$)				FS
$VVV+\text{jets}$	SHERPA	NLO	NNPDF3.0nnlo	SHERPA	SHERPA	FS
VH	POWHEGBOX v2	NLO	NNPDF3.0aznlo	PYTHIA8	A14	FS

1021 4% for $c_t = 1$.

1022 5.2.2 Background samples

1023 SM $t\bar{t}t\bar{t}$ background

1024 Nominal SM $t\bar{t}t\bar{t}$ sample was generated with MADGRAPH5_AMC@NLO [71] at NLO
1025 in QCD with the NNPDF3.0nlo [72] PDF set and interfaced with PYTHIA8.230 [73] using
1026 A14 tune [74]. Decays for top quarks are simulated LO with MADSPIN [75, 76] to preserve
1027 spin information, while decays for b - and c -hadrons are simulated with EVTGEN v1.6.0
1028 [77]. The renormalization and factorization scales μ_R and μ_F are set to $\sqrt{m^2 + p_T^2}/4$, which
1029 represents the sum of transverse mass of all particles generated from the ME calculation [78].
1030 The ATLAS detector response was simulated with AF3. Additional auxiliary $t\bar{t}t\bar{t}$ samples
1031 are also generated to evaluate the impact of generator and PS uncertainties as shown in 5.2.

1032 $t\bar{t}W$ background

1033 Nominal $t\bar{t}W$ sample was generated using SHERPA v2.2.10 [79] at NLO in QCD with
1034 the NNPDF3.0nnlo [72] PDF with up to one extra parton at NLO and two at LO, which
1035 are matched and merged with SHERPA PS based on Catani-Seymour dipole factorization
1036 [80] using the MEPS@NLO prescription [81–84] and a merging scale of 30 GeV. Higher-
1037 order ME corrections are provided in QCD by the OpenLoops 2 library [85–87] and in EW
1038 from $\mathcal{O}(\alpha^3) + \mathcal{O}(\alpha_S^2\alpha^2)$ (LO3 & NLO2) via two sets of internal event weights. An alternative
1039 sample with only EW corrections at LO from $\mathcal{O}(\alpha_S\alpha^3)$ (NLO3) diagrams were also simulated
1040 with the same settings.

1041 $t\bar{t}(Z/\gamma^*)$ background

1042 Nominal $t\bar{t}(Z/\gamma^*)$ samples were generated separately for different ranges of dilepton in-
1043 variant mass $m_{\ell\ell}$ to account for on-shell and off-shell Z/γ^* production. Sample for $m_{\ell\ell}$
1044 between 1 and 5 GeV was produced using MADGRAPH5_AMC@NLO [71] at NLO with
1045 the NNPDF3.0nnlo [72] PDF set, interfaced with PYTHIA8.230 [73] using A14 tune [74] and
1046 NNPDF2.31lo PDF set. Sample for $m_{\ell\ell} < 5$ GeV was produced with SHERPA v2.2.10 [79]
1047 at NLO using NNPDF3.0nnlo PDF set. To account for generator uncertainty, an alternative
1048 $m_{\ell\ell} > 5$ GeV sample was generated with identical settings to the low $m_{\ell\ell}$ sample. The
1049 ATLAS detector response was simulated with full detector simulation (FS).

Chapter 6. Analysis Strategy

6.1 Event selection

Events for the analysis first are preselected following a list of criteria to optimize for event quality and background rejection. The following criteria are applied sequentially from top to bottom along with cleaning and veto cuts

1. **Good Run List (GRL):** data events must be part of a predefined list of suitable runs and luminosity blocks [88].
2. **Primary vertex:** events must have at least one reconstructed vertex matched to 2 or more associated tracks with $p_T > 500$ MeV.
3. **Trigger:** events must be selected by at least one trigger in Table 5.1.
4. **Kinematic selection:** events must have exactly two *Tight* leptons with the same electric charge, or at least three *Tight* leptons of any charge. The leading lepton must have $p_T > 28$ GeV, and all leptons must satisfy $p_T > 15$ GeV.

Events are separated into two channels based on the number of leptons: same-sign dilepton (SS2L) for events with exactly two leptons of the same charge, or multilepton (ML) for events with three or more leptons. The channels are further separated into regions defined in section 6.2 to prepare for analysis.

Additional selections are applied based on the lepton flavors present. In the SS2L channel, if both leptons are electrons, the invariant mass m_{ll} must satisfy $m_{ll} < 81$ GeV and $m_{ll} > 101$ GeV to suppress background involving Z -bosons. In the ML channel, the same criteria must be satisfied for every opposite-sign same-flavor pair of leptons in an event.

6.1.1 Object definition

Table 6.1 shows the selections used in this analysis. Each selection comes with associated calibration scale factors (SFs) to account for discrepancies between data and MC simulation, and are applied multiplicatively to MC event weights.

Table 6.1: Summary of object selection criteria used in this analysis.

Selection	Electrons	Muons	Jets
p_T [GeV]	> 15 $p_T(l_0) > 28$	> 15	> 20
$ \eta $	$1.52 \leq \eta < 2.47$ < 1.37	< 2.5	< 2.5
Identification	<i>TightLH</i> pass ECIDS ($ee/e\mu$)	<i>Medium</i>	NNJvt <i>FixedEffPt</i> ($p_T < 60$, $ \eta < 2.4$)
Isolation	<i>Tight_VarRad</i>	<i>PflowTight_VarRad</i>	
Track-vertex assoc.			
$ d_0^{\text{BL}}(\sigma) $	< 5	< 3	
$ \Delta z_0^{\text{BL}} \sin \theta $ [mm]	< 0.5	< 0.5	

6.1.2 Event categorization

Simulated events are categorized using truth information of leptons (e/μ) and their originating MC particle (mother-particle). Each lepton can be classified as either prompt or non-prompt, with non-prompt leptons further categorized for background estimation purposes. If an event contains only prompt leptons, the event is classified as its corresponding process. If the event contains one non-prompt lepton, the event is classified as the corresponding type of the non-prompt lepton. If the event contains more than one non-prompt lepton, the event is classified as other.

- **Prompt:** if the lepton originates from $W/Z/H$ boson decays, or from a mother-particle created by a final state photon.

- **Non-prompt:**

- **Charge-flip (e only):** if the reconstructed charge of the lepton differs from that of the first mother-particle.

- **Material conversion (e only):** if the lepton originated from a photon conversion and the mother-particle is an isolated prompt photon, non-isolated final state photon, or heavy boson.

- **γ^* -conversion (e only):** if the lepton originated from a photon conversion and the mother-particle is a background electron.

- **Heavy flavor decay:** if the lepton originated from a b - or c -hadron.

- **Fake:** if the lepton originated from a light- or s -hadron, or if the truth type of the lepton is hadron.

- **Other:** any lepton that does not belong to one of the above categories.

6.2 Analysis regions

Events are selected and categorized into analysis regions belonging to one of two types: control regions (CRs) enriched in background events, and signal regions (SRs) enriched in signal events. This allows for the examination and control of backgrounds and systematic uncertainties, as well as study of signal sensitivities. The signal is then extracted from the SRs with a profile LH fit using all regions. The full selection criteria for each region are summarized in Table 6.2.

Table 6.2: Definitions of signal, control and validation regions (VR) used in this analysis. N_{jets} and N_b refers to the number of jets and number of b -tagged jets respectively. ℓ_1 refers to the leading lepton, ℓ_2 refers to the subleading lepton and so on. H_T refers to the p_T scalar sum of all leptons and jets in the event. $m_{\ell\ell}$ refers to the dilepton invariant mass, which must not coincide with the Z -boson mass range of 81-101 GeV for SS2L+3L events.

Region	Channel	N_{jets}	N_b	Other selections	Fitted variable
CR Low m_{γ^*}	SS $e\ell$	[4, 6)	≥ 1	ℓ_1/ℓ_2 is from virtual photon decay $\ell_1 + \ell_2$ not from material conversion	event yield
CR Mat. Conv.	SS $e\ell$	[4, 6)	≥ 1	ℓ_1/ℓ_2 is from material conversion	event yield
CR HF μ	$\ell\mu\mu$	≥ 1	1	$\ell_1 + \ell_2$ not conversion candidates $100 < H_T < 300$ GeV $E_T^{\text{miss}} > 35$ GeV total charge = ± 1	$p_T(\ell_3)$
CR HF e	$e\ell\ell$	≥ 1	1	$\ell_1 + \ell_2$ not conversion candidates $100 < H_T < 275$ GeV $E_T^{\text{miss}} > 35$ GeV total charge = ± 1	$p_T(\ell_3)$
CR $t\bar{t}W^+$	SS $\ell\mu$	≥ 4	≥ 2	$ \eta(e) < 1.5$ for $N_b = 2$: $H_T < 500$ GeV or $N_{\text{jets}} < 6$ for $N_b \geq 3$: $H_T < 500$ GeV total charge > 0	N_{jets}
CR $t\bar{t}W^-$	SS $\ell\mu$	≥ 4	≥ 2	$ \eta(e) < 1.5$ for $N_b = 2$: $H_T < 500$ GeV or $N_{\text{jets}} < 6$ for $N_b \geq 3$: $H_T < 500$ GeV total charge < 0	N_{jets}
CR 1b(+)	SS2L+3L	≥ 4	1	$\ell_1 + \ell_2$ not from material conversion $H_T > 500$ GeV total charge > 0	N_{jets}
CR 1b(-)	SS2L+3L	≥ 4	1	$\ell_1 + \ell_2$ not from material conversion $H_T > 500$ GeV total charge < 0	N_{jets}
VR $t\bar{t}Z$	3L $\ell^\pm\ell^\mp$	≥ 4	≥ 2	$m_{\ell\ell} \in [81, 101]$ GeV	$N_{\text{jets}}, m_{\ell\ell}$
VR $t\bar{t}W + 1b$	SS2L+3L			CR $t\bar{t}W^\pm$ CR 1b(\pm)	N_{jets}
VR $t\bar{t}W + 1b + \text{SR}$	SS2L+3L			CR $t\bar{t}W^\pm$ CR 1b(\pm) SR	N_{jets}
SR	SS2L+3L	≥ 6	≥ 2	$H_T > 500$ GeV $m_{\ell\ell} \notin [81, 101]$ GeV	H_T

6.2.1 Signal regions

All events selected for the SR must satisfy the following criteria:

- Contains 6 or more jets, with at least 2 jets b -tagged at the 85% OP.
- Scalar sum of the transverse momenta of all leptons and jets $H_T > 500$ GeV.
- Dilepton invariant mass $m_{\ell\ell}$ does not coincide with the Z -boson mass range of $81 - 101$ GeV

The SR is further divided into sub-regions by the number of b -jets and leptons as shown in Table 6.3 to further study signal behavior and improve sensitivity.

Table 6.3: Definitions of SR sub-regions. Events are sorted into different sub-regions based on the number of b -tagged jets and leptons present.

Sub-region	Selection criteria	
	b -jets	leptons
SR 2b2l	$N_b = 2$	$N_l = 2$
SR 2b3l4l	$N_b = 2$	$N_l \geq 3$
SR 3b2l	$N_b = 3$	$N_l = 2$
SR 3b3l4l	$N_b = 3$	$N_l \geq 3$
SR 4b	$N_b \geq 4$	

6.2.2 Control regions

Control regions are defined for each background to be enriched in the targeted process, in order to maximize the background's purity and minimize contamination from other sources within the region. This helps to constrain and reduce correlation between background normalization factors in the final fit. Fit variables and selection criteria are determined via

optimization studies performed on CRs that aimed to achieve the largest discriminating power possible between the target background and other event types.

$t\bar{t}W$ background CRs

Theoretical modeling for $t\bar{t}W$ +jets background in the phase space of this analysis suffers from large uncertainties, especially at high jet multiplicities [89]. A data-driven method was employed in a similar manner to the SM $t\bar{t}t\bar{t}$ observation analysis [36] to mitigate this effect, and are described in further details in section 6.3.3. The method necessitates the definition of two groups of dedicated CRs to estimate the flavor composition and normalization of $t\bar{t}W$ +jets background: CR $t\bar{t}W$ +jets to constrain flavor composition, and CR 1b to constrain the jet multiplicity spectrum. These are further split into CR $t\bar{t}W^\pm$ and CR 1b(\pm) due to the pronounced asymmetry in $t\bar{t}W$ production from pp collisions, with $t\bar{t}W^+$ being produced at approximately twice the rate of $t\bar{t}W^-$ [90].

Events in CR $t\bar{t}W^\pm$ are required to contain at least two b -tagged jets similar to the SR to determine the $t\bar{t}W$ normalization within an SR-related phase space. Orthogonality with SR is ensured by requiring $H_T < 500$ GeV or $N_{\text{jets}} < 6$ when $N_b = 2$, and $H_T < 500$ GeV when $N_b \geq 3$. Events in CR 1b(\pm) are required to have $H_T > 500$ GeV and at least four jets to encompass events with high N_{jets} , which can be used to determine the $t\bar{t}W$ jet multiplicity spectrum for fitting $a_{0,1}$. The selection criteria also include exactly one b -tagged jet to maintain orthogonality with the SR.

Fake/non-prompt background CRs

Selection for fake/non-prompt CRs are determined using the `DFCommonAddAmbiguity` (DFCAA) variable for reconstructed leptons.

Table 6.4: List of possible assigned values for DFCAA.

DFCAA	Description
-1	No 2nd track found
0	2nd track found, no conversion found
1	Virtual photon conversion candidate
2	Material conversion candidate

Four CRs are defined for the three main types of fake/non-prompt backgrounds in the analysis - virtual photon (γ^*) conversion, photon conversion in detector material (Mat. Conv.) and heavy flavor decays (HF). The full selection criteria for fake/non-prompt CRs are shown in Table 6.2.

- **Low m_γ^* :** events with an e^+e^- pair produced from a virtual photon.

Events are selected if there are two same-sign leptons with at least one electron reconstructed as an internal conversion candidate, and neither reconstructed as a material conversion candidate.

- **Mat. Conv.:** events with an electron originating from photon conversion within the detector material.

Events are selected if there are two same-sign leptons with at least one electron reconstructed as a material conversion candidate.

- **HF $e(\mu)$:** events with a reconstructed non-prompt lepton from semi-leptonic decays of b - and c -hadrons (heavy flavor decays).

Events are selected if there are three leptons with at least two electrons (muons), with no lepton reconstructed as a conversion candidate.

6.3 Background estimation

Background in this analysis consist of SM processes that can result in a signal signature similar to a $t\bar{t}t\bar{t}$ SSML final state and can be divided into two types, reducible and irreducible. Reducible background consists of processes that do not result in a SSML final state physically, but are reconstructed as such due to detector and reconstruction effects. Three main types of reducible background are considered: charge misidentification (QmisID) and fake/non-prompt leptons. Fake/non-prompt lepton backgrounds are estimated using template fitting method, where MC simulations are normalized to their theoretical SM cross section via floating normalization factors (NFs) constrained by the corresponding CRs. Lepton charge misidentification background contaminates the SR with opposite-sign events, and are estimated using a data-driven method described in section 6.3.2 along with ECIDS described in section 4.3.1.

Irreducible background consists of SM processes that result in SSML final states physically with all leptons being prompt. The dominating background in the SR are SM $t\bar{t}t\bar{t}$, $t\bar{t}W$, $t\bar{t}Z$, and $t\bar{t}H$ production with smaller contributions from VV , VVV , VH and rarer processes like $t\bar{t}VV$, tWZ , tZq and $t\bar{t}t$. Most irreducible backgrounds are estimated using template fitting method, with the exception of $t\bar{t}W$ +jets background. The $t\bar{t}W$ +jets background is instead given four dedicated CRs, and estimated using a data-driven method with a fitted function parameterized in N_{jets} . All CRs and SR are included in the final profile LH fit to data.

6.3.1 Template fitting for fake/non-prompt estimation

Template fitting method is a semi-data-driven approach [89] that estimates fake/non-prompt background distributions by fitting the MC kinematic profile of background processes arising from fake/non-prompt leptons to data. Each of the four main sources of fake/non-prompt leptons is assigned a free-floating NF constrained by a CR enriched with the corresponding background resulting in four NFs: $\text{NF}_{\text{HF } e}$, $\text{NF}_{\text{HF } \mu}$, $\text{NF}_{\text{Mat. Conv.}}$, $\text{NF}_{\text{Low } m_{\gamma^*}}$. The NFs are fitted simultaneously with the signal.

6.3.2 Charge misidentification data-driven estimation

The ee and $e\mu$ channels in the SS2L region are contaminated with opposite-sign (OS) dilepton events with one misidentified charge. Charge misidentification (QmisID) largely affects electrons due to muons' precise curvature information using ID and MS measurements and low bremsstrahlung rate. The charge flip rates are significant at higher p_T and varies with $|\eta|$ which is proportional to the amount of detector material the electron interacted with.

The charge flip probability ϵ is estimated in this analysis with a data-driven method [91] using a sample of $Z \rightarrow e^+e^-$ events with additional constraints on the invariant mass m_{ee} to be within 10 GeV of the Z -boson mass. The Z -boson mass window is defined to be within 4σ to include most events within the peak, and is determined by fitting the m_{ee} spectrum of the two leading electrons to a Breit-Wigner function, resulting in a range of [65.57, 113.49] for SS events and [71.81, 109.89] for OS events. Background contamination near the peak is assumed to be uniform and subtracted using a sideband method. Since the Z -boson decay products consist of a pair of opposite-sign electrons, all same-sign electron

1197 pairs are considered affected by charge misidentification.

1198 Let N_{ij}^{SS} be the number of events with SS electrons with the leading electron in the
 1199 i^{th} 2D bin in $(p_{\text{T}}, |\eta|)$ and the sub-leading electron in the j^{th} bin. Assuming the QmisID
 1200 probabilities of electrons in an event are uncorrelated, N_{ij}^{SS} can be estimated as

$$N_{ij}^{\text{SS}} = N_{ij}^{\text{tot}}(\epsilon_i(1 - \epsilon_j) + \epsilon_j(1 - \epsilon_i)), \quad (6.1)$$

1201 where N_{ij}^{tot} is the total number of events in the i^{th} and j^{th} bin regardless of charge, and
 1202 $\epsilon_{i(j)}$ is the QmisID rate in the $i^{\text{th}}(j^{\text{th}})$ bin. Assuming N_{ij}^{SS} follows a Poisson distribution
 1203 around the expectation value \bar{N}_{ij}^{SS} , the $(i, n)^{\text{th}}$ rate ϵ can be estimated by minimizing a
 1204 negative-LLH function parameterized in p_{T} and $|\eta|$,

$$\begin{aligned} -\ln(\mathcal{L}(\epsilon|N_{\text{SS}})) &= -\ln \prod_{ij} \frac{(N_{ij}^{\text{tot}})^{N_{ij}^{\text{SS}}} \cdot e^{-N_{ij}^{\text{tot}}}}{N_{ij}^{\text{SS}}!} \\ &= -\sum_{ij} \left[N_{ij}^{\text{SS}} \ln(N_{ij}^{\text{tot}}(\epsilon_i(1 - \epsilon_j) + \epsilon_j(1 - \epsilon_i))) - N_{ij}^{\text{tot}}(\epsilon_i(1 - \epsilon_j) + \epsilon_j(1 - \epsilon_i)) \right]. \end{aligned} \quad (6.2)$$

1205 The QmisID rates are then calculated separately for SR and CRs with different electron
 1206 definitions i.e. CR Low m_{γ^*} , CR Mat. Conv., CR $t\bar{t}W^{\pm}$, using events from data after
 1207 applying region-specific lepton selections and ECIDS. The events are required to satisfy
 1208 SS2L kinematic selections but contains OS electrons. The following weight is applied to OS
 1209 events to correct for misidentified SS events within the region,

$$w = \frac{\epsilon_i + \epsilon_j - 2\epsilon_i\epsilon_j}{1 - \epsilon_i - \epsilon_j + 2\epsilon_i\epsilon_j}. \quad (6.3)$$

1210 The QmisID rates calculated for SR and CR $t\bar{t}W$ are shown in Figure 6.1




Figure 6.1: Charge flip rate calculated for SR and CR $t\bar{t}W$ in bins of $|\eta|$ and p_T .

1211 The QmisID rates obtained after applying w contain a dependency on jet multiplicity
 1212 and are underestimated at higher N_{jets} . This dependency affect the SR which require events
 1213 with ≥ 6 jets, and is corrected by applying a correction factor $\text{SF}_{i,n} = \epsilon_{i,n}/\epsilon_{i,N}$ where N is
 1214 the inclusive bin containing all N_{jets} and $\epsilon_{i,n}$ is the QmisID rate obtained from Equation 6.2
 1215 in the $(i, n)^{\text{th}}$ 2D bin in (p_T, N_{jets}) . Jet multiplicity and consequently the obtained SFs are
 1216 assumed to be independent of $|\eta|$.

1217 6.3.3 $t\bar{t}W$ background data-driven estimation

1218 Previously, $t\bar{t}W$ background in $t\bar{t}t\bar{t}$ final state analysis was handled by assigning large
 1219 ad-hoc systematic uncertainties to $t\bar{t}W$ events with 7 or more jets [92]. A semi-data-driven
 1220 method [93] was shown to be effective in the SM $t\bar{t}t\bar{t}$ observation analysis [36] by improving
 1221 $t\bar{t}W$ modeling, especially in the showering step and switching $t\bar{t}W$ systematic uncertainties
 1222 from predominantly modeling to statistical.

1223 The data-driven method applies correction factors obtained from a fitted function pa-
 1224 rameterized in N_{jets} to $t\bar{t}W$ MC kinematic distributions. The QCD scaling patterns [94] can
 1225 be represented by ratio of successive exclusive jet cross-sections

$$R_{(n+1)/n} = \frac{\sigma_{n+1}}{\sigma_n} = e^{-b} + \frac{\bar{n}}{n+1} = a_0 + \frac{a_1}{1+(j-4)}, \quad (6.4)$$

1226 where $a_{0(1)}$ and b are constants, n is the number of jets in addition to the hard process, j
 1227 is the inclusive number of jets, and \bar{n} is the expectation value for the Poisson distribution
 1228 of exclusive jet cross-section at jet multiplicity n . The $t\bar{t}W$ ME for SS2L events gives 4 jets
 1229 in the hard process, so n is defined starting from the 5th jets and the inclusive number of
 1230 jets $j = n + 4$. The two terms in Equation 6.4 correspond to staircase and Poisson scaling
 1231 in cross section between successive jet multiplicities and are sensitive to high and low jet
 1232 multiplicity events respectively [94]. The scaling pattern can then be reparameterized in a_0
 1233 and a_1 to obtain the $t\bar{t}W$ yield at $j' \equiv j + 1$ jets

$$\text{Yield}_{t\bar{t}W(j')} = \text{Yield}_{t\bar{t}W(N_{\text{jets}}=4)} \times \prod_{j=4}^{j'-1} \left(a_0 + \frac{a_1}{1+(j-4)} \right) \quad (6.5)$$

1234 with $j \geq 4$. The $t\bar{t}W$ yield in the 4-jet bin can be represented by a NF applied to $t\bar{t}W$ MC
 1235 simulation

$$\text{Yield}_{t\bar{t}W(N_{\text{jets}}=4)} = \text{NF}_{t\bar{t}W(N_{\text{jets}}=4)} \times \text{MC}_{t\bar{t}W(N_{\text{jets}}=4)}. \quad (6.6)$$

1236 To account for the asymmetry in $t\bar{t}W^+$ and $t\bar{t}W^-$ cross-sections, $\text{NF}_{t\bar{t}W(N_{\text{jets}}=4)}$ is further
 1237 split into $\text{NF}_{t\bar{t}W^\pm(N_{\text{jets}}=4)}$ assuming the scaling is the same for both processes. Both NFs
 1238 are left free-floating to constrain $t\bar{t}W$ yields in the 4-jet bin within CR 1b(+) and CR 1b(-).

1239 The final N_{jets} -parameterized function can then be represented by $\text{NF}_{t\bar{t}W(j')}$ as

$$\text{NF}_{t\bar{t}W(j')} = \left(\text{NF}_{t\bar{t}W^+(N_{\text{jets}}=4)} + \text{NF}_{t\bar{t}W^-(N_{\text{jets}}=4)} \right) \times \prod_{j=4}^{j'-1} \left(a_0 + \frac{a_1}{1+(j-4)} \right). \quad (6.7)$$

1240 The normalization is calculated and applied separately for each sub-sample of $t\bar{t}W^+$ and
 1241 $t\bar{t}W^-$ in a N_{jets} bin for $4 \leq N_{\text{jets}} < 10$. Due to small contributions in the CRs, events with
 1242 $N_{\text{jets}} < 4$ and $N_{\text{jets}} \geq 10$ are not normalized with this scheme. Instead, $N_{\text{jets}} < 4$ events are
 1243 fitted by propagating normalization in the 4-jet bin without additional shape correction. The
 1244 correction factor for $t\bar{t}W$ events with $N_{\text{jets}} \geq 10$ is obtained by summing up the overflow
 1245 from $N_{\text{jets}} = 10$ to $N_{\text{jets}} = 12$, described as $\sum_{j'=10}^{12} \prod_{j=4}^{j'-1} \left(a_0 + \frac{a_1}{1+(j-4)} \right)$. Events with
 1246 $N_{\text{jets}} \geq 13$ are negligible and are not included in the sum.

1247 The four CRs, CR $t\bar{t}W^\pm$ and CR 1b(\pm), are constructed to fit $\text{NF}_{t\bar{t}W^\pm(N_{\text{jets}}=4)}$ and
 1248 the scaling parameters $a_{0(1)}$, as well as validating the parameterization. Assuming the N_{jets}
 1249 distribution of $t\bar{t}W$ is similar across bins of $N_{b\text{-jets}}$, a fitted N_{jets} distribution in CR 1b(\pm)
 1250 can be used to describe the $t\bar{t}W$ parameterization at higher N_{jets} .

Chapter 7. Systematic Uncertainties

Physics analysis inherently incurs uncertainties in the form of statistical and systematic uncertainties, depending on the source. Statistical uncertainties occur in this analysis from sample size of collected data and simulated MC samples, and from the maximizing of the LH function. Systematic uncertainties depend on identifiable sources in the analysis i.e. from detector and reconstruction effects (experimental uncertainties) or theoretical modeling (theoretical uncertainties). Systematic uncertainties are represented as nuisance parameters (NP) in the profile LH fit. During the fit, systematic uncertainties with negligible impact on the final results can be pruned to simplify the statistical model and reduce computational complexity. This section outlines all uncertainties considered in this analysis.

7.1 Experimental uncertainties

7.1.1 Luminosity & pile-up reweighting

Uncertainty on the integrated luminosity of the 2015-2018 Run 2 data set is 0.83% [70], obtained by the LUCID-2 detector [95] for the primary luminosity measurements and complemented by the ID and calorimeters. Pile-up was modeled in MC and calibrated to data through pile-up reweighting, resulting in a set of calibration SFs and associated uncertainties.

7.1.2 Leptons

In general, calibrating MC simulations to match performance in data incurs uncertainties associated obtaining the MC-to-data calibration SFs, which are in turn propagated to observables in the analysis. The data-to-MC calibration of trigger, reconstruction, identification

Table 7.1: Summary of the experimental systematic uncertainties considered in this analysis.

Systematic uncertainty	Components
Event	
Luminosity	1
Pile-up reweighting	1
Electrons	
Trigger efficiency	1
Reconstruction efficiency [†]	1
Identification efficiency [†]	1
Isolation efficiency [†]	1
Energy scale	1
Energy resolution	1
Charge identification (ECIDS) efficiency [†]	1
Muons	
Trigger efficiency	2
Track-to-vertex association efficiency	2
Reconstruction/identification efficiency	2
Low- p_T (< 15 GeV) reconstruction/identification efficiency	2
Isolation efficiency	2
Charge-independent momentum scale	1
Charge-dependent momentum scale	4
Energy resolution (CB)	1
Energy resolution (ID & MS) [†]	2
Jets	
JES effective NP	15
JES η intercalibration	3
JES flavor composition	2
JES flavor response	1
JES pile-up	4
JES punch-through (FS/AF3)	2
JES non-closure	1
JES high- p_T single particle	1
JES b -jet response	1
JER effective NP	12
JER data/MC (FS/AF3)	2
JVT efficiency	1
GN2v01 b -tagging efficiency	85
GN2v01 c -tagging efficiency	56
GN2v01 light-tagging efficiency	42
E_T^{miss} track-based soft terms	
Transversal resolution	1
Longitudinal resolution	1
Longitudinal energy scale	1

[†]Not ready for the analysis, but will be included

and isolation efficiencies for electrons and muons incur associated uncertainties, with separate systematic and statistical components for those related to muons. Similarly, electron and muon energy-momentum scale and resolution are also subjected to calibration uncertainties estimated by varying the corresponding calibration quantity during simulation. Electron has an additional uncertainty related to ECIDS efficiency. Muon has additional uncertainties for charge-independent and charge-dependent momentum scale, as well as detector-specific track resolution. Systematic uncertainties for electron reconstruction, identification, isolation, ECIDS efficiencies and muon ID/MS energy resolution were not ready for the sample version used in this analysis, and are therefore not included.

7.1.3 Jets

Experimental uncertainties for jets are dominated by flavor tagging-related uncertainties, with subleading contributions from uncertainties related to JES [61], JER [60] and JVT [96] calibrations.

Jet energy scale

Uncertainties associated with JES are determined using data from LHC collisions along with MC simulated samples [61], decomposed into uncorrelated components:

- **Effective NPs:** 15 total p_T -dependent uncertainty components measured in situ, grouped based on their origin (2 detector-related, 4 modeling-related, 3 mixed, 6 statistical-related)
- **η intercalibration:** 6 total components (1 modeling-related, 4 non-closure and 1 statistical-related) associated with the correction of the forward jets' ($0.8 \leq |\eta| < 4.5$)

energy scale to that of the central jets ($|\eta| < 0.8$).

- **Flavor composition & response:** 2 components for relative quark-gluon flavor compositions in background and signal samples, and 2 components for responses to gluon-initiated versus quark-initiated jets.
- **Pile-up subtraction:** 4 components, 2 for μ (`OffsetMu`) and N_{PV} (`OffsetNPV`) modeling, 1 for residual p_T -dependency (`PtTerm`) and 1 for topology dependence on the per-event p_T density modeling (`RhoTopology`).
- **Punch-through effect treatment:** 2 terms for **GSC!** punch-through jet response deviation between data and MC, one for each detector response simulation method (AF3 and FS).
- **Non-closure:** 1 term applied to AF3 sample to account for the difference between AF3 and FS simulation.
- **High- p_T single-particle response:** 1 term for the response to high- p_T jets from single-particle and test-beam measurements.
- **b -jets response:** 1 term for the difference between b -jets and light-jets response.

Jet energy resolution

Measurements of JER were performed in bins of p_T and η , separately in data using in-situ techniques and in MC simulation using dijet events [60]. This analysis uses the full correlation JER uncertainty scheme provided for Run 2 analysis with 14 total components: 12 for effective NPs and 2 for difference between data and MC, separately for AF3 and FS [60].

Jet vertex tagging

The uncertainty associated with JVT is obtained by varying the JVT efficiency SFs within their uncertainty range [96]. This uncertainty accounts for remaining contamination from pile-up jets after applying pile-up suppression and MC generator choice.

Flavor tagging

Calibration SFs for b -tagging efficiencies and c -/light-jets mistagging rates are derived as a function of p_T for b -, c -, light-jets and PCBT score. The full set of flavor tagging-related uncertainties was reduced in dimensions by diagonalizing the uncertainty covariance matrix via eigendecomposition [64], resulting in a compact set of orthogonal NPs for this analysis: 85 for b -jets, 56 for c -jets and 42 for light-jets.

7.1.4 Missing transverse energy

Uncertainties on E_T^{miss} arise from possible mis-calibration of the soft-track component and are estimated using data-to-MC comparison of the p_T scale and resolution between the hard and soft E_T^{miss} components [68]. These uncertainties are represented by three independent terms: 1 for scale uncertainty and 2 for resolution uncertainty of the parallel and perpendicular components.

7.2 Modeling uncertainties

7.2.1 Signal and irreducible background uncertainties

The signal and background samples used were modeled using MC simulation. Most uncertainties on simulation parameters (e.g. generator choice, PS model) are estimated by varying the relevant parameters and comparing them with the nominal sample. Uncertainties involving PDF in particular for most processes in the analysis are set to a flat 1% uncertainty. Cross-section uncertainties were considered for all irreducible background except $t\bar{t}W$. Extra uncertainties for the production of four or more b -jets (additional b -jets) in association with $t\bar{t}X$ and HF jets were also considered due to a lack of theoretical predictions or dedicated measurements, rendering MC modeling challenging. Uncertainties from missing higher-order QCD corrections in MC simulation are estimated by varying the renormalization scale μ_R and factorization scale μ_F within seven different combinations

$$(\mu_R, \mu_F) = \{(0.5, 0.5), (0.5, 1), (1, 0.5), (1, 1), (1, 2), (2, 1), (2, 2)\}.$$

Process-specific uncertainty treatments are detailed below.

SM $t\bar{t}t\bar{t}$ background

The generator uncertainty for SM $t\bar{t}t\bar{t}$ background was evaluated between a nominal sample of MADGRAPH5_AMC@NLO and SHERPA. Parton shower uncertainty was evaluated between PYTHIA8 and HERWIG. The cross-section uncertainty was estimated to be 20% computed from a prediction at NLO in QCD+EW [78].

1347 $t\bar{t}t$ background

1348 The cross-section uncertainty for $t\bar{t}t$ was estimated to be 30% computed from a prediction
1349 at NLO in QCD+EW [78]. Events with additional b -jets also incur a 50% uncertainty.

1350 $t\bar{t}W$, $t\bar{t}Z$, $t\bar{t}H$ backgrounds

1351 For $t\bar{t}W$, $t\bar{t}Z$ and $t\bar{t}H$ backgrounds, an uncertainty of 50% is assigned to events with
1352 one additional truth b -jets that did not originate from a top quark decay, and an added
1353 50% uncertainty is assigned to events with two or more [97]. The generator uncertainty was
1354 estimated for $t\bar{t}Z$ using a MADGRAPH5_AMC@NLO nominal sample and a SHERPA sample,
1355 and for $t\bar{t}H$ using POWHEGBOX samples interfaced with PYTHIA8 (nominal) and HERWIG7.
1356 Cross-section uncertainties of 12% and 10% were applied to $t\bar{t}Z$ and $t\bar{t}H$ respectively [98].
1357 No $t\bar{t}W$ cross-section or PDF uncertainty was considered since the normalizations and jet
1358 multiplicity spectrum for $t\bar{t}W$ are estimated using the data-driven method described in
1359 section 6.3.3.

1360 Other backgrounds

1361 Other backgrounds include processes with small overall contribution in the SR. The
1362 cross-section uncertainty for tZ and tWH is considered to be 30% [99, 100]. A conservative
1363 cross-section uncertainty of 50% is applied to $t\bar{t}VV$, VVV and VH . For VV , the cross-
1364 section uncertainty is dependent on jet multiplicity and is considered to be 20%/50%/60%
1365 for events with $\leq 3/4/\geq 5$ jets [101]. For VV , $t\bar{t}VV$, VVV and VH events with additional
1366 truth b -jets, an uncertainty of 50% is applied.

7.2.2 Reducible background uncertainties

Reducible backgrounds consist of $t\bar{t}/V$ +HF jets and single top events. Reducible background has small contamination within the SR, thus uncertainties related to reducible background have minor impact. Treatment for reducible background in this analysis largely follows Ref. [36], except for QmisID.

Charge misidentification

Uncertainties on the QmisID background originate from the charge flip rates obtained using the data-driven method described in section 6.3.2. Four sources of uncertainty were considered: statistical uncertainty from the maximum LLH estimation using Equation 6.2; uncertainty from choice of the Z -mass window and sidebands; non-closure uncertainty defined as the relative difference between the number of SS and OS events; and statistical uncertainty from the N_{jets} dependency correction SFs. The combined uncertainties from all four sources are calculated separately for each region involved in section 6.3.2, and are treated as correlated across all regions. Figure 7.1 shows the uncertainty calculated for SR.

Internal (low γ^*) and material conversion

The normalization for internal and material conversion background are free parameters in the fit, as a result the only uncertainties evaluated are from the shape of the distributions used in the template fit method (see section 6.3.1). The uncertainties on internal (material) conversion are estimated based on the difference between data and MC prediction in a region enriched in $Z + \gamma \rightarrow \mu^+\mu^- + e^+e^-$ events.




Figure 7.1: Combined QmisID uncertainty rate for SR in bins of $|\eta|$ and p_T .

Heavy-flavor non-prompt lepton

Similar to the conversion backgrounds, the uncertainties on non-prompt HF decays come from the shape of the distributions, and are estimated by comparing data and MC prediction between all regions in the analysis on a per bin basis. The events used are required to contain at least one *Loose* reconstructed lepton used in the region selection criteria detailed in Table 6.2 to maintain orthogonality with the SR.

Light-flavor decays and other fake/non-prompt backgrounds

A conservative normalization uncertainty of 100% is assigned for light-flavor non-prompt lepton background [89], and an ad-hoc normalization uncertainty of 30% is applied to all other fake and non-prompt backgrounds. The shape uncertainties for these backgrounds are negligible.

Chapter 8. Results

8.1 Statistical analysis

This section provides an overview of the statistical methods needed to interpret the collected and simulated data to estimate unknown physics parameters and determine compatibility between data and the analysis hypothesis. For the BSM resonance search, the null hypothesis H_0 assumes only SM background contributions and none from any new resonance in the data.

8.1.1 Profile likelihood fit

Given a set of observed data points $\mathbf{x} = [x_1, x_2, \dots]$ and unknown parameters $\boldsymbol{\theta} = [\theta_1, \theta_2, \dots, \theta_n]$, the maximum likelihood method aims to find an estimate $\hat{\boldsymbol{\theta}}$ that maximizes the joint probability function $f(\mathbf{x}, \boldsymbol{\theta})$, or in other words the set of parameters that gives the highest probability of observing the collected data points for a particular model. The function to be maximized for this purpose is the log-likelihood (LLH) function $\ln \mathcal{L}(\mathbf{x}, \boldsymbol{\theta})$ where $\mathcal{L}(\mathbf{x}, \boldsymbol{\theta}) \equiv \prod_i f(x_i, \boldsymbol{\theta})$ is defined as the likelihood (LH) function. The LLH is maximized when $\partial/\partial\theta_i (\ln \mathcal{L}) = 0$ for each parameter θ_i .

For an usual binned physics analysis, the above variables for the LH function \mathcal{L} can be expressed as nuisance parameters (NP) $\boldsymbol{\theta}$ and number of events for a model $N_i(\mu)$ for the i^{th} bin, where μ is the targeted parameter of interest (POI). In this analysis, N_i is assumed to follow a Poisson distribution and depends on the following quantities: the signal strength μ defined as the ratio of observed to expected cross sections $\sigma_{\text{obs}}/\sigma_{\text{exp}}$; nuisance parameters $\boldsymbol{\theta}$ which represents the effects of systematic uncertainties, implemented in the

1419 LH function as Gaussian constraints; and normalization factors (NFs) $\boldsymbol{\lambda}$ that control the
 1420 normalization of background components that do not have a well-known cross section. The
 1421 Poisson probability of observing exactly N_i events for an expected number of event n_i is

$$\mathcal{P}(N_i|n_i(\mu, \boldsymbol{\lambda})) = \frac{n_i^{N_i} e^{-n_i}}{N_i!}. \quad (8.1)$$

1422 The expected Poisson event number in a bin i can be parameterized as

$$n_i = \mu s_i(\boldsymbol{\theta}) + \sum_j \lambda_j b_{ij}(\boldsymbol{\theta}), \quad (8.2)$$

1423 where s_i is the number of signal events in bin i of every region, and b_{ij} is the number of
 1424 events for a certain background source index j in bin i . The LH function in this analysis
 1425 can be written as

$$\mathcal{L}(\mathbf{N}|\mu, \boldsymbol{\theta}, \boldsymbol{\lambda}) = \left(\prod_i \mathcal{P}(N_i|n_i) \right) \cdot \prod_k \mathcal{G}(\theta_k), \quad (8.3)$$

1426 where $\mathcal{G}(\theta_k)$ is the Gaussian constraint for a NP k . The signal significance μ and NFs $\boldsymbol{\lambda}$ are
 1427 left unconstrained and are fitted simultaneously in the profile LH fit. From Neyman-Person
 1428 lemma [citation](#), the optimal test statistic for hypothesis testing is a function dependent on
 1429 the profile LH ratio defined as

$$q_\mu \equiv -2 \ln \frac{\mathcal{L}(\mu, \hat{\boldsymbol{\theta}}_\mu, \hat{\boldsymbol{\lambda}}_\mu)}{\mathcal{L}(\hat{\mu}, \hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\lambda}})}, \quad (8.4)$$

1430 where $\hat{\mu}$, $\hat{\boldsymbol{\theta}}$ and $\hat{\boldsymbol{\lambda}}$ are parameter values that optimally maximizes the LH function, and $\hat{\boldsymbol{\theta}}_\mu$,
 1431 $\hat{\boldsymbol{\lambda}}_\mu$ are NP and NF values respectively that maximize the LH function for a given μ .

8.1.2 Exclusion limits

8.2 Fit results

Fit setup

- Plain Asimov fit (**only mentioning briefly**): all regions included; simulated data used in the fit match exactly to MC prediction with nominal $\mu_{t\bar{t}Z'}$ set to 0 and allowed to free-float.

Purpose: to perform studies on optimizing fitted parameters and expected sensitivity; refining background estimation techniques; optimizing region definition and object definition

- Real SRs-blinded fit: similar to plain Asimov, but use observed data in CRs.

Purpose: study the behavior of background estimation using real observed data in CRs on Asimov data in SRs and assessing the influence of statistical effects on fitted parameters and expected sensitivity

- Real SRs-unblinded/ H_T fit: all regions included,

Limits

1447 Chapter 9. Summary

References

- [1] ATLAS Collaboration. *The ATLAS Experiment at the CERN Large Hadron Collider*. JINST 3 (2008), S08003 (cit. on p. 2).
- [2] D. H. Perkins. *Introduction to High Energy Physics*. 4th. Cambridge, UK: Cambridge University Press, Apr. 2000. ISBN: 9780521621960 (cit. on p. 3).
- [3] C. Burgard and D. Galbraith. *Standard Model of Physics*. URL: <https://texample.net/model-physics/> (visited on 06/02/2025) (cit. on p. 4).
- [4] H. Georgi. *Lie Algebras in Particle Physics: from Isospin to Unified Theories*. 2nd. CRC Press, 2000. ISBN: 9780429499210 (cit. on pp. 5, 8).
- [5] S. Navas et al. *Review of particle physics*. Phys. Rev. D 110.3 (2024), p. 030001 (cit. on p. 6).
- [6] M. Cristinziani and M. Mulders. *Top-quark physics at the Large Hadron Collider*. Journal of Physics G: Nuclear and Particle Physics 44.6 (2017), p. 063001. arXiv: 1606.00327 [hep-ex] (cit. on p. 6).
- [7] H. de la TorreTrisha Farooque and T. Farooque. *Looking beyond the Standard Model with Third Generation Quarks at the LHC*. Symmetry 14.3 (2022), p. 444 (cit. on p. 6).

- [8] Q.-H. Cao, J.-N. Fu, Y. Liu, X.-H. Wang, and R. Zhang. *Probing Topphilic New Physics via FourTopQuark Production*. [Chinese Physics C 45.9 \(2021\), p. 093107](#). arXiv: [2105.03372 \[hep-ph\]](#) (cit. on p. 6).
- [9] H. Beauchesne et al. *A case study about BSM vector resonances with direct couplings to the third quark generation*. [European Physical Journal C 80.5 \(2020\), p. 485](#). arXiv: [1908.11619 \[hep-ph\]](#) (cit. on p. 6).
- [10] F. Maltoni, D. Pagani, and S. Tentori. *Topquark pair production as a probe of light topphilic scalars and anomalous Higgs interactions*. [Journal of High Energy Physics 2024.9 \(Sept. 2024\), p. 098](#). arXiv: [2406.06694 \[hep-ph\]](#) (cit. on p. 6).
- [11] CMS Collaboration. *Search for $t\bar{t}H$ production in the $H \rightarrow b\bar{b}$ decay channel with leptonic $t\bar{t}$ decays in proton–proton collisions at $\sqrt{s} = 13 \text{ TeV}$* . [JHEP 03 \(2019\), p. 026](#). arXiv: [1804.03682 \[hep-ex\]](#) (cit. on p. 6).
- [12] M. E. Peskin and D. V. Schroeder. *An Introduction to Quantum Field Theory*. 1st. Reading, MA, USA: AddisonWesley, 1995. ISBN: 978-0-201-50397-5 (cit. on p. 7).
- [13] D. J. Gross. *The role of symmetry in fundamental physics*. [Proceedings of the National Academy of Sciences of the United States of America 93.25 \(Dec. 1996\), pp. 14256–14259](#) (cit. on p. 7).
- [14] C. Yang and R. Mills. *Conservation of Isotopic Spin and Isotopic Gauge Invariance*. [Phys. Rev. 96 \(1 1954\), pp. 191–195](#) (cit. on p. 8).
- [15] A. Milsted and T. J. Osborne. *Quantum Yang-Mills theory: An overview of a program*. [Phys. Rev. D 98 \(1 2018\), p. 014505](#) (cit. on p. 8).

- [16] A. Pich. *The Standard Model of electroweak interactions. 2004 European School of High-Energy Physics*. Feb. 2005, pp. 1–48. arXiv: [hep-ph/0502010](#) [[hep-ex](#)] (cit. on pp. 11, 12).
- [17] P. Dev and A. Pilaftsis. *High-temperature electroweak symmetry non-restoration from new fermions and implications for baryogenesis*. [Journal of High Energy Physics](#) **2020.9** (Sept. 2020), p. 012. arXiv: [2002.05174](#) [[hep-ph](#)] (cit. on p. 11).
- [18] P. Higgs. *Broken symmetries and the masses of gauge bosons*. [Phys. Rev. Lett.](#) **13** (16 1964), pp. 508–509 (cit. on p. 12).
- [19] P. Higgs. *Broken symmetries, massless particles and gauge fields*. [Physics Letters](#) **12.2** (1964), pp. 132–133. ISSN: 0031-9163 (cit. on p. 12).
- [20] F. Englert and R. Brout. *Broken Symmetry and the Mass of Gauge Vector Mesons*. [Phys. Rev. Lett.](#) **13** (9 1964), pp. 321–323 (cit. on p. 12).
- [21] ATLAS Collaboration. *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*. [Phys. Lett. B](#) **716** (2012), p. 1. arXiv: [1207.7214](#) [[hep-ex](#)] (cit. on p. 12).
- [22] CMS Collaboration. *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*. [Phys. Lett. B](#) **716** (2012), p. 30. arXiv: [1207.7235](#) [[hep-ex](#)] (cit. on p. 12).
- [23] J. Riebesell. *Higgs Potential*. URL: <https://tikz.net/higgs-potential/> (visited on 07/07/2025) (cit. on p. 13).
- [24] J. Goldstone, A. Salam, and S. Weinberg. *Broken Symmetries*. [Phys. Rev.](#) **127** (3 1962), pp. 965–970 (cit. on p. 14).

- [25] J. Ellis. *Higgs Physics. 2013 European School of High-Energy Physics*. 2015, pp. 117–168. arXiv: [1312.5672 \[hep-ph\]](#) (cit. on pp. 14, 15).
- [26] C. Collaboration. *Search for New Physics in High-Mass Electron-Positron Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV*. *Phys. Rev. Lett.* 99 (17 2007), p. 171802. arXiv: [0707.2524 \[hep-ex\]](#) (cit. on p. 15).
- [27] N. Arkani-Hamed, A. G. Cohen, and H. Georgi. *Electroweak symmetry breaking from dimensional deconstruction*. *Physics Letters B* 513.1-2 (July 2001), pp. 232–240. arXiv: [hep-ph/0105239 \[hep-ph\]](#) (cit. on p. 15).
- [28] T. Han, H. E. Logan, B. McElrath, and L.-T. Wang. *Phenomenology of the little Higgs model*. *Phys. Rev. D* 67 (9 2003), p. 095004. arXiv: [hep-ph/0301040 \[hep-ph\]](#) (cit. on p. 15).
- [29] P. Langacker and M. Plümacher. *Flavor changing effects in theories with a heavy Z' boson with family nonuniversal couplings*. *Phys. Rev. D* 62 (1 2000), p. 013006. arXiv: [hep-ph/0001204 \[hep-ph\]](#) (cit. on p. 15).
- [30] P. Langacker. *The Physics of Heavy Z' Gauge Bosons*. *Rev. Mod. Phys.* 81 (2009), pp. 1199–1228. arXiv: [0801.1345 \[hep-ph\]](#) (cit. on p. 15).
- [31] G. Ferretti and D. Karateev. *Fermionic UV completions of composite Higgs models*. *Journal of High Energy Physics* 2014.3 (Mar. 2014). ISSN: 1029-8479 (cit. on pp. 15, 18).
- [32] L. Vecchi. *A dangerous irrelevant UV-completion of the composite Higgs*. *JHEP* 02 (2017), p. 094. arXiv: [1506.00623 \[hep-ph\]](#) (cit. on pp. 15, 18).

- [33] K. Agashe, A. Delgado, M. J. May, and R. Sundrum. *RS1, custodial isospin and precision tests*. *JHEP* 08 (2003), p. 050. arXiv: [hep-ph/0308036 \[hep-ph\]](#) (cit. on p. 15).
- [34] K. Agashe, R. Contino, and A. Pomarol. *The Minimal composite Higgs model*. *Nucl. Phys. B* 719 (2005), pp. 165–187. arXiv: [hep-ph/0412089 \[hep-ph\]](#) (cit. on p. 15).
- [35] ATLAS Collaboration. *Search for top-philic heavy resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. *Eur. Phys. J. C* 84 (2024), p. 157. arXiv: [2304.01678 \[hep-ex\]](#) (cit. on p. 16).
- [36] ATLAS Collaboration. *Observation of four-top-quark production in the multilepton final state with the ATLAS detector*. *Eur. Phys. J. C* 83 (2023), p. 496. arXiv: [2303.15061 \[hep-ex\]](#) (cit. on pp. 16, 46, 56, 61, 71).
- [37] CMS Collaboration. *Observation of four top quark production in protonproton collisions at $\sqrt{s} = 13$ TeV*. *Physics Letters B* 847 (2023), p. 138290. arXiv: [2305.13439 \[hep-ex\]](#) (cit. on p. 16).
- [38] N. Greiner, K. Kong, J.-C. Park, S. C. Park, and J.-C. Winter. *Model-independent production of a top-philic resonance at the LHC*. *Journal of High Energy Physics* 2015.4 (2015), p. 29. ISSN: 1029-8479. arXiv: [1410.6099 \[hep-ph\]](#) (cit. on pp. 16, 17).
- [39] J. H. Kim, K. Kong, S. J. Lee, and G. Mohlabeng. *Probing TeV scale top-philic resonances with boosted top-tagging at the high luminosity LHC*. *Phys. Rev. D* 94 (3 2016), p. 035023. arXiv: [1604.07421 \[hep-ph\]](#) (cit. on p. 16).
- [40] G. C. Branco et al. *Theory and phenomenology of two-Higgs-doublet models*. *Phys. Rept.* 516 (2012), pp. 1–102. arXiv: [1106.0034 \[hep-ph\]](#) (cit. on p. 18).

- [41] S. Gori, I.-W. Kim, N. R. Shah, and K. M. Zurek. *Closing the wedge: Search strategies for extended Higgs sectors with heavy flavor final states*. *Phys. Rev. D* **93** (7 2016), p. 075038. URL: <https://link.aps.org/doi/10.1103/PhysRevD.93.075038> (cit. on p. 18).
- [42] P. S. B. Dev and A. Pilaftsis. *Maximally symmetric two Higgs doublet model with natural Standard Model alignment*. *Journal of High Energy Physics* **2014.12** (Dec. 2014), p. 024. arXiv: 1408.3405 [hep-ph] (cit. on p. 18).
- [43] P. Sabatini. *Evidence for four-top-quarks production with the ATLAS detector at the Large Hadron Collider*. Tech. rep. Geneva: CERN, 2021. URL: <https://cds.cern.ch/record/2784150> (cit. on p. 19).
- [44] ATLAS Collaboration. *Standard Model Summary Plots October 2023*. ATL-PHYS-PUB-2023-039. 2023. URL: <https://cds.cern.ch/record/2882448> (cit. on p. 22).
- [45] ATLAS Collaboration. *Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton–proton collision data at $\sqrt{s} = 13$ TeV*. *Eur. Phys. J. C* **79** (2019), p. 639. arXiv: 1902.04655 [physics.ins-det] (cit. on pp. 24, 39–41).
- [46] ATLAS Collaboration. *Performance of the ATLAS track reconstruction algorithms in dense environments in LHC Run 2*. *Eur. Phys. J. C* **77** (2017), p. 673. arXiv: 1704.07983 [hep-ex] (cit. on p. 30).
- [47] T. Cornelissen et al. *Concepts, design and implementation of the ATLAS New Tracking (NEWT)*. Tech. rep. Geneva: CERN, 2007. URL: <https://cds.cern.ch/record/1020106> (cit. on p. 30).

- [48] A. Salzburger and on behalf of the ATLAS Collaboration. *Optimisation of the ATLAS Track Reconstruction Software for Run-2*. [Journal of Physics: Conference Series 664.7 \(2015\), p. 072042](#) (cit. on p. 30).
- [49] R. Frühwirth. *Application of Kalman filtering to track and vertex fitting*. [Nucl. Instrum. Methods Phys. Res. A 262.2 \(1987\), pp. 444–450](#). ISSN: 0168-9002 (cit. on p. 30).
- [50] T. Cornelissen et al. *The global χ^2 track fitter in ATLAS*. [Journal of Physics: Conference Series 119.3 \(2008\), p. 032013](#) (cit. on p. 30).
- [51] ATLAS Collaboration. *Improved electron reconstruction in ATLAS using the Gaussian Sum Filter-based model for bremsstrahlung*. ATLAS-CONF-2012-047. 2012. URL: <https://cds.cern.ch/record/1449796> (cit. on p. 30).
- [52] D. Wicke. *A new algorithm for solving tracking ambiguities*. Tech. rep. Oct. 1998. URL: <https://cds.cern.ch/record/2625731> (cit. on p. 31).
- [53] ATLAS Collaboration. *Reconstruction of primary vertices at the ATLAS experiment in Run 1 proton–proton collisions at the LHC*. [Eur. Phys. J. C 77 \(2017\), p. 332](#). arXiv: [1611.10235 \[physics.ins-det\]](#) (cit. on p. 31).
- [54] W. Waltenberger, R. Frühwirth, and P. Vanlaer. *Adaptive vertex fitting*. [Journal of Physics G: Nuclear and Particle Physics 34.12 \(2007\), N343](#) (cit. on p. 31).
- [55] ATLAS Collaboration. *Secondary vertex finding for jet flavour identification with the ATLAS detector*. ATL-PHYS-PUB-2017-011. 2017. URL: <https://cds.cern.ch/record/2270366> (cit. on p. 31).

- [56] ATLAS Collaboration. *Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8 \text{ TeV}$ using the ATLAS detector*. *Eur. Phys. J. C* **76** (2016), p. 581. arXiv: [1510.03823 \[hep-ex\]](#) (cit. on pp. 32, 35).
- [57] ATLAS Collaboration. *Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1*. *Eur. Phys. J. C* **77** (2017), p. 490. arXiv: [1603.02934 \[hep-ex\]](#) (cit. on pp. 32, 33).
- [58] ATLAS Collaboration. *Jet reconstruction and performance using particle flow with the ATLAS Detector*. *Eur. Phys. J. C* **77** (2017), p. 466. arXiv: [1703.10485 \[hep-ex\]](#) (cit. on p. 34).
- [59] M. Cacciari, G. P. Salam, and G. Soyez. *The anti-kt jet clustering algorithm*. *Journal of High Energy Physics* **2008.04** (2008), p. 063 (cit. on p. 34).
- [60] ATLAS Collaboration. *Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*. *Eur. Phys. J. C* **81** (2021), p. 689. arXiv: [2007.02645 \[hep-ex\]](#) (cit. on pp. 35, 66, 67).
- [61] ATLAS Collaboration. *Jet energy scale measurements and their systematic uncertainties in proton–proton collisions at $\sqrt{s} = 13 \text{ TeV}$ with the ATLAS detector*. *Phys. Rev. D* **96** (2017), p. 072002. arXiv: [1703.09665 \[hep-ex\]](#) (cit. on pp. 35, 66).
- [62] ATLAS Collaboration. *Transforming jet flavour tagging at ATLAS*. Tech. rep. Submitted to: Nature Communications. Geneva: CERN, 2025. arXiv: [2505.19689](#) (cit. on pp. 36–38).
- [63] A. Vaswani et al. *Attention Is All You Need*. 2023. arXiv: [1706.03762 \[cs.CL\]](#) (cit. on p. 37).

- [64] ATLAS Collaboration. *Measurements of b-jet tagging efficiency with the ATLAS detector using $t\bar{t}$ events at $\sqrt{s} = 13$ TeV*. *JHEP* 08 (2018), p. 089. arXiv: 1805.01845 [hep-ex] (cit. on pp. 38, 68).
- [65] ATLAS Collaboration. *Electron Identification with a Convolutional Neural Network in the ATLAS Experiment*. ATL-PHYS-PUB-2023-001. 2023. URL: <https://cds.cern.ch/record/2850666> (cit. on p. 41).
- [66] ATLAS Collaboration. *Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV*. *Eur. Phys. J. C* 81 (2021), p. 578. arXiv: 2012.00578 [hep-ex] (cit. on pp. 42, 43).
- [67] ATLAS Collaboration. *Muon reconstruction performance of the ATLAS detector in proton–proton collision data at $\sqrt{s} = 13$ TeV*. *Eur. Phys. J. C* 76 (2016), p. 292. arXiv: 1603.05598 [hep-ex] (cit. on p. 43).
- [68] ATLAS Collaboration. *Performance of missing transverse momentum reconstruction with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV*. *Eur. Phys. J. C* 78 (2018), p. 903. arXiv: 1802.08168 [hep-ex] (cit. on pp. 44, 68).
- [69] ATLAS Collaboration. *E_T^{miss} performance in the ATLAS detector using 2015–2016 LHC pp collisions*. ATLAS-CONF-2018-023. 2018. URL: <https://cds.cern.ch/record/2625233> (cit. on p. 45).
- [70] ATLAS Collaboration. *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*. *Eur. Phys. J. C* 83 (2023), p. 982. arXiv: 2212.09379 [hep-ex] (cit. on pp. 46, 64).

- [71] J. Alwall et al. *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*. [JHEP 07 \(2014\), p. 079](#). arXiv: [1405.0301 \[hep-ph\]](#) (cit. on pp. 47, 49, 50).
- [72] NNPDF Collaboration, R. D. Ball, et al. *Parton distributions for the LHC run II*. [JHEP 04 \(2015\), p. 040](#). arXiv: [1410.8849 \[hep-ph\]](#) (cit. on pp. 47, 49, 50).
- [73] T. Sjöstrand et al. *An introduction to PYTHIA 8.2*. [Comput. Phys. Commun. 191 \(2015\), p. 159](#). arXiv: [1410.3012 \[hep-ph\]](#) (cit. on pp. 47, 49, 50).
- [74] ATLAS Collaboration. *ATLAS Pythia 8 tunes to 7 TeV data*. ATL-PHYS-PUB-2014-021. 2014. URL: <https://cds.cern.ch/record/1966419> (cit. on pp. 49, 50).
- [75] S. Frixione, E. Laenen, P. Motylinski, and B. R. Webber. *Angular correlations of lepton pairs from vector boson and top quark decays in Monte Carlo simulations*. [JHEP 04 \(2007\), p. 081](#). arXiv: [hep-ph/0702198](#) (cit. on p. 49).
- [76] P. Artoisenet, R. Frederix, O. Mattelaer, and R. Rietkerk. *Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations*. [JHEP 03 \(2013\), p. 015](#). arXiv: [1212.3460 \[hep-ph\]](#) (cit. on p. 49).
- [77] D. J. Lange. *The EvtGen particle decay simulation package*. [Nucl. Instrum. Meth. A 462 \(2001\), p. 152](#) (cit. on p. 49).
- [78] R. Frederix, D. Pagani, and M. Zaro. *Large NLO corrections in $t\bar{t}W^\pm$ and $t\bar{t}t\bar{t}$ hadroproduction from supposedly subleading EW contributions*. [JHEP 02 \(2018\), p. 031](#). arXiv: [1711.02116 \[hep-ph\]](#) (cit. on pp. 49, 69, 70).
- [79] E. Bothmann et al. *Event generation with Sherpa 2.2*. [SciPost Phys. 7.3 \(2019\), p. 034](#). arXiv: [1905.09127 \[hep-ph\]](#) (cit. on pp. 49, 50).

- [80] S. Schumann and F. Krauss. *A parton shower algorithm based on Catani–Seymour dipole factorisation*. *JHEP* 03 (2008), p. 038. arXiv: 0709.1027 [hep-ph] (cit. on p. 49).
- [81] S. Höche, F. Krauss, M. Schönherr, and F. Siegert. *A critical appraisal of NLO+PS matching methods*. *JHEP* 09 (2012), p. 049. arXiv: 1111.1220 [hep-ph] (cit. on p. 49).
- [82] S. Höche, F. Krauss, M. Schönherr, and F. Siegert. *QCD matrix elements + parton showers. The NLO case*. *JHEP* 04 (2013), p. 027. arXiv: 1207.5030 [hep-ph] (cit. on p. 49).
- [83] S. Catani, F. Krauss, B. R. Webber, and R. Kuhn. *QCD Matrix Elements + Parton Showers*. *JHEP* 11 (2001), p. 063. arXiv: hep-ph/0109231 (cit. on p. 49).
- [84] S. Höche, F. Krauss, S. Schumann, and F. Siegert. *QCD matrix elements and truncated showers*. *JHEP* 05 (2009), p. 053. arXiv: 0903.1219 [hep-ph] (cit. on p. 49).
- [85] F. Cascioli, P. Maierhöfer, and S. Pozzorini. *Scattering Amplitudes with Open Loops*. *Phys. Rev. Lett.* 108 (2012), p. 111601. arXiv: 1111.5206 [hep-ph] (cit. on p. 49).
- [86] A. Denner, S. Dittmaier, and L. Hofer. *COLLIER: A fortran-based complex one-loop library in extended regularizations*. *Comput. Phys. Commun.* 212 (2017), pp. 220–238. arXiv: 1604.06792 [hep-ph] (cit. on p. 49).
- [87] F. Buccioni et al. *OpenLoops 2*. *Eur. Phys. J. C* 79.10 (2019), p. 866. arXiv: 1907.13071 [hep-ph] (cit. on p. 49).
- [88] ATLAS Collaboration. *ATLAS data quality operations and performance for 2015–2018 data-taking*. *JINST* 15 (2020), P04003. arXiv: 1911.04632 [physics.ins-det] (cit. on p. 51).

- [89] ATLAS Collaboration. *Analysis of $t\bar{t}H$ and $t\bar{t}W$ production in multilepton final states with the ATLAS detector*. ATLAS-CONF-2019-045. 2019. URL: <https://cds.cern.ch/record/2693930> (cit. on pp. 56, 59, 72).
- [90] ATLAS Collaboration. *Measurement of the total and differential cross-sections of $t\bar{t}W$ production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. JHEP 05 (2024), p. 131. arXiv: 2401.05299 [hep-ex] (cit. on p. 56).
- [91] ATLAS Collaboration. *Search for new phenomena in events with same-charge leptons and b -jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. JHEP 12 (2018), p. 039. arXiv: 1807.11883 [hep-ex] (cit. on p. 59).
- [92] ATLAS Collaboration. *Evidence for $t\bar{t}t\bar{t}$ production in the multilepton final state in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. Eur. Phys. J. C 80 (2020), p. 1085. arXiv: 2007.14858 [hep-ex] (cit. on p. 61).
- [93] ATLAS Collaboration. *Search for R -parity-violating supersymmetry in a final state containing leptons and many jets with the ATLAS experiment using $\sqrt{s} = 13$ TeV proton–proton collision data*. Eur. Phys. J. C 81 (2021), p. 1023. arXiv: 2106.09609 [hep-ex] (cit. on p. 61).
- [94] E. Gerwick, T. Plehn, S. Schumann, and P. Schichtel. *Scaling Patterns for QCD Jets*. JHEP 10 (2012), p. 162. arXiv: 1208.3676 [hep-ph] (cit. on p. 62).
- [95] G. Avoni et al. *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*. JINST 13.07 (2018), P07017 (cit. on p. 64).
- [96] ATLAS Collaboration. *Tagging and suppression of pileup jets*. ATL-PHYS-PUB-2014-001. 2014. URL: <https://cds.cern.ch/record/1643929> (cit. on pp. 66, 68).

- [97] ATLAS Collaboration. *Measurements of inclusive and differential fiducial cross-sections of $t\bar{t}$ production with additional heavy-flavour jets in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. *JHEP* 04 (2019), p. 046. arXiv: 1811.12113 [hep-ex] (cit. on p. 70).
- [98] D. de Florian et al. *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*. *CERN Yellow Rep. Monogr.* 2 (2017), pp. 1–869. arXiv: 1610.07922 [hep-ph] (cit. on p. 70).
- [99] ATLAS Collaboration. *Measurement of the production cross-section of a single top quark in association with a Z boson in proton–proton collisions at 13 TeV with the ATLAS detector*. ATLAS-CONF-2017-052. 2017. URL: <https://cds.cern.ch/record/2273868> (cit. on p. 70).
- [100] F. Demartin, B. Maier, F. Maltoni, K. Mawatari, and M. Zaro. *tWH associated production at the LHC*. *EPJC* 77.1 (2017). arXiv: 1607.05862 [hep-ph]. URL: <https://doi.org/10.1140/epjc/s10052-017-4601-7> (cit. on p. 70).
- [101] ATLAS Collaboration. *Measurement of $W^\pm Z$ production cross sections and gauge boson polarisation in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*. *Eur. Phys. J. C* 79 (2019), p. 535. arXiv: 1902.05759 [hep-ex] (cit. on p. 70).