

## Boolean Lattice

A Boolean lattice is a relation satisfying a long list of conditions. This can be decomposed into the following ascending chain of definitions.

**Definition.** A *preorder*<sup>1</sup> is a relation  $\leq$  that is reflexive and transitive.

A *partial order* is a preorder  $\leq$  that is antisymmetric, i.e  $x \leq y \leq x$  implies  $x = y$ .

A *lattice* is a partial order in which every pair of elements  $x, y$  have a meet  $x \wedge y := \inf\{x, y\}$  and a join  $x \vee y := \sup\{x, y\}$ . A *bounded* lattice is a lattice which has a minimum  $\hat{0}$  and a maximum  $\hat{1}$ .

A *distributive lattice* is a lattice  $D$  in which

$$x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$$

and


$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

for all  $x, y, z \in D$ .

A *complement* of an element  $x$  of a bounded lattice  $L$  is an  $x' \in L$  such that  $x \wedge x' = \hat{0}$  and  $x \vee x' = \hat{1}$ . A *Boolean lattice* is a bounded distributive lattice in which every element has a unique complement.

**Example.** Taking the power set of a set  $X$  yields a Boolean lattice  $(2^X, \subseteq)$ . Meets are intersections, joins are unions, and the complement of a  $Y \subseteq X$  is  $X \setminus Y$ .

**Example.** The lattice  $N_5$  is given below.



Here,


$$x \wedge y = \hat{0} = x \wedge z$$

and


$$x \vee y = \hat{1} = x \vee z,$$

so  $y, z$  are both complements of  $x$ .

**Example.** The lattice  $M_3$  is given below.



A lattice is distributive if and only if it has no sublattices isomorphic to  $N_5$  or  $M_3$ . So, the lattice below is distributive.



But, this lattice is not Boolean since 2 has no complement. More generally, it is possible to show that the distributive lattice of positive divisors of an  $n \in \mathbb{Z}^+$  ordered by divisibility is Boolean if and only if  $n$  is squarefree.

**Example.** The two-element lattice  $\{0, 1\}$  with  $0 < 1$  is Boolean. The complement of a  $x \in \{0, 1\}$  is  $1 - x$ .

The chain at the beginning of this section also has two branches we will use.

**Definition.** An *equivalence relation* is a pre-order  $\equiv$  that is symmetric, i.e.  $x \equiv y$  implies  $y \equiv x$ .

A *linear order* is a partial order  $\leq$  such that every pair of elements  $x, y$  are comparable, i.e.  $x \leq y$  or  $y \leq x$ .

<sup>1</sup>Equivalently, a preorder is a category in which, for each pair of objects  $x, y$ , there is at most one morphism from  $x$  to  $y$ . Then, meets are products and joins are coproducts.

## Formulas

For each set  $A$ , we can construct the set  $W_A$  of well-formed formulas on  $A$  as follows. Let  $A$  be a set which is arbitrary unless otherwise specified.

Let  $C = \{\neg, \vee, \wedge, \rightarrow, (, )\}$  be a set of 6 currently meaningless symbols. Let  $S$  be the set of strings on  $A \cup C$ . Define maps  $\varepsilon_{\neg} : S \rightarrow S$  and  $\varepsilon_{\vee}, \varepsilon_{\wedge}, \varepsilon_{\rightarrow} : S \times S \rightarrow S$  by

$$\begin{aligned}\varepsilon_{\neg}(\psi) &= \neg(\psi) \\ \varepsilon_{\vee}(\phi, \psi) &= (\phi \vee \psi) \\ \varepsilon_{\wedge}(\phi, \psi) &= (\phi \wedge \psi) \\ \varepsilon_{\rightarrow}(\phi, \psi) &= (\phi \rightarrow \psi)\end{aligned}$$

for all  $\phi, \psi \in S$ . Inductively define subsets  $W_i$  of  $S$  for  $i \in \mathbb{Z}^+$  as follows. Let  $W_1 = A$ . If  $i \in \mathbb{Z}^+$  such that  $W_i$  has been defined, set

$$W_{i+1} = W_i \cup \varepsilon_{\neg}(W_i) \cup \bigcup_{\oplus \in \{\vee, \wedge, \rightarrow\}} \varepsilon_{\oplus}(W_i \times W_i).$$

Let  $W_A = \bigcup_{i \in \mathbb{Z}^+} W_i$ . The elements of  $A$  will be called *atoms*, and the elements of  $W_A$  will be called *well-formed formulas*. Elements of  $S \setminus W_A$  such as  $\rightarrow)((\wedge \neg$  are indeed ill-formed. Next, we see how to assign some meaning to the elements of  $W_A$ .

**Definition.** A map  $v : W_A \rightarrow \{0, 1\}$  is a *valuation* if and only if

$$\begin{aligned}v(\neg\phi) &= 1 - v(\phi) \\ v(\phi \vee \psi) &= \max\{v(\phi), v(\psi)\} \\ v(\phi \wedge \psi) &= \min\{v(\phi), v(\psi)\} \\ v(\phi \rightarrow \psi) &= \max\{1 - v(\phi), v(\psi)\}\end{aligned}$$

for all  $\phi, \psi \in W_A$ .

So, a valuation is a map which assigns a truth-value to each well-formed formula in a way that respects the connectives.

Recall that vector spaces are free over their bases. If  $V$  is a vector space with basis  $B$ , and  $U$  is another vector space, then any

map  $\alpha : B \rightarrow U$  can be uniquely extended to a linear map  $\bar{\alpha} : V \rightarrow U$ .

$$\begin{array}{ccc} B & \xrightarrow{\alpha} & U \\ \downarrow & \nearrow \bar{\alpha} & \\ V & & \end{array}$$

Similarly, the set of well-formed formulas  $W_A$  is free over the atoms  $A$ .

**Proposition.** Suppose  $v : A \rightarrow \{0, 1\}$  is a map. Then, there exists a unique valuation  $\bar{v} : W_A \rightarrow \{0, 1\}$  such that  $\bar{v}(a) = v(a)$  for all  $a \in A$ .

$$\begin{array}{ccc} A & \xrightarrow{v} & \{0, 1\} \\ \downarrow & \nearrow \bar{v} & \\ W_A & & \end{array}$$

Now, let's see an example of how well-formed formulas can be used to express useful things. Let  $P$  be a set. We have the following bijective correspondence between the relations on  $P$  and the valuations  $W_{P \times P} \rightarrow \{0, 1\}$ . Using the previous proposition, for each relation  $\sim$  on  $P$ , we can let  $v_{\sim} : W_{P \times P} \rightarrow \{0, 1\}$  be the valuation with  $v_{\sim}((y, z)) = 1$  if and only if  $y \sim z$ . Then

$$\sim \mapsto v_{\sim}$$

is a bijection from the set of relations on  $P$  to the set of valuations  $W_{P \times P} \rightarrow \{0, 1\}$ . If relations on  $P$  are viewed as subsets of  $P \times P$ , then the inverse bijection is

$$v \mapsto v^{-1}(\{1\}) \cap P \times P.$$

Observe that a relation  $\sim$  on  $P$  is reflexive if and only if

$$1 = v_{\sim}((p, p))$$

for all  $p \in P$ . A relation  $\sim$  on  $P$  is transitive if and only if

$$1 = v_{\sim}(((p, q) \wedge (q, r)) \rightarrow (p, r))$$

for all  $p, q, r \in P$ . Indeed, if  $p, q, r \in P$  with  $p \sim q$  and  $q \sim r$  and

$$1 = v_{\sim}(((p, q) \wedge (q, r)) \rightarrow (p, r)),$$

then  $v(p, q) = v(q, r) = 1$  and

$$\begin{aligned} 1 &= \max\{1 - \min\{v(p, q), v(q, r)\}, v(p, r)\} \\ &= \max\{0, v(p, r)\}, \end{aligned}$$

whence  $v(p, r) = 1$  and  $p \sim r$ . If  $\sim$  is transitive and  $p, q, r \in P$  and  $v(p, r) = 0$ , then  $v(p, q) = 0$  or  $v(q, r) = 0$ , so

$$\begin{aligned} v_\sim(((p, q) \wedge (q, r)) \rightarrow (p, r)) \\ &= \max\{1 - \min\{v(p, q), v(q, r)\}, 0\} \\ &= 1 - \min\{v(p, q), v(q, r)\} \\ &= 1. \end{aligned}$$

Similarly for antisymmetry and comparability. Let

$$\begin{aligned} T = &\{(p, q) \in P \times P \mid p = q\} \\ &\cup \{((p, q) \wedge (q, r)) \rightarrow (p, r) \mid p, q, r \in P\} \\ &\cup \{\neg((p, q) \wedge (q, p)) \mid p, q \in P \text{ and } p \neq q\} \\ &\cup \{(p, q) \vee (q, p) \mid p, q \in P\}. \end{aligned}$$

So, a relation  $\sim$  on  $P$  is a linear order if and only if  $1 = \inf v_\sim(T)$ .

### Preorder to Partial Order

The following is a natural<sup>2</sup> way to obtain a partial order from a preorder. Suppose  $\leq$  is a preorder on a set  $X$ . Define an equivalence relation  $\equiv$  on  $X$  by  $x \equiv y$  if and only if  $x \leq y$  and  $y \leq x$ . For each  $x \in X$ , let

$$[x] = \{y \in X \mid y \equiv x\}$$

denote the equivalence class of  $x$ . Let

$$X^* = (X / \equiv) = \{[x] \mid x \in X\}$$

denote the quotient of  $X$  by  $\equiv$ . Define the relation  $\leq^*$  on  $X^*$  by  $[x] \leq^* [y]$  if and only if  $x \leq y$ .

<sup>2</sup>This way of turning preorders into partial orders is functorial, and is left adjoint to the forgetful functor from the category of partial orders to the category of preorders.

**Proposition.** The relation  $\equiv$  is indeed an equivalence relation. The relation  $\leq^*$  is a well-defined partial order.

*Proof.* The relation  $\equiv$  is reflexive and transitive since  $\leq$  is reflexive and transitive. The relation  $\equiv$  is symmetric since its definition is symmetric.

If  $a \equiv y \leq z \equiv b$ , then  $a \leq y \leq z \leq b$  and  $a \leq b$ . The relation  $\leq^*$  is a partial order since  $\leq$  is a partial order.  $\square$

Define a relation  $\models$  on  $W_A$  by  $\phi \models \psi$  if and only if  $v(\phi) \leq v(\psi)$  for all valuations  $v$ . Equivalently,  $\phi \models \psi$  if and only if, for all valuations  $v$ , we have  $v(\phi) = 1$  implies  $v(\psi) = 1$ .

**Proposition.** The relation  $\models$  is a preorder.

*Proof.* Suppose  $\phi \in W_A$ . Since  $v(\phi) \leq v(\phi)$  for all valuations  $v$ , we have  $\phi \models \phi$ .

Suppose  $\phi, \psi, \chi \in W_A$  with  $\phi \models \psi$  and  $\psi \models \chi$ . If  $v$  is a valuation, then  $v(\phi) \leq v(\psi)$  and  $v(\psi) \leq v(\chi)$ , and thus  $v(\phi) \leq v(\chi)$ . So  $\phi \models \chi$ .  $\square$

So, we have a partial order  $\models^*$  with  $\phi \models \psi$  if and only if  $[\phi] \models^* [\psi]$ . Note that  $v(\phi) = v(\psi)$  for all valuations  $v$  if and only if  $[\phi] = [\psi]$ .

**Proposition.** The partial order  $\models^*$  is a Boolean lattice.

*Proof.* Suppose  $[\phi], [\psi] \in W_A^*$ . For all valuations  $v$ , we have

$$v(\phi) \leq \max\{v(\phi), v(\psi)\} = v(\phi \vee \psi),$$

so  $\phi \models \phi \vee \psi$ . Hence  $[\phi] \models^* [\phi \vee \psi]$ . Similarly,  $[\psi] \models^* [\phi \vee \psi]$ . So  $[\phi \vee \psi]$  is a common upper bound for  $[\phi]$  and  $[\psi]$ .

Suppose  $[\chi] \in W_A^*$  is another common upper bound for  $[\phi]$  and  $[\psi]$ . Then, for all valuations  $v$ , we have  $v(\phi) \leq v(\chi)$  and  $v(\psi) \leq v(\chi)$ , so

$$v(\phi \vee \psi) = \max\{v(\phi), v(\psi)\} \leq v(\chi).$$

Hence  $\phi \vee \psi \models \chi$ , i.e.  $[\phi \vee \psi] \models^* [\chi]$ .

So,  $[\phi \vee \psi]$  is the least upper bound of  $[\phi]$  and  $[\psi]$ , i.e.

$$[\phi] \vee [\psi] = [\phi \vee \psi].$$

Similarly,

$$[\phi] \wedge [\psi] = [\phi \wedge \psi].$$

So,  $\models^*$  is a lattice. Also, this shows that it makes sense to use the same symbols for disjunction and conjunction for logic as for join and meet for orders.

A *tautology* is a  $\phi \in W_A$  such that  $v(\phi) = 1$  for all valuations  $v$ . Pick an  $a \in W_A$  and set  $\top = (a \vee (\neg a))$ . For all valuations  $v$ , we have

$$\begin{aligned} v(\top) &= v(a \vee (\neg a)) \\ &= \max\{v(a), v(\neg a)\} \\ &= \max\{v(a), 1 - v(a)\} \\ &= 1 \end{aligned}$$

So, for all  $\phi \in W_A$ , we have

$$v(\phi) \leq 1 = v(\top)$$

for all valuations  $v$ , and thus  $\phi \models \top$ , i.e.  $[\phi] \models^* [\top]$ . So,  $[\top]$  is the maximum element of  $W_A^*$ .

A *contradiction* is a  $\phi \in W_A$  such that  $v(\phi) = 0$  for all valuations  $v$ . Pick an  $a \in W_A$ , set  $\perp = (a \wedge (\neg a))$ , and observe that similarly  $\perp$  is a contradiction and hence  $[\perp]$  is the minimum element of  $W_A^*$ .

Suppose  $[\phi] \in W_A^*$ . For all valuations  $v$ , we have

$$\begin{aligned} v(\phi \vee (\neg \phi)) &= \max\{v(\phi), v(\neg \phi)\} \\ &= \max\{v(\phi), 1 - v(\phi)\} \\ &= 1 \\ &= v(\top). \end{aligned}$$

It follows that

$$[\phi] \vee [\neg \phi] = [\phi \vee (\neg \phi)] = [\top],$$

Similarly,

$$[\phi] \wedge [\neg \phi] = [\perp].$$

So  $[\neg \phi]$  is a complement of  $[\phi]$ . Suppose  $[\psi] \in W^*$  is another complement of  $[\phi]$ . Then

$$[\phi \vee \psi] = [\phi] \vee [\psi] = [\top].$$

Suppose  $v$  is a valuation. Then

$$\begin{aligned} 1 &= v(\top) \\ &= v(\phi \vee \psi) \\ &= \max\{v(\phi), v(\psi)\}. \end{aligned}$$

If  $v(\phi) = 0$ , then this implies

$$v(\psi) = 1 = 1 - 0 = v(\neg \phi).$$

Otherwise, if  $v(\phi) = 1$ , consider instead

$$\begin{aligned} 0 &= v(\perp) \\ &= v(\phi \wedge \psi) \\ &= \min\{v(\phi), v(\psi)\}, \end{aligned}$$

whence

$$v(\psi) = 0 = 1 - 1 = v(\neg \phi).$$

So  $v(\psi) = v(\neg \phi)$  for all valuations  $v$ , i.e.  $[\psi] = [\neg \phi]$ . So, the complement of  $[\phi]$  is unique.  $\square$

We now have a counterexample to the infinite extension of the following theorem.

**Theorem.** Each finite Boolean lattice is isomorphic to  $(2^{[n]}, \subseteq)$  for some  $n \in \mathbb{Z}^+ \cup \{0\}$ .

Recall that Cantor's Theorem says that if  $X$  is a set, then there are no injections  $2^X \rightarrow X$ . In particular, the powerset of a set is either finite or uncountable. If  $A$  is chosen to be countably infinite, then the Boolean lattice  $W_A^*$  is countably infinite and hence not isomorphic to a powerset lattice.

## Partial Order to Linear Order

**Definition.** A subset  $S \subseteq W_A$  is *satisfiable* if and only if there exists a valuation  $v$  such that  $\inf v(S) = 1$ , i.e. such that  $v(\phi) = 1$  for all  $\phi \in S$ .

**Example.** Suppose  $a \in A$ . Then  $\{a, (\neg a)\}$  is not satisfiable since, if  $v$  is a valuation with  $v(a) = 1$ , then  $v(\neg a) = 1 - 1 = 0 \neq 1$ .

**Theorem.** A subset  $S \subseteq W$  is satisfiable if and only if every finite subset of  $S$  is satisfiable.

The theorem above is the Compactness Theorem.<sup>3</sup> It can be used to extend various results to the infinite case. Dilworth's Theorem for partial orders of finite width is one example. Another is the following.

**Definition.** A *linear extension* of a partial order  $\preceq$  on a set  $P$  is a linear order  $\leq$  on  $P$  such that  $p \preceq q$  implies  $p \leq q$ . If relations on  $P$  are viewed as subsets of  $P \times P$ , then this is the same as saying  $\preceq \subseteq \leq$ .

**Lemma.** Every finite partial order has a linear extension.

*Proof.* Suppose  $(P, \preceq)$  is a partial order, and  $y, z \in P$  are incomparable. Define a relation  $\leq$  on  $P$  by  $p \leq q$  if and only if  $p \preceq q$ , or  $p \preceq y$  and  $q \preceq z$ . So,  $\leq$  is an extension of  $\preceq$  having fewer pairs of incomparable elements than  $\preceq$ . Observe that it is possible to use the fact that  $\preceq$  is a partial order to check that  $\leq$  is a partial order. So, induction can be used to obtain the desired result.  $\square$

**Proposition.** Every partial order has a linear extension.

*Proof.* Suppose  $(P, \preceq)$  is a partial order. Let

$$S_e = \{(p, q) \in P \times P \mid p \preceq q\}$$

$$S_t = \{((p, q) \wedge (q, r)) \rightarrow (p, r) \mid p, q, r \in P\}$$

$$S_a = \{\neg((p, q) \wedge (q, p)) \mid p, q \in P \text{ and } p \neq q\}$$

$$S_c = \{(p, q) \vee (q, p) \mid p, q \in P\}.$$

Let

$$S = S_e \cup S_t \cup S_a \cup S_c \subseteq W_{P \times P}.$$

Suppose  $F \subseteq S$  is finite. Let  $Q \subseteq P$  be the set of consisting of all elements of  $P$  appearing in  $F$ . Then  $Q$  is finite since  $F$  is finite and the elements of  $F$  are finite strings. Then  $Q$  is a finite subposet of  $P$ , so the lemma yields a linear extension  $\leq$  of the ordering on  $Q$  induced by  $\preceq$ . Let  $v : W_{Q \times Q} \rightarrow \{0, 1\}$  be the valuation with  $v(r, s) = 1$  for  $r \leq s$  and  $v(r, s) = 0$  for  $r \not\leq s$ .

Suppose  $(p, q) \in S_e \cap F$ . Then  $p \preceq q$  and  $p, q \in Q$ . Since  $p \preceq q$  and  $\leq$  is an extension of  $\preceq$ , we have  $p \leq q$ . Then  $v(p, q) = 1$ . Similarly, since  $\leq$  is transitive, antisymmetric, and has comparability, it is possible to show that  $v(\phi) = 1$  for all  $\phi \in (S_t \cup S_a \cup S_c) \cap F$ .

It follows that  $F$  is satisfiable. Using the Compactness Theorem, there exists a valuation  $v : W_{P \times P} \rightarrow \{0, 1\}$  such that  $v(\phi) = 1$  for all  $\phi \in S$ . Using the last paragraph of the second section, the relation  $\leq$  on  $P$  given by

$$p \leq q \quad \text{if and only if} \quad v(p, q) = 1$$

for all  $p, q \in P$  is a linear extension of  $\preceq$ .  $\square$

## Video

[youtu.be/f3a-o-Vn7Fg](https://youtu.be/f3a-o-Vn7Fg)

## References

- [1] B.A. Davey and H.A. Priestly, *Introduction to Lattices and Order*, first edition
- [2] J.B. Nation, *Notes on Lattice Theory*. [math.hawaii.edu/~jb/math618/Nation-LatticeTheory.pdf](http://math.hawaii.edu/~jb/math618/Nation-LatticeTheory.pdf)
- [3] P. Johnstone, *Notes on Logic and Set Theory*

<sup>3</sup>The name of this theorem makes sense since it can be proved using Tychonoff's theorem.