


Antimicrobial Stewardship Strategy:

Identification of inappropriate pathogen/ antimicrobial combination (“bug-drug mismatch”)

Identification and modification of therapy in patients with positive cultures who are receiving ineffective therapy.



@istock.com/zmeel

Priority Level: **B**

Difficulty Level: **3**

Program Stage:

- Early
- Intermediate
- ✓ Advanced

For more information on these criteria and how they were developed, please see the

[Antimicrobial Stewardship Strategy Criteria Reference Guide](#).

Description

This is an overview and not intended to be an all-inclusive summary. As a general principle, patients must be monitored by the health care team after changes to therapy resulting from recommendations made by the antimicrobial stewardship team.

“Bug-drug mismatch” refers to a situation in which the antimicrobial(s) a patient is receiving do(es) not provide adequate therapy (e.g., is resistant) for the microbiologically identified organism presumed to be causing the clinical infection. For example, culture and susceptibility results of drainage from an intra-abdominal abscess may isolate *Klebsiella sp.* resistant to piperacillin/tazobactam in a patient empirically started on piperacillin/tazobactam for an intra-abdominal infection. Or a patient may be receiving intravenous ciprofloxacin for pyelonephritis, with *Escherichia coli* resistant to ciprofloxacin cultured from the blood and urine. An extreme example could include the identification of a significant organism in a clinical culture from a patient who is not receiving therapy. It is important, of course, to consider whether the organism cultured by the microbiology laboratory is of clinical significance (i.e., not colonization or contamination) when assessing therapy.

Although identification of bug-drug mismatch is part of a comprehensive assessment of a patient’s therapy, it may also be performed as a standalone antimicrobial stewardship strategy.

Implementation

- Stewardship team/individual must establish a process for identifying and targeting patients with potential bug-drug mismatch. This can be facilitated by a computer program with pharmacy and microbiology interfaces. An alternative but more labour-intensive approach would be for the antimicrobial stewardship team to obtain a list of patients with positive culture results (could begin with blood cultures, and then expand to additional cultures as resources allow) and manually review current antimicrobial therapies.
- Certain commercial clinical decision support systems can facilitate the identification of bug-drug mismatch.

Bug-drug mismatch has the potential to significantly compromise patient outcomes, so communication (preferably verbal) with the prescriber is required. Messages left in the chart may not be noticed in a timely fashion.

Advantages

- Identifies patients at highest risk of therapeutic failure and subsequent morbidity and mortality.
- Because the incidence of bug-drug mismatch is generally not high, this may be an appealing strategy for a program wishing to initiate a prospective intervention, and/or for programs with fewer resources, providing the process for patient identification is not too onerous.
- Can improve the appropriateness of therapy in patients whose culture results were inadvertently overlooked, and can expedite changes to more appropriate therapy.
- Provides an opportunity to reinforce with prescribers the importance of checking final culture and susceptibility results, and using microbiologic data to modify/optimize antimicrobial prescribing.

Disadvantages

- May be labour-intensive if manual review of culture results and patient therapy is required.
- Chart review is still required for proper interpretation and assessment of the clinical issue.
- Requires ongoing resources of the antimicrobial stewardship team.
- Because the incidence of bug-drug mismatch is generally not high, the time spent identifying potential mismatches may not justify the benefits.

Requirements

- Access to microbiology results:
 - Microbiology laboratory system capable of providing a list of patients with new, positive microbiology results.
 - An integrated information technology system to identify patients with potential mismatch.
- Availability of a clinician (usually pharmacist or physician) able to assess the clinical significance of culture results and make recommendations for patients' therapy.

Associated Metrics

- Number of patients identified with bug-drug mismatch.
- Rating of potential consequences if not detected (patient safety perspective).

Useful References

Select articles to provide supplemental information and insight into the strategy described and/or examples of how the strategy was applied; not a comprehensive reference list. URLs are provided when materials are freely available on the Internet.

- Griffith M, Postelnick M, Scheetz M. Antimicrobial stewardship programs: methods of operation and suggested outcomes. *Expert Rev Anti Infect Ther.* 2012;10(1):63–73.

Discusses identification of bug-drug mismatch as a stewardship intervention.

Samples/Examples

- [Example 1: Grey Bruce Health Services - Microbiology Condensed Line Listing Report](#)
- [Example 2: Royal Victoria Regional Health Centre - Pharmacy/Microbiology Report](#)
- [Example 3: The Scarborough Hospital - ICNet System Sample Automated Alerts](#)

These documents have been generously shared by various health care institutions to help others develop and build their antimicrobial stewardship programs. We recommend crediting an institution when adopting a specific tool/form/pathway in its original form.

Examples that contain clinical or therapeutic recommendations may not necessarily be consistent with published guidelines, or be appropriate or directly applicable to other institutions. All examples should be considered in the context of the institution's population, setting and local antibiogram.

The materials and information in this section are not owned by Public Health Ontario. Neither Public Health Ontario nor the institution sharing the document shall be responsible for the use of any tools and resources by a third party.

Links with Other Strategies

- [Clinical decision support systems/computerized physician order entry](#)
- [Prospective audit with intervention and feedback](#)
- [Targeted review of patients with bacteremia/fungemia](#)

Disclaimer

This document may be freely used without permission for non-commercial purposes only and provided that appropriate credit is given to Public Health Ontario. No changes and/or modifications may be made to the content without explicit written permission from Public Health Ontario.

Citation

Ontario Agency for Health Protection and Promotion (Public Health Ontario). Antimicrobial Stewardship Strategy: Identification of inappropriate pathogen/antimicrobial combination (“bug-drug mismatch”). Toronto, ON: Queen’s Printer for Ontario; 2016.

©Queen’s Printer for Ontario, 2016

For further information

[Antimicrobial Stewardship Program](#), Infection Prevention and Control, Public Health Ontario.

Email: asp@oahpp.ca

Public Health Ontario acknowledges the financial support of the Ontario Government.



Example 1: Grey Bruce Health Services - Microbiology Condensed Line Listing Report



Date: 09/23/15 06:35:15

Microbiology Condensed Line Listing Report

Id: Operations

Sequence: OS Pos Suscep Condensed Line Listing

GBRHC

Page: 1

Filter:

1: Date/Time - Complete - 09/22/15 00:00 - 09/23/15 23:59
2: Positive Indicator - Positive:
3: Service Resource - Owen Sound Lab, Saugeen Lab, Meaford Lab,
Markdale Lab, Wiarton Lab, Lions Head Lab
4: Task Result - Susceptibility Interp Result - I, MS, NA, NS, R,
S

Sort(s):

1: Client [A]
2: Acscn# [A]
3: Orderable [A]

Name:	MRN:	Receive Date:	Procedure:	Accession:	Source:	Final Isolate:
		09/18/15 13:12	Culture-Wou		Wound	MRSA, *, *, *
		09/20/15 09:07	Culture-Uri		Urine	EC
		09/20/15 16:57	Culture-Uri		Urine	EC
		09/20/15 09:07	Culture-Uri		Urine	urpred1, EC
		09/20/15 09:07	Culture-Uri		U Cath	MRSA, *
		09/19/15 17:13	Culture-Uri		Urine	EC
		09/20/15 16:57	Culture-Uri		Urine	EC, *
		09/20/15 16:57	Culture-Uri		Urine	urpred1, EC
Total accession selected:		8				

***** END OF REPORT *****

Disclaimer

This resource was created by Grey Bruce Health Services. PHO is not the owner of this content and does not take responsibility for the information provided within this document. Neither PHO nor Grey Bruce Health Services shall be responsible for the subsequent use of any tools and resources by any third party.

Example 2: Royal Victoria Regional Health Centre - Pharmacy/Microbiology Report



DATE: 20/10/15 @ 1615		Royal Victoria Pharmacy **LIVE**		PAGE 9																																																																																					
USER: MCAL		Microbiology Report																																																																																							
Patient	[REDACTED]	Responsible Doctor	[REDACTED]																																																																																						
Account Number	[REDACTED]	Location	Unit Number	[REDACTED]																																																																																					
Age/Sex	[REDACTED]	Room	Registered Date	[REDACTED]																																																																																					
Status	ADM IN	Bed	Discharged Date	[REDACTED]																																																																																					
Rx Data																																																																																									
Rx Number	Medications	Dose	Directions	Start Stop Status																																																																																					
	0.9% sodium chloride	250 mL	Q12 IV	18/10 25/10																																																																																					
	vancomycin	1,000 mg																																																																																							
	piperacillin/tazo 3.375 g 100 mL		Q6 IV	19/10 26/10																																																																																					
Microbiology Specimens																																																																																									
Specimen:	Collected: 17/10/15-2336	Status: RES	Req#:																																																																																						
	Received: 17/10/15-2348	Source: BLOOD	Sp Desc:																																																																																						
> Blood culture Preliminary																																																																																									
No growth after 2 days incubation																																																																																									
Specimen:	Collected: 17/10/15-2336	Status: RES	Req#:																																																																																						
	Received: 17/10/15-2348	Source: BLOOD	Sp Desc:																																																																																						
> Blood culture Preliminary																																																																																									
No growth after 2 days incubation																																																																																									
Specimen:	Collected: 18/10/15-0145	Status: RES	Req#:																																																																																						
	Received: 18/10/15-0152	Source: BLOOD	Sp Desc:																																																																																						
Specimen:	Collected: 18/10/15-UNK	Status: COMP	Req#:																																																																																						
	Received: 18/10/15-1159	Source: FINGER	Sp Desc: Unknown																																																																																						
<table border="0"> <tr> <td></td> <td>S aureus</td> <td></td> <td>P rettgeri</td> <td></td> </tr> <tr> <td></td> <td>M.I.C.</td> <td>RX</td> <td>M.I.C.</td> <td>RX</td> </tr> <tr> <td>TRIMETH/SULFA</td> <td><=10</td> <td>S</td> <td><=20</td> <td>S</td> </tr> <tr> <td>AMPICILLIN</td> <td></td> <td></td> <td><=2</td> <td>R</td> </tr> <tr> <td>CEFAZOLIN</td> <td></td> <td></td> <td><=4</td> <td>R</td> </tr> <tr> <td>CEFTAZIDIME</td> <td></td> <td></td> <td><=1</td> <td>S</td> </tr> <tr> <td>CEFTRIAXONE</td> <td></td> <td></td> <td><=1</td> <td>S</td> </tr> <tr> <td>CIPROFLOXACIN</td> <td></td> <td></td> <td><=0.25</td> <td>S</td> </tr> <tr> <td>CLINDAMYCIN</td> <td><=0.25</td> <td>S</td> <td></td> <td></td> </tr> <tr> <td>ERYTHROMYCIN</td> <td><=0.25</td> <td>S</td> <td></td> <td></td> </tr> <tr> <td>GENTAMICIN</td> <td></td> <td></td> <td><=1</td> <td>S</td> </tr> <tr> <td>ERTAPENEM</td> <td></td> <td></td> <td><=0.5</td> <td>S</td> </tr> <tr> <td>OXACILLIN</td> <td><=0.25</td> <td>S</td> <td></td> <td></td> </tr> <tr> <td>PENICILLIN G</td> <td>>=0.5</td> <td>R</td> <td></td> <td></td> </tr> <tr> <td>TOBRAMYCIN</td> <td></td> <td></td> <td><=1</td> <td>S</td> </tr> <tr> <td>VANCOMYCIN</td> <td><=0.5</td> <td>S</td> <td></td> <td></td> </tr> <tr> <td>PIPERA/TAZOACT</td> <td></td> <td></td> <td><=4</td> <td>S</td> </tr> </table>						S aureus		P rettgeri			M.I.C.	RX	M.I.C.	RX	TRIMETH/SULFA	<=10	S	<=20	S	AMPICILLIN			<=2	R	CEFAZOLIN			<=4	R	CEFTAZIDIME			<=1	S	CEFTRIAXONE			<=1	S	CIPROFLOXACIN			<=0.25	S	CLINDAMYCIN	<=0.25	S			ERYTHROMYCIN	<=0.25	S			GENTAMICIN			<=1	S	ERTAPENEM			<=0.5	S	OXACILLIN	<=0.25	S			PENICILLIN G	>=0.5	R			TOBRAMYCIN			<=1	S	VANCOMYCIN	<=0.5	S			PIPERA/TAZOACT			<=4	S
	S aureus		P rettgeri																																																																																						
	M.I.C.	RX	M.I.C.	RX																																																																																					
TRIMETH/SULFA	<=10	S	<=20	S																																																																																					
AMPICILLIN			<=2	R																																																																																					
CEFAZOLIN			<=4	R																																																																																					
CEFTAZIDIME			<=1	S																																																																																					
CEFTRIAXONE			<=1	S																																																																																					
CIPROFLOXACIN			<=0.25	S																																																																																					
CLINDAMYCIN	<=0.25	S																																																																																							
ERYTHROMYCIN	<=0.25	S																																																																																							
GENTAMICIN			<=1	S																																																																																					
ERTAPENEM			<=0.5	S																																																																																					
OXACILLIN	<=0.25	S																																																																																							
PENICILLIN G	>=0.5	R																																																																																							
TOBRAMYCIN			<=1	S																																																																																					
VANCOMYCIN	<=0.5	S																																																																																							
PIPERA/TAZOACT			<=4	S																																																																																					

Disclaimer

This resource was created by Royal Victoria Regional Health Centre. PHO is not the owner of this content and does not take responsibility for the information provided within this document. Neither PHO nor Royal Victoria Regional Health Centre shall be responsible for the subsequent use of any tools and resources by any third party.

Example 2: Royal Victoria Regional Health Centre - Pharmacy/Microbiology Report (continued)



DATE: 20/10/15 @ 1615		Royal Victoria Pharmacy **LIVE**		PAGE 10
USER: MCAL		Microbiology Report		
Patient	[REDACTED]	Responsible Doctor	[REDACTED]	
Account Number	[REDACTED]	Location	Unit Number	[REDACTED]
Age/Sex	[REDACTED]	Room	Registered Date	[REDACTED]
Status	[REDACTED]	Bed	Discharged Date	[REDACTED]

Wound culture Final (continued)
 Providencia rettgeri: AST-N219
 This organism may develop resistance during therapy with B-lactam agents. Retesting in 3-4 days is recommended.

1. Staphylococcus aureus

Target	Route	Dose	RX	AB	Cost	M.I.C.	IQ
TRIMETH/SULFA			S			<=10	
CLINDAMYCIN			S			<=0.25	
ERYTHROMYCIN			S			<=0.25	
OXACILLIN			S			<=0.25	
PENICILLIN G			R			>=0.5	
VANCOMYCIN			S			<=0.5	

2. Providencia rettgeri

Target	Route	Dose	RX	AB	Cost	M.I.C.	IQ
TRIMETH/SULFA			S			<=20	
AMPICILLIN			R			<=2	
CEFAZOLIN			R			<=4	
CEFTAZIDIME			S			<=1	
CEFTRIAXONE			S			<=1	
CIPROFLOXACIN			S			<=0.25	
GENTAMICIN			S			<=1	
ERTAPENEM			S			<=0.5	
TOBRAMYCIN			S			<=1	
PIPERA/TAZOACT			S			<=4	

This organism may develop resistance during therapy with B-lactam agents. Retesting in 3-4 days is recommended.

Specimen:	Collected: 18/10/15-UNK	Status: COMP	Req#:
	Received: 18/10/15-1353	Source: NARES	Sp Desc: BOTH NARES

> **MRSA screen** Final
 NOT isolated.

Specimen:	Collected: 18/10/15-UNK	Status: COMP	Req#:
	Received: 18/10/15-1216	Source: RECTUM	Sp Desc:

> **MRSA screen** Final
 NOT isolated.

> **VRE screen** Final
 NOT isolated.

> **ESBL screen** Final
 NOT isolated.


Disclaimer

This resource was created by Royal Victoria Regional Health Centre. PHO is not the owner of this content and does not take responsibility for the information provided within this document. Neither PHO nor Royal Victoria Regional Health Centre shall be responsible for the subsequent use of any tools and resources by any third party.


Example 3: The Scarborough Hospital - ICNet System Sample Automated Alerts



Sample Automated Email Alert for +ve cultures



Other alerts (browser view)



Disclaimer

This resource was created by The Scarborough Hospital. PHO is not the owner of this content and does not take responsibility for the information provided within this document. Neither PHO nor The Scarborough Hospital shall be responsible for the subsequent use of any tools and resources by any third party.