

Humpback whales: Northern Atlantic

POPULATION SIZE, MIGRATION, DIVERGENCE, ASSIGNMENT, HISTORY

Bayesian inference using the structured coalescent

Migrate-n version 5.0.3a [January-08-2018]

Using Intel AVX (Advanced Vector Extensions)

Compiled for PARALLEL computer architectures

One master and 10 compute nodes are available.

Program started at Sun Oct 28 09:28:20 2018

Program finished at Sun Oct 28 10:14:08 2018 [Runtime:0000:00:45:48]



Options

Inheritance scalers in use for Thetas:

All loci use an inheritance scaler of 1.0

[The locus with a scaler of 1.0 used as reference]

Data set was subsampled: used a random sample of size:

20 and seed 163

Random number seed:

(with internal timer) 753380850

Start parameters:

Theta values were generated

Using a percent value of the prior

M values were generated

Using a percent value of the prior

Connection matrix:

m = average (average over a group of Thetas or M,

s = symmetric migration M, S = symmetric 4Nm,

0 = zero, and not estimated,

* = migration free to vary, Thetas are on diagonal

d = row population split off column population, D = split and then migration

Population 1

1 north_atlantic *

Order of parameters:

1 Θ_1 <displayed>

Mutation rate among loci:

Mutation rate is constant

Analysis strategy:

Bayesian inference

-Population size estimation:

Exponential Distribution

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling
M	Metropolis sampling
Divergence	Metropolis sampling
Divergence Spread	Metropolis sampling
Genealogy	Metropolis-Hastings

Prior distribution for parameter

Parameter	Prior	Minimum	Mean	Maximum	Delta	Bins	UpdateFreq
1 Theta	-1 Exponential	0.000000	0.020	0.900	-	1500	0.20000

[-1 -1 means priors were set globally]

Markov chain settings:

Long chain

Number of chains	1
Recorded steps [a]	10000
Increment (record every x step [b])	1000
Number of concurrent chains (replicates) [c]	10
Visited (sampled) parameter values [a*b*c]	100000000
Number of discard trees per chain (burn-in)	5000

Multiple Markov chains:

Static heating scheme

4 chains with temperatures
1000000.00 3.00 1.50 1.00
Swapping interval is 1

Print options:

Data file: infile.humpback_northern
 Haplotyping is turned on: NO
 Output file: outfile_humpbacknorth_0.6
 Posterior distribution raw histogram file: bayesfile
 Raw data from the MCMC run: bayesallfile_humpbacknorth_0.6
 Print data: No
 Print genealogies [only some for some data type]: None

Data summary

Data file: infile.humpback_northern
 Datatype: Haplotype data
 Number of loci: 1

Mutationmodel:

Locus	Sublocus	Mutationmodel	Mutationmodel parameters
1	1	Felsenstein 84	[Bf:0.33 0.19 0.13 0.35, t/t ratio=18.080]

Data set was subsampled: used a random sample of size: 20

Sites per locus

Locus	Sites
1	283

Site rate variation and probabilities:

Locus	Sublocus	Region type	Rate of change	Probability	Patch size
-------	----------	-------------	----------------	-------------	------------

1	1	1	0.418	0.706	1.000
1	1	2	2.260	0.283	1.000
1	1	3	6.161	0.011	1.000

Population	Locus	Gene copies data	(missing)
1 north_atlantic	1	188	
Total of all populations	1	188	(0)

Subsampled dataset

Data set was subsampled randomly per population: 20 samples taken

Locus	Population	Individuals
-------	------------	-------------

1	north_atla	MMASKLIKE 13pals01 04pals04 17pals23 17pals04 28pals02 02pals14 17pals11 17pals20 27pals01 26pals07 MTORCH 23pals19 MCLIPPER DR15 DR13 02pals03 Ice02 07pals01 02pals10
---	------------	---


Bayesian Analysis: Posterior distribution table

Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	Θ_1	0.04380	0.06780	0.07770	0.08760	0.12000	0.08250	0.08458

Citation suggestions:

Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters.
Bioinformatics 22:341-345

Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use?
In Population Genetics for Animal Conservation, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli,
and C. Vernesi, eds., vol. 17 of Conservation Biology, Cambridge University Press, Cambridge UK, pp. 42-79.

Bayesian Analysis: Posterior distribution for locus 1

Log-Probability of the data given the model (marginal likelihood)

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel}))]$

or as $LBF = 2 (\ln(\text{Prob}(D \mid \text{thisModel}) - \ln(\text{Prob}(D \mid \text{otherModel})))$

shows the support for thisModel]

Locus	TI(1a)	BTI(1b)	SS(2)	HS(3)
1	-643.71	-542.41	-391.42	-526.61

(1a) TI: Thermodynamic integration: $\log(\text{Prob}(D|\text{Model}))$: Good approximation with many temperatures

(1b) BTI: Bezier-approximated Thermodynamic integration: when using few temperatures USE THIS!

(2) SS: Steppingstone Sampling (Xie et al 2011)

(3) HS: Harmonic mean approximation: Overestimates the marginal likelihood, poor variance

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, *Genetics*, 185: 313-326.

Palczewski M. and P. Beerli, 2014. Population model comparison using multi-locus datasets.

In M.-H. Chen, L. Kuo, and P. O. Lewis, editors, *Bayesian Phylogenetics: Methods, Algorithms, and Applications*, pages 187-200. CRC Press, 2014.

Xie W., P. O. Lewis, Y. Fan, L. Kuo, and M.-H. Chen. 2011. Improving marginal likelihood estimation for Bayesian phylogenetic model selection. *Systematic Biology*, 60(2):150â 160, 2011.

Acceptance ratios for all parameters and the genealogies

Parameter	Accepted changes	Ratio
Θ_1	1017824/20002866	0.05088
Genealogies	8738664/79997134	0.10924

Average temperatures during the run

Chain	Temperatures
-------	--------------

1	0.00000
2	0.00000
3	0.00000
4	0.00000

Adaptive heating often fails, if the average temperatures are very close together
try to rerun using static heating! If you want to compare models using marginal
likelihoods then you **MUST** use static heating

Potential Problems

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is very common that some parameters for some loci will not be very informative, triggering suggestions (for example to increase the prior range) that are not sensible. This suggestion tool will improve with time, therefore do not blindly follow its suggestions. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. For example, if you run a Bayesian inference with sequence data, for macroscopic species there is rarely the need to increase the prior for Theta beyond 0.1; but if you use microsatellites it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3) it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the range will not help in such situations, reducing number of parameters may help in such situations.

No warning was recorded during the run