


two (fake) Swiss towns

POPULATION SIZE, MIGRATION, DIVERGENCE, ASSIGNMENT, HISTORY
Bayesian inference using the structured coalescent
Migrate-n version 5.0.7 [May-01-2025]
Compiled for PARALLEL computer architectures
One master and 4 compute nodes are available.
Program started at Mon Jul 28 10:59:57 2025
Program finished at Mon Jul 28 11:00:49 2025 [Runtime:0000:00:00:52]



Options

Datatype: DNA sequence data

Inheritance multipliers in use for Thetas:
All loci use an inheritance multiplier of 1.0

Random number seed: (with internal timer) 2159615964

Start parameters:

Theta values were generated Using a percent value of the prior

M values were generated Using a percent value of the prior

Connection matrix:
m = average (average over a group of Thetas or M,
s = symmetric migration M, S = symmetric 4Nm,
0 = zero, and not estimated,
* = migration free to vary, Thetas are on diagonal
d = row population split off column population, D = split and then migration

Population	1	2
1 Aadorf	*	*
2 Bern	*	*

Order of parameters:
1 Θ_1 <displayed>

2	Θ_2	<displayed>
3	$M_{2 \rightarrow 1}$	<displayed>
4	$M_{1 \rightarrow 2}$	<displayed>

Mutation rate among loci:

Mutation rate is constant for all loci

Analysis strategy:

Bayesian inference

-Population size estimation:

Exponential Distribution

-Geneflow estimation:

Exponential Distribution

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling
M	Metropolis sampling
Genealogy	Metropolis-Hastings

Prior distribution for parameter

Parameter		Prior	Minimum	Mean*	Maximum	Delta	Bins	UpdateFreq
1 Theta	* *	Exponential	0.000000	0.010	0.100	-	1500	0.10417
2 Theta	* *	Exponential	0.000000	0.010	0.100	-	1500	0.10417
3 M	2 1	Uniform	0.000000	100.0	1000.	100.00000	2000	0.10417
4 M	1 2	Uniform	0.000000	100.0	1000.	100.00000	2000	0.10417

[* * means priors were set globally]

Posterior distribution:

Parameter values were collected using MCMC, these values

were then used to generate the posterior histograms using KERNEL SMOOTHING (window=41)

and subsequent SAVITZKY-GOLAY SMOOTHING (window=41) for combination over loci

Markov chain settings:

Long chain

Number of chains	1
Recorded steps [a]	10000
Increment (record every x step [b])	50
Number of concurrent chains (replicates) [c]	1
Visited (sampled) parameter values [a*b*c]	500000
Number of discard trees per replicate (burn-in * b)	250000

Multiple Markov chains:

Static heating scheme

4 chains with temperatures
1000000.00 3.00 1.50 1.00
Swapping interval is 1

Print options:





Data file:	twoswisstowns
	parmfile.twoswisstowns
Haplotyping is turned on:	NO
Output file:	outfile-twoswisstowns
Posterior distribution raw histogram file:	bayesfile
Raw data from the MCMC run:	bayesallfile.txt
Print data:	No
Print genealogies [only some for some data type]:	None

Bayesian Analysis: Posterior distribution table

Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	Θ_1	0.00300	0.00753	0.01077	0.01453	0.02580	0.01237	0.01329
1	Θ_2	0.00600	0.01180	0.01577	0.02193	0.03947	0.01903	0.02074
1	$M_{2 \rightarrow 1}$	530.5	820.5	965.8	988.0	999.5	830.8	807.5
1	$M_{1 \rightarrow 2}$	567.5	824.5	957.2	984.5	999.5	836.8	818.1
2	Θ_1	0.00440	0.00993	0.01430	0.01933	0.03507	0.01663	0.01800
2	Θ_2	0.00707	0.01527	0.02050	0.02873	0.04900	0.02457	0.02640
2	$M_{2 \rightarrow 1}$	576.5	828.0	965.2	985.5	999.5	839.8	821.9
2	$M_{1 \rightarrow 2}$	564.0	838.5	952.8	984.0	999.5	829.8	812.6
3	Θ_1	0.00493	0.01020	0.01397	0.01940	0.03440	0.01683	0.01808
3	Θ_2	0.01020	0.01853	0.02517	0.03327	0.05553	0.02877	0.03058
3	$M_{2 \rightarrow 1}$	614.5	845.0	968.8	987.0	999.5	856.2	839.5
3	$M_{1 \rightarrow 2}$	601.5	832.0	960.2	983.5	999.5	845.2	828.8
All	Θ_1	0.00827	0.01247	0.01550	0.01867	0.02707	0.01650	0.01706
All	Θ_2	0.01587	0.02407	0.02830	0.03567	0.04680	0.03083	0.03164
All	$M_{2 \rightarrow 1}$	732.5	891.0	966.8	984.5	999.5	902.2	887.0
All	$M_{1 \rightarrow 2}$	724.0	880.0	956.2	982.5	999.5	892.8	878.8

Citation suggestions:

- Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 22:341-345
- Beerli, P., H. Ashki, S. Mashayekhi, and M. Palczewski, 2022. Population divergence time estimation using individual lineage label switching. *G3 Genes & Genomes & Genetics*, 12(4), 02 2022.
- Beerli P., 2007. Estimation of the population scaled mutation rate from microsatellite data, *Genetics*, 177:1967-1968.
- Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use? In *Population Genetics for Animal Conservation*, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli, and C. Vernesi, eds., vol. 17 of *Conservation Biology*, Cambridge University Press, Cambridge UK, pp. 42-79.
- Beerli, P., S. Mashayekhi, M. Sadeghi, M. Khodaei, and K. Shaw, 2019. Population genetic inference with migrate. *Current Protocols in Bioinformatics*, 68(1):e87.

Bayesian Analysis: Posterior distribution over all loci

Log-Probability of the data given the model (marginal likelihood)

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D \mid \text{thisModel})) - \ln(\text{Prob}(D \mid \text{otherModel}))]$

or as $LBF = 2 (\ln(\text{Prob}(D \mid \text{thisModel})) - \ln(\text{Prob}(D \mid \text{otherModel})))$

shows the support for thisModel]

Locus	TI(1a)	BTI(1b)	HS(3)
1	-2612.85	-2308.70	-2406.18
2	-1432.95	-1257.42	-1327.26
3	-1579.35	-1351.29	-1409.90
All	-5637.72	-4929.99	-5155.91

(1a) TI: Thermodynamic integration: $\log(\text{Prob}(D \mid \text{Model}))$: Good approximation with many temperatures

(1b) BTI: Bezier-approximated Thermodynamic integration: when using few temperatures USE THIS!

(3) HS: Harmonic mean approximation: Overestimates the marginal likelihood, poor variance

[Scaling factor = -12.571520]

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, *Genetics*, 185: 313-326.

Palczewski M. and P. Beerli, 2014. Population model comparison using multi-locus datasets.

In M.-H. Chen, L. Kuo, and P. O. Lewis, editors, *Bayesian Phylogenetics: Methods, Algorithms, and Applications*, pages 187-200. CRC Press, 2014.

Acceptance ratios for all parameters and the genealogies

Parameter	Accepted changes	Ratio
Θ_1	41436/156474	0.26481
Θ_2	29976/156408	0.19165
$M_{2 \rightarrow 1}$	68723/155975	0.44060
$M_{1 \rightarrow 2}$	87917/156281	0.56256
Genealogies	69568/156281	0.44515

MCMC-Autocorrelation and Effective MCMC Sample Size

Parameter	Autocorrelation	Effective Sample Size
Θ_1	0.57061	8222.44
Θ_2	0.64277	6661.98
$M_{2 \rightarrow 1}$	0.56188	8417.76
$M_{1 \rightarrow 2}$	0.56272	8417.23
Genealogies	0.57061	8222.44

Static temperatures during the run using standard heating

Chain	Temperatures	log(marginal likelihood)	log(mL_steppingstone)
1	1.00000	-1582.68860	365.85794
2	1.00000	-1599.69025	106.72915
3	1.00000	-1628.61981	112.74756
4	1.00000	-3211.01132	6.65434

Potential Problems

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is common that some parameters for some loci will not be informative. These parameters then trigger suggestions to increase the prior range that are not sensible. Do not blindly follow the suggestions given. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. Suppose you run a Bayesian inference with sequence data for macroscopic species. In that case, there is rarely the need to increase the prior for Theta beyond 0.1. If you use microsatellites data, it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3), it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the prior range will not help in such situations, but reducing the number of parameters may help.

Param 3 (Locus 1): Upper prior boundary seems too low!
 Param 4 (Locus 1): Upper prior boundary seems too low!
 Param 3 (Locus 2): Upper prior boundary seems too low!
 Param 4 (Locus 2): Upper prior boundary seems too low!
 Param 3 (Locus 3): Upper prior boundary seems too low!
 Param 4 (Locus 3): Upper prior boundary seems too low!
 Param 3 (all loci): Upper prior boundary seems too low!
 Param 4 (all loci): Upper prior boundary seems too low!

Summary Assignment of Individuals to Populations

Individual	Population	
	1	2
?BAH0	0.326	0.674
?BAF0	0.343	0.657
?BAG1	0.363	0.637
?BAJ1	0.336	0.664
?BAH1	0.382	0.618
?BAI1	0.381	0.619
?BAF1	0.342	0.658

Detailed Assignment of Individuals to Populations

Individual	Locus	Population	
		1	2
?BAH0	1	0.452	0.548
?BAH0	2	0.434	0.566
?BAH0	3	0.434	0.566
?BAH0	All	0.326	0.674
?BAF0	1	0.431	0.569
?BAF0	2	0.462	0.538
?BAF0	3	0.446	0.554
?BAF0	All	0.343	0.657
?BAG1	1	0.440	0.560
?BAG1	2	0.458	0.542
?BAG1	3	0.461	0.539
?BAG1	All	0.363	0.637
?BAJ1	1	0.408	0.592
?BAJ1	2	0.450	0.550
?BAJ1	3	0.473	0.527
?BAJ1	All	0.336	0.664
?BAH1	1	0.438	0.562
?BAH1	2	0.468	0.532
?BAH1	3	0.474	0.526
?BAH1	All	0.382	0.618
?BAI1	1	0.479	0.521
?BAI1	2	0.447	0.553
?BAI1	3	0.454	0.546
?BAI1	All	0.381	0.619
?BAF1	1	0.446	0.554
?BAF1	2	0.458	0.542
?BAF1	3	0.433	0.567
?BAF1	All	0.342	0.658