


Japan

POPULATION SIZE, MIGRATION, DIVERGENCE, ASSIGNMENT, HISTORY
Bayesian inference using the structured coalescent
Migrate-n version 5.0.7 [May-01-2025]
Compiled for PARALLEL computer architectures
One master and 9 compute nodes are available.
Program started at Mon Jul 28 11:01:58 2025
Program finished at Mon Jul 28 11:02:58 2025 [Runtime:0000:00:01:00]



Options

Datatype: Single nucleotide polymorphism data(Hapmap formatting)

Inheritance multipliers in use for Thetas:
All loci use an inheritance multiplier of 1.0

Random number seed: (with internal timer) 3632703597

Start parameters:

Theta values were generated Using a percent value of the prior

M values were generated Using a percent value of the prior

Connection matrix:
m = average (average over a group of Thetas or M,
s = symmetric migration M, S = symmetric 4Nm,
0 = zero, and not estimated,
* = migration free to vary, Thetas are on diagonal
d = row population split off column population, D = split and then migration

Population 1
1 Japan *

Order of parameters:
1 Θ_1 <displayed>

Mutation rate among loci: Mutation rate is constant for all loci

Analysis strategy: Bayesian inference

-Population size estimation: Exponential Distribution

Proposal distributions for parameter

Parameter	Proposal
Theta	Metropolis sampling

M	Metropolis sampling
---	---------------------

Genealogy	Metropolis-Hastings
-----------	---------------------

Prior distribution for parameter

Parameter	Prior	Minimum	Mean*	Maximum	Delta	Bins	UpdateFreq
1 Theta * *	Uniform	0.000000	0.050	0.100	0.010000	1500	0.50000

[* * means priors were set globally]

Posterior distribution:

Parameter values were collected using MCMC, these values

were then used to generate the posterior histograms using KERNEL SMOOTHING (window=41)

and subsequent SAVITZKY-GOLAY SMOOTHING (window=41) for combination over loci

Markov chain settings: Long chain

Number of chains 1

Recorded steps [a] 10000

Increment (record every x step [b] 100

Number of concurrent chains (replicates) [c] 1

Visited (sampled) parameter values [a*b*c] 1000000

Number of discard trees per replicate (burn-in * b) 100000

Multiple Markov chains:

Static heating scheme 4 chains with temperatures

1000000.00 3.00 1.50 1.00

Swapping interval is 1

Print options:

Data file: infile.hapmap

parmfile.hapmap

Haplotyping is turned on: NO


Output file: outfile_hapmap

Posterior distribution raw histogram file: bayesfile

Raw data from the MCMC run: bayesallfile.txt

Print data: No

Print genealogies [only some for some data type]: None




Bayesian Analysis: Posterior distribution table

Locus	Parameter	2.5%	25.0%	Mode	75.0%	97.5%	Median	Mean
1	Θ_1	0.04560	0.06693	0.08050	0.09047	0.09980	0.07450	0.07348
2	Θ_1	0.03040	0.05067	0.05977	0.07767	0.09673	0.06323	0.06301
3	Θ_1	0.05200	0.07520	0.08463	0.09440	0.09993	0.07957	0.07815
4	Θ_1	0.05140	0.07713	0.08830	0.09600	0.09993	0.08043	0.07869
5	Θ_1	0.05060	0.07587	0.08183	0.09587	0.09993	0.07910	0.07755
6	Θ_1	0.05067	0.07207	0.08177	0.09240	0.09993	0.07817	0.07696
7	Θ_1	0.02787	0.03993	0.05390	0.06600	0.09153	0.05697	0.05804
8	Θ_1	0.05393	0.07453	0.08710	0.09393	0.09993	0.07950	0.07851
9	Θ_1	0.04787	0.07713	0.08950	0.09747	0.09993	0.07910	0.07727
All	Θ_1	0.06613	0.07453	0.07957	0.08340	0.09313	0.07930	0.07934

Citation suggestions:

- Beerli P., 2006. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 22:341-345
- Beerli, P., H. Ashki, S. Mashayekhi, and M. Palczewski, 2022. Population divergence time estimation using individual lineage label switching. *G3 Genes & Genomes & Genetics*, 12(4), 02 2022.
- Beerli P., 2009. How to use MIGRATE or why are Markov chain Monte Carlo programs difficult to use? In *Population Genetics for Animal Conservation*, G. Bertorelle, M. W. Bruford, H. C. Hauffe, A. Rizzoli, and C. Vernesi, eds., vol. 17 of *Conservation Biology*, Cambridge University Press, Cambridge UK, pp. 42-79.
- Beerli, P., S. Mashayekhi, M. Sadeghi, M. Khodaei, and K. Shaw, 2019. Population genetic inference with migrate. *Current Protocols in Bioinformatics*, 68(1):e87.

Bayesian Analysis: Posterior distribution over all loci



Log-Probability of the data given the model (marginal likelihood)

Use this value for Bayes factor calculations:

$BF = \text{Exp}[\ln(\text{Prob}(D \mid \text{thisModel})) - \ln(\text{Prob}(D \mid \text{otherModel}))]$

or as $LBF = 2 (\ln(\text{Prob}(D \mid \text{thisModel})) - \ln(\text{Prob}(D \mid \text{otherModel})))$

shows the support for thisModel]

Locus	TI(1a)	BTI(1b)	HS(3)
1	-61.46	-39.72	-15.36
2	-8.15	-7.99	-8.09
3	-43.45	-35.75	-19.17
4	-33.84	-30.97	-17.33
5	-16.67	-16.37	-14.47
6	-16.55	-16.23	-14.73
7	-0.44	-0.43	-0.48
8	-24.36	-22.90	-14.52
9	-28.67	-26.71	-19.64
All	-238.03	-201.52	-128.24

(1a) TI: Thermodynamic integration: $\log(\text{Prob}(D \mid \text{Model}))$: Good approximation with many temperatures

(1b) BTI: Bezier-approximated Thermodynamic integration: when using few temperatures USE THIS!

(3) HS: Harmonic mean approximation: Overestimates the marginal likelihood, poor variance

[Scaling factor = -4.450263]

Citation suggestions:

Beerli P. and M. Palczewski, 2010. Unified framework to evaluate panmixia and migration direction among multiple sampling locations, *Genetics*, 185: 313-326.

Palczewski M. and P. Beerli, 2014. Population model comparison using multi-locus datasets.

In M.-H. Chen, L. Kuo, and P. O. Lewis, editors, *Bayesian Phylogenetics: Methods, Algorithms, and Applications*, pages 187-200. CRC Press, 2014.

Acceptance ratios for all parameters and the genealogies

Parameter	Accepted changes	Ratio
Θ_1	2180530/4503174	0.48422
Genealogies	3834602/4503174	0.85153

MCMC-Autocorrelation and Effective MCMC Sample Size

Parameter	Autocorrelation	Effective Sampe Size
Θ_1	0.65057	19127.73
Genealogies	0.65057	19127.73

Static temperatures during the run using standard heating

Chain	Temperatures	log(marginal likelihood)	log(mL_steppingstone)
1	1.00000	-5.97065	313.95272
2	1.00000	-10.24909	206.89874
3	1.00000	-31.17826	108.03103
4	1.00000	-66.89428	15.52049

Potential Problems

This section reports potential problems with your run, but such reporting is often not very accurate. With many parameters in a multilocus analysis, it is common that some parameters for some loci will not be informative. These parameters then trigger suggestions to increase the prior range that are not sensible. Do not blindly follow the suggestions given. If some parameters are flagged, inspect the tables carefully and judge whether an action is required. Suppose you run a Bayesian inference with sequence data for macroscopic species. In that case, there is rarely the need to increase the prior for Theta beyond 0.1. If you use microsatellites data, it is rather common that your prior distribution for Theta should have a range from 0.0 to 100 or more. With many populations (>3), it is also very common that some migration routes are estimated poorly because the data contains little or no information for that route. Increasing the prior range will not help in such situations, but reducing the number of parameters may help.

Param 1 (Locus 1): Upper prior boundary seems too low!
Param 1 (Locus 3): Upper prior boundary seems too low!
Param 1 (Locus 4): Upper prior boundary seems too low!
Param 1 (Locus 5): Upper prior boundary seems too low!
Param 1 (Locus 6): Upper prior boundary seems too low!
Param 1 (Locus 8): Upper prior boundary seems too low!
Param 1 (Locus 9): Upper prior boundary seems too low!