

The LMTO object of the CP-PAW code

Peter E. Blöchl

Copyright Peter E. Blöchl; Sept.2, 2013-May 22, 2019
Institute of Theoretical Physics; Clausthal University of Technology;
D-38678 Clausthal Zellerfeld; Germany;
<http://www.pt.tu-clausthal.de/atp/>

Contents

1	Todo	2
1.1	Fixes	2
1.2	Ideas	2
2	Purpose and theoretical background of the LMTO Object	4
2.1	Augmentation	4
2.2	Structure constants	5
2.2.1	Hankel functions as envelope function	5
2.2.2	Hankel and Bessel functions as head and tail functions	5
2.2.3	Bare structure constants	6
2.2.4	Screened structure constants	6
2.3	Screening on finite clusters	8
2.4	Derivatives	9
2.4.1	Lower and higher angular momenta	9
2.5	Augmentation and Potential parameters	10
2.5.1	Local orbitals	10
2.6	Coefficients of the tight-binding orbital	11
2.6.1	Introduction	11
2.6.2	Transformation between local-orbital and partial-wave projections	12
2.7	Tailed representation of the natural tight-binding orbitals	14
2.8	How to choose the parameters	14
2.8.1	Augmentation radius must be large for semi-core states	14
3	Contributions to the Hamiltonian	15
3.1	Core-valence exchange	15
3.2	U-tensor	16
3.3	Double-counting correction	16
3.3.1	HSE-like	16
3.3.2	Blöchl, Walther, Pruschke	17
4	Description of Subroutines	19
4.1	Workflow	19

4.2	LMTO_POTPAR	19
4.3	LMTO\$CLUSTERSTRUCTURECONSTANTS	20
4.3.1	LMTO\$STRUCTURECONSTANTS	21
4.3.2	LMTO\$SCREEN	21
4.4	Waves object	22
4.5	Offsite matrix elements	23
4.6	Matrix elements using Gaussians	24
4.7	Matrix elements on an adaptive grid	24
4.8	Routines for reporting	24
4.9	Routines for plotting orbitals	25
5	Benchmarks	27
5.1	Silicon	27
5.1.1	Summary	29
6	Implementation of simpler orbitals	30
6.1	Local orbitals	30
6.2	Projector functions	31
6.2.1	Fitting	31
6.3	Expressions	31
6.3.1	Linear approximation in structure constants	32
6.4	Failed procedure	32
6.5	Code structure	33
6.5.1	Local orbitals	34
6.5.2	Projections	34
6.6	Transformation of density matrix and Hamiltonian	35
6.6.1	Formally correct derivation	35
6.6.2	Coding of direct inversion	36
6.6.3	Approximate inversion	36
6.6.4	Otherstuff	40
6.7	Analysis of the code structure in paw_lmto	41
6.8	New code	41
6.8.1	Determine off-site density matrix in partial-wave representation	41
6.8.2	Density matrix in orbital representation	42
6.8.3	Bare structure constants	42
6.8.4	Routine radial\$helmholtz	42
6.9	Helmholtz equation	43
7	Off-site matrix elements NDDO	47
7.1	Exchange energy and U-tensor	47
7.1.1	Division of exchange energy	47

7.1.2	Interaction	48
7.2	Long-range expansions	49
7.2.1	Long-range expansion of the potential	49
7.2.2	Long-range expansion of the energy	50
7.2.3	Division of U-tensor into short- and long-ranged parts	51
7.3	Expansion of the density in angular momenta	53
7.3.1	Tabulating data	57
7.3.2	Integrals with the point density	58
7.3.3	Tests	60
7.3.4	Complete long-range expansion	60
A	Definition of solid Hankel functions	62
A.1	Modified spherical Bessel functions	63
A.2	Bare structure constants	64
A.3	Consistency checks	65
B	Bloch theorem revisited	68
C	Offsite matrix elements using Gaussian integrals	70
D	Double counting	71
D.1	Other double-counting schemes	71
D.1.1	Around mean field (AMF)	73
D.1.2	Definition of U and J parameters	76
E	Changelog, Bugfixes	77

Chapter 1

Todo

1.1 Fixes

- `Imto__overlapphi` calculates the onsite overlap matrix of partial waves in a sphere.
- using only sp like tight-binding orbitals and local exchange lead to an increase of the band gap of silicon above 1.3 eV. After adding the d-orbitals to the HF term collapsed the band gap dramatically below the dft value. Core-valence exchange seems to have an important effect on the band gap too.
- The charge sumrule is not correct! The calculation for an H-atom yields $Tr[\rho O] \approx 0.2$. The problem is not the difference between tailed orbitals and the multicenter expansion, because $(r * \chi)^2$ agrees quite well.

1.2 Ideas

- With the introduction of the tailed representation of the NTBO's we **switch from a multicenter expansion to a one-center expansion**. This, however, requires to increase the number of angular momenta for the tail functions beyond that used for the augmentation: Each (screening) Hankel function centered on the neighboring site contributes arbitrary many angular momenta at the central site.

The tail functions for the higher angular momenta behave, at the central site, like a bare Bessel function. Beyond the central site, we can add the same pairs of exponential tails as for the lower tailed partial waves.

(The following is probably no more true. please check!) Currently we use the following rule: Rule: (1) Each head function has exactly one tail function attributed to it. Thus we can identify the phidot function uniquely by looking for the tail function with the correct angular momentum. From (1) follows, that there is at most one tail function per site and angular momentum.

- Suggestion: the exponential functions for the tailed expansion are determined by trial and error. A more systematic approach for the tail function would be to (1) build a Hankel function on a neighboring site, which is augmented by (a) a bare Bessel function and (b) by

a screened scattering partial wave $|\dot{\vec{\phi}}\rangle$. (1a) In order to make the formulation less dependent on the neighbor distance, we could also include the gradients of these functions. (2) These functions are expanded about the central site into an angular-momentum expansion. (3) a reasonably weighted least-square fit of the exponential tails to these functions will provide a “optimum shape” of the tails for each angular momentum.

- The orbitals in the tailed one-center expansion and as multicenter expansions can be compared using `LMT0_PLOTORBS`. Calculations for hydrogen suggest that a good choice is $K2 = -1$, and $(\lambda_1, \lambda_2) = (4, 2)$. The rationale for $K2$ is that the energy of the hydrogen orbital is at $-1Ry = -\frac{1}{2}H = \frac{1}{2}K2$. (see [Dummy/Sitest/Test/Occtest/H2test](#))

Chapter 2

Purpose and theoretical background of the LMTO Object

The LMTO object maps the wave functions expressed in augmented plane waves onto a basiset of **natural tight-binding orbitals**. The natural tight-binding orbitals are a kind of LMTO's, screened such that the tails exhibit only scattering character in the context of nodeless wave functions[?].

2.1 Augmentation

The concept of linear augmented waves[?] is as follows:

1. At first, a so-called **envelope function** $|K_\alpha^\infty\rangle$ is defined.¹
2. In a second step, this envelope function is expanded about each atomic site R_α into spherical harmonics. More generally, they are expanded into **head functions** $|K_\alpha^\Omega\rangle$ and **tail functions** $|J_\alpha^\Omega\rangle$. The head function is the dominant contribution and carries the quantum number of the final orbital, while the tail functions are the minor contributions with different quantum numbers. In practice, the head functions are solid Hankel functions and the tail functions are solid Bessel functions.

$$|K_\alpha^\infty\rangle = |K_\alpha^\Omega\rangle - \sum_\beta |J_\beta^\Omega\rangle S_{\beta,\alpha}^\dagger + |K_\alpha^I\rangle \quad (2.1)$$

The coefficients $S_{\alpha,\beta}$ of the tail functions are called **structure constants**.

The difference between the full envelope function and its expansion into head and tail functions is the **interstitial envelope function**² $|K_\alpha^I\rangle$.

¹The superscript ∞ denotes that the function extends over all space, a superscript Ω denotes that the function is truncated (set to zero) outside the augmentation sphere Ω_R centered at the site R_α denoted by the index α . The superscript I denotes that the function is non-zero only in the interstitial region, that is outside all augmentation spheres. If the augmentation spheres overlap, the function in the interstitial region is defined by subtraction of all sphere contributions. Similarly, the interstitial function contains terms from the higher angular momenta within the augmentation regions.

²The interstitial envelope function is confined mostly in the region in between the atoms, but it also accounts for the overlap of the atomic regions and so-called higher partial waves not taken care of in the regular partial-wave expansion.

3. In the third step, the head and tail functions are replaced differentiably at some sphere radius by **partial waves** of the atomic potential. For that purpose, we use as partial waves a solution of the Schrödinger equation for some energy, denoted as $|\phi_\alpha\rangle$ and its energy derivative function $|\dot{\phi}_\alpha\rangle$.

The matching parameters are called **potential parameters**.

2.2 Structure constants

2.2.1 Hankel functions as envelope function

In practice, we will use solid Hankel functions $H_L(\vec{r})$ as envelope functions, so that

$$\langle \vec{r} | K_\alpha^\infty \rangle = H_{L_\alpha}(\vec{r} - \vec{R}_\alpha) \quad (2.2)$$

Solid Hankel functions are irregular solutions of the the inhomogeneous Helmholtz equation³

$$[\vec{\nabla}^2 + k^2] H_L(\vec{r}) = -4\pi(-1)^\ell \mathcal{Y}(\vec{\nabla}) \delta(\vec{r}) \quad (2.3)$$

Here $\mathcal{Y}_L(\vec{r}) \stackrel{\text{def}}{=} r^\ell Y_L(\vec{r})$ is a polynomial. With a gradient as argument, it becomes a differential operator. With $Y_L(\vec{r})$ we denote a spherical or real harmonic function and $L \stackrel{\text{def}}{=} (\ell, m)$ is a composite index of angular momentum quantum number ℓ and magnetic quantum number m .

Further details about the Hankel and Bessel functions can be found in appendix A.

2.2.2 Hankel and Bessel functions as head and tail functions

Defining the envelope function via an isotropic and translationally invariant differential equation of second order has the advantage that the solution can be expanded about different centers into regular solutions of the same differential equation with specific angular momenta. The regular solutions of the Helmholtz equation are the Bessel functions.

Hankel and Bessel functions are defined⁴ so that they behave at the origin as

$$K_{R,L}^\Omega(\vec{r}) = \left[(2\ell - 1)!! \frac{1}{|\vec{r} - \vec{R}|^{\ell+1}} + \dots \right] Y_L(\vec{r} - \vec{R}) \theta_{\Omega_R}(\vec{r}) \quad (2.4)$$

$$J_{R,L}^\Omega(\vec{r}) = \left[\frac{1}{(2\ell + 1)!!} |\vec{r} - \vec{R}|^{\ell+1} + \dots \right] Y_L(\vec{r} - \vec{R}) \theta_{\Omega_R}(\vec{r}) \quad (2.5)$$

$\theta_{\Omega_R}(\vec{r})$ is a step function, which equals unity within the augmentation region Ω_R centered at site R , while it vanishes outside. The terms neglected are higher orders in $|\vec{r} - \vec{R}|$.

³I am not sure whether also the three dimensional differential equation or only the one-dimensional differential equation for the radial part is called Helmholtz equation.

⁴This is our definition, not a generally accepted convention.

2.2.3 Bare structure constants

The **bare structure constants** $S_{\beta,\alpha}^\dagger$ are the expansion constants for an off-center expansion of solid spherical Hankel functions $|K_\alpha^\infty\rangle$ into **solid Bessel functions** $|J_\beta^\Omega\rangle$.

$$|K_\alpha^\infty\rangle = |K_\alpha^\Omega\rangle - \sum_\beta |J_\beta^\Omega\rangle S_{\beta,\alpha}^\dagger + |K_\alpha^I\rangle \quad (2.6)$$

The index α denotes here an atomic site R and a set of angular momenta $L = (\ell, m)$.

The superscript ∞ denotes that the function extends over all space, a superscript Ω denotes that the function is truncated (set to zero) outside the augmentation sphere Ω_R centered at the site denoted by the index. The superscript I denotes that the function is limited to the interstitial region, that is, outside all augmentation spheres. If the augmentation spheres overlap, the function in the interstitial region is defined by subtraction of all sphere contributions.

BARE STRUCTURE CONSTANTS

The bare structure constants have the form

$$S_{RL,R'L'} = (-1)^{\ell'+1} 4\pi \sum_{L''} C_{L,L',L''} H_{L''}(\vec{R}' - \vec{R}) \begin{cases} (-ik)^{\ell+\ell'-\ell''} & \text{for } k^2 > 0 \\ \delta_{\ell+\ell'-\ell''} & \text{for } k^2 = 0 \\ \kappa^{\ell+\ell'-\ell''} & \text{for } k^2 = -\kappa^2 < 0 \end{cases} \quad (2.7)$$

With $C_{L,L',L''}$, we denote the **Gaunt coefficients** defined by

$$Y_{L'}(\vec{r}) Y_{L''}(\vec{r}) = \sum_L Y_L(\vec{r}) C_{L,L',L''} \quad (2.8)$$

Note, that the Gaunt coefficients for spherical and real spherical harmonics differ.⁵

The bare structure constants are hermitean⁶, i.e.

$$S_{RL,R'L'} = S_{R'L',RL} \quad (2.9)$$

This is, however, not true for each angular-momentum block individually, i.e. in general we have $S_{RL,R'L'} \neq S_{R,L',R'L}$.

2.2.4 Screened structure constants

A so-called **screened LMTO representation** is defined by a set of screened scattering partial waves $|\dot{\phi}_\alpha\rangle$. In what we call the nodeless-representation, the scattering partial waves are defined as nodeless scattering wave functions.

The node-less scattering partial wave $|\dot{\phi}_\alpha\rangle$ define the screening constants \bar{Q}_α such that the screened tail functions $|\bar{J}_\alpha\rangle$ match with value and derivative to the scattering partial wave

$$|\dot{\phi}_\alpha\rangle \rightarrow |\bar{J}_\alpha^\Omega\rangle \stackrel{\text{def}}{=} |J_\alpha^\Omega\rangle - |K_\alpha^\Omega\rangle \bar{Q}_\alpha \quad (2.10)$$

⁵In practice, we use real spherical harmonics and the corresponding Gaunt coefficients.

⁶We use that $H_L(\vec{r}) = (-1)^\ell H_L(-\vec{r})$ and that the Gaunt coefficients $C_{L,L',L''}$ vanish unless $\ell + \ell' + \ell''$ is even.

A screened solid Hankel function $|\bar{K}_\alpha^\infty\rangle$ is a superposition of bare solid Hankel functions on a set of atomic positions

$$|\bar{K}_\alpha^\infty\rangle = \sum_{\beta} |K_\beta^\infty\rangle c_{\beta,\alpha} \quad (2.11)$$

with the property that the tail functions are made entirely from screened Bessel functions $|\bar{J}_\beta^\Omega\rangle$, i.e.

$$|\bar{K}_\alpha^\infty\rangle = |K_\alpha^\Omega\rangle - \sum_{\beta} |\bar{J}_\beta^\Omega\rangle \bar{S}_{\beta,\alpha}^\dagger + |\bar{K}_\alpha^I\rangle \quad (2.12)$$

The expansion coefficients \bar{S} are the **screened structure constants**.

By equating the two expressions for the screened Hankel functions, namely Eq. 2.11 and Eq. 2.12, we can extract the screened structure constants $\bar{S}_{\beta,\alpha}^\dagger$ and the superposition coefficients $c_{\beta,\alpha}$.

$$\begin{aligned} \sum_{\beta} \left[|K_\beta^\Omega\rangle - \sum_{\gamma} |J_\gamma^\Omega\rangle S_{\gamma,\beta}^\dagger + |K_\beta^I\rangle \right] c_{\beta,\alpha} &= |K_\alpha^\Omega\rangle - \sum_{\beta} \underbrace{\left[|J_\beta^\Omega\rangle - |K_\beta^\Omega\rangle \bar{Q}_\beta \right]}_{|\bar{J}_\beta^\Omega\rangle} \bar{S}_{\beta,\alpha}^\dagger + |\bar{K}_\alpha^I\rangle \\ \stackrel{\text{Eq. 2.11}}{\Rightarrow} \sum_{\beta} |K_\beta^\Omega\rangle c_{\beta,\alpha} - \sum_{\beta,\gamma} |J_\gamma^\Omega\rangle S_{\gamma,\beta}^\dagger c_{\beta,\alpha} &= \sum_{\beta} |K_\beta^\Omega\rangle \left[\delta_{\beta,\alpha} + \bar{Q}_\beta \bar{S}_{\beta,\alpha}^\dagger \right] - \sum_{\beta} |J_\beta^\Omega\rangle \bar{S}_{\beta,\alpha}^\dagger \end{aligned} \quad (2.13)$$

By comparing the coefficients, we obtain

$$c_{\beta,\alpha} = \delta_{\beta,\alpha} + \bar{Q}_\beta \bar{S}_{\beta,\alpha}^\dagger \quad (2.14)$$

$$\bar{S}_{\gamma,\alpha}^\dagger = \sum_{\beta} S_{\gamma,\beta}^\dagger c_{\beta,\alpha} \quad (2.15)$$

which can be resolved to ⁷ the defining equation of the screened structure constants

SCREENED STRUCTURE CONSTANTS

$$\bar{S}^\dagger = \left[\mathbf{1} - \mathbf{s}^\dagger \bar{\mathbf{Q}} \right]^{-1} \mathbf{s}^\dagger \quad (2.17)$$

and the expression of the screened Hankel functions

$$|\bar{K}_\alpha^\infty\rangle = \sum_{\beta} |K_\beta^\infty\rangle \left[\delta_{\beta,\alpha} + \bar{Q}_\beta \bar{S}_{\beta,\alpha}^\dagger \right]. \quad (2.18)$$

Because we calculate the screened structure constants on finite clusters, Eq. 2.17 should be considered of only formal value and should not be used in the actual calculations. Rather, the defining equations Eq. 2.15 shall be used as shown in the following section.

$$\bar{S}^\dagger \stackrel{\text{Eq. 2.15}}{=} \mathbf{s}^\dagger \mathbf{c} \stackrel{\text{Eq. 2.14}}{=} \mathbf{s}^\dagger (\mathbf{1} + \bar{\mathbf{Q}} \bar{S}^\dagger) \Rightarrow (\mathbf{1} - \mathbf{s}^\dagger \bar{\mathbf{Q}}) \bar{S}^\dagger = \mathbf{s}^\dagger \Rightarrow \bar{S}^\dagger = (\mathbf{1} - \mathbf{s}^\dagger \bar{\mathbf{Q}})^{-1} \mathbf{s}^\dagger \quad (2.16)$$

2.3 Screening on finite clusters

The screened structure constants are calculated on a cluster of atomic sites. The calculation can, in principle, be done for each single screened Hankel function independently. In practice, we do the calculations for all atoms centered on a given site in one step.

We go back to the defining equation system Eq. 2.14, Eq. 2.15 and rewrite it in terms of vectors, which are defined on the cluster B . The index α labeling the vectors counts the envelope functions centered at the central site. The dimension of the vectors and matrices defined below is given by angular momentum channels and sites with non-zero screening charge⁸ \bar{Q} .

The equations attain the form

$$\vec{c}_\alpha \stackrel{\text{Eq. 2.14}}{=} \vec{e}_\alpha + \bar{Q} \vec{s}_\alpha \quad (2.19)$$

$$\vec{s}_\alpha \stackrel{\text{Eq. 2.15}}{=} \mathbf{S}^\dagger \vec{c}_\alpha \quad (2.20)$$

where the vectors \vec{c}_α , \vec{s}_α and \vec{e}_α are defined by its components

$$\begin{aligned} (\vec{c}_\alpha)_\beta &= c_{\beta,\alpha} \\ (\vec{s}_\alpha)_\beta &= \bar{S}_{\beta,\alpha}^\dagger \\ (\vec{e}_\alpha)_\beta &= \delta_{\beta,\alpha} \end{aligned} \quad (2.21)$$

In matrix form, all these quantities are non-square matrices.

$$\begin{aligned} \vec{c}_\alpha &\stackrel{\text{Eq. 2.19}}{=} \vec{e}_\alpha + \bar{Q} \vec{s}_\alpha \stackrel{\text{Eq. 2.20}}{=} \vec{e}_\alpha + \bar{Q} \mathbf{S}^\dagger \vec{c}_\alpha \\ \Rightarrow \quad [\mathbf{1} - \bar{Q} \mathbf{S}^\dagger] \vec{c}_\alpha &= \vec{e}_\alpha \\ \Rightarrow \quad \vec{c}_\alpha &= [\mathbf{1} - \bar{Q} \mathbf{S}^\dagger]^{-1} \vec{e}_\alpha \\ \vec{s}_\alpha &\stackrel{\text{Eq. 2.20}}{=} \mathbf{S}^\dagger \vec{c}_\alpha = \mathbf{S}^\dagger [\mathbf{1} - \bar{Q} \mathbf{S}^\dagger]^{-1} \vec{e}_\alpha \end{aligned} \quad (2.22)$$

Interestingly, the vector on the right-hand side \vec{e}_α can not be simply ignored as the matrix form seems to suggest. This is specific to the calculation on the cluster. Because of this, we cannot identify the contribution of these vectors with a unit matrix.

⁸Inclusion of orbitals with vanishing screening charge does not affect the expansion coefficients $c_{\alpha,\beta}$. However, it also includes the structure constant matrix elements for the higher angular momenta with vanishing screening charge. The contribution of the higher angular momenta can be calculated after the screening has been done.

CALCULATION OF SCREENED STRUCTURE CONSTANTS

Thus, we first evaluate the bare structure constants \mathbf{S}^\dagger on the cluster, and from that $[\mathbf{1} - \bar{\mathbf{Q}}\mathbf{S}^\dagger]$. Then we solve the equation

$$[\mathbf{1} - \bar{\mathbf{Q}}\mathbf{S}^\dagger]\vec{c}_\alpha = \vec{e}_\alpha \quad (2.23)$$

$$\vec{s}_\alpha = \mathbf{S}^\dagger \vec{c}_\alpha \quad (2.24)$$

for \vec{c}_α first from Eq. 2.23 using a standard routine for linear equation systems. Using the result for \vec{c}_α , we obtain the screened structure constants \vec{s}_α from Eq. 2.24 by multiplication with the bare structure constants.

Finally we obtain the screened structure constants as

$$\bar{s}_{\gamma,\alpha}^\dagger = \left(\vec{s}_\alpha \right)_\gamma \quad (2.25)$$

Note that the screened structure constants calculated on finite clusters are no more exactly hermitean.

2.4 Derivatives

In order to evaluate forces, we need to investigate the derivatives of the structure constants with respect to changes of the atomic structure.

$$\begin{aligned} -\bar{\mathbf{Q}}d\mathbf{S}^\dagger \vec{c}_\alpha + [\mathbf{1} - \bar{\mathbf{Q}}\mathbf{S}^\dagger]d\vec{c}_\alpha &\stackrel{\text{Eq. 2.23}}{=} 0 \\ [\mathbf{1} - \bar{\mathbf{Q}}\mathbf{S}^\dagger]d\vec{c}_\alpha &= \bar{\mathbf{Q}}d\mathbf{S}^\dagger \vec{c}_\alpha \\ \Rightarrow d\vec{s}_\alpha &\stackrel{\text{Eq. 2.24}}{=} d\mathbf{S}^\dagger \vec{c}_\alpha + \mathbf{S}^\dagger d\vec{c}_\alpha = \bar{\mathbf{Q}}^{-1}d\vec{c}_\alpha \\ &= \bar{\mathbf{Q}}^{-1}[\mathbf{1} - \bar{\mathbf{Q}}\mathbf{S}^\dagger]^{-1}\bar{\mathbf{Q}}d\mathbf{S}^\dagger \vec{c}_\alpha \\ &= [\mathbf{1} - \mathbf{S}^\dagger\bar{\mathbf{Q}}]^{-1}d\mathbf{S}^\dagger \vec{c}_\alpha \end{aligned} \quad (2.26)$$

If this is correct, it implies that the screened structure constants depend only in second order on a displacement of the atoms.

2.4.1 Lower and higher angular momenta

The angular momentum expansion of a Hankel function $|K_\alpha^\infty\rangle$ about a neighboring site has infinitely many angular momentum contributions $|J_\alpha^\Omega\rangle$. The Hankel functions $|K_\alpha^\Omega\rangle$, on the other hand, are limited to a subset. We call the angular momenta, for which the Hankel functions $|K_\alpha^\Omega\rangle$ are taken into account, lower angular momenta, and those for which only the Bessel function $|J_\alpha^\Omega\rangle$ are considered, the higher angular momenta. The distinction reflects directly to the values of $\bar{\mathbf{Q}}$, which are nonzero for the lower angular momenta and which vanish for the higher angular momenta.

To avoid the confusion between the choice of screening charges and the selection of tight-binding orbitals, we assume that there is an orbital considered for each non-zero screening charge. A truncation of the basisset is postponed. With this choice, the left index of the expansion coefficients $c_{\beta,\alpha}$ in Eq. 2.11 runs over all sites in the cluster and angular momenta for which there is a non-zero screening charge \bar{Q}_β . The right index includes all orbital indices on the central site of the cluster.

If there are two orbitals with the same leading angular momentum, they will have, per definition, the same envelope function and the same tails.⁹ Thus, we consider only the angular momenta during the construction and expand the structure-constant array after the screening has been completed.

A	indices of screened orbitals
B	indices of non-zero screening charges
C	indices of partial waves
D	indices of angular momenta considered one-center expansion of the tailed representation

Screened orbitals (A) can only be formed in channels with partial waves (C), because they require augmentation of the envelope function.

Non-zero screening charges (B) are only possible where partial waves (C) exist, because the irregular contribution must be augmented.

The number of screened orbitals (A) may be a subset of (B) the channels with non0zero screening charges.

2.5 Augmentation and Potential parameters

2.5.1 Local orbitals

The local orbitals have the form

$$\begin{aligned}
 |\chi_\alpha\rangle = & |\phi_\alpha^K\rangle - \sum_{R,L} |\phi_{R,L}^J\rangle \bar{S}_{R,L,R_\alpha,L_\alpha}^\dagger \\
 & + \sum_{R',L'} |K_{R',L'}^I\rangle \left[\delta_{R',L',R_\alpha,L_\alpha} - \bar{Q}_{R',L'} \bar{S}_{R',L',R_\alpha,L_\alpha}^\dagger \right]
 \end{aligned} \tag{2.27}$$

⁹The difference between two such orbitals is thus confined to the augmentation region of the central site, and can probably be identified with a local orbital as they are used in LAPW.

where the new partial waves $|\phi_\alpha^K\rangle$ and $|\phi_\alpha^{\bar{J}}\rangle$ are superpositions of the valence and scattering partial waves that match differentiably to the head and tail functions $|K_\alpha\rangle$ and $|\bar{J}_\alpha\rangle$.

$$\begin{aligned}
 |\phi_\alpha^K\rangle &= |\phi_\alpha\rangle \overbrace{\frac{W_\alpha[K, \dot{\phi}]}{W_\alpha[\phi, \dot{\phi}]} - |\dot{\phi}_\alpha\rangle \frac{W_\alpha[K, \phi]}{W_\alpha[\phi, \dot{\phi}]}}^{\rightarrow |K_\alpha^\Omega\rangle} \\
 &\quad \underbrace{\hspace{1.5cm}}_{Ktphi} \quad \underbrace{\hspace{1.5cm}}_{-Ktophidot} \\
 |\phi_{R,L}^{\bar{J}}\rangle &= |\dot{\phi}_\beta\rangle \overbrace{\left(-\frac{W_\beta[\bar{J}, \phi]}{W_\beta[\phi, \dot{\phi}]} \right)}^{\rightarrow |\bar{J}_\beta^\Omega\rangle} \\
 &\quad \underbrace{\hspace{1.5cm}}_{JBARTophidot}
 \end{aligned} \tag{2.28}$$

Note, that in the factor $JBARTOPHIDOT$ does not depend on the choice of $|\phi\rangle$.

With $W_\alpha[f, g]$ we denote the **Wronskian**

$$W_\alpha[f, g] \stackrel{\text{def}}{=} f_\alpha(\partial_r g_\alpha) - (\partial_r f_\alpha)g_\alpha = \det \begin{bmatrix} f & g \\ \partial_r f & \partial_r g \end{bmatrix} \tag{2.29}$$

which is used to match two functions differentiably to a third via

$$y(x) \rightarrow f(x) \frac{W[y, g]}{W[f, g]} + g(x) \frac{W[y, f]}{W[g, f]} \tag{2.30}$$

The matrix elements $\langle \tilde{p}_\gamma | \tilde{\chi}_\alpha \rangle$, which will be needed later, have the form

$$\begin{aligned}
 \langle \tilde{p}_\gamma | \tilde{\chi}_\alpha \rangle &= \langle \tilde{p}_\gamma | \tilde{\phi}_\alpha^K \rangle - \sum_{R', L'} \langle \tilde{p}_\gamma | \tilde{\phi}_{R', L'}^{\bar{J}} \rangle \bar{S}_{R, L, R_\alpha, L_\alpha}^\dagger \\
 &= \langle \tilde{p}_\gamma | \tilde{\phi}_\alpha^K \rangle - \langle \tilde{p}_\gamma | \tilde{\phi}_{R_\gamma, L_\gamma}^{\bar{J}} \rangle \bar{S}_{R_\gamma, L_\gamma, R_\alpha, L_\alpha}^\dagger
 \end{aligned} \tag{2.31}$$

2.6 Coefficients of the tight-binding orbital

2.6.1 Introduction

In this section, I describe how to determine the wave functions in terms of local orbitals, if the projections $\langle \tilde{p}_\gamma | \tilde{\psi} \rangle$ onto the pseudo wave functions are known.

The basic idea is to find a representation of the wave function in terms of local orbitals $|\chi_\alpha\rangle$

$$|\psi'_n\rangle = \sum_\alpha |\chi_\alpha\rangle q_\alpha, \tag{2.32}$$

such that the deviation from the true wave function $|\psi_n\rangle$ is as small as possible.

Ideally, this would amount to minimizing the mean-square deviation of the orbital expansion from the wave function.

$$Q'[\vec{q}] := \left(\langle \psi_n | - \sum_\alpha q_\alpha^* \langle \chi_\alpha | \right) \left(| \psi_n \rangle - \sum_\beta | \chi_\beta \rangle q_\beta \right)$$

Because evaluating the mean square deviation as integral over all space is time consuming, we limit the integral to the augmentation spheres.

$$\begin{aligned} Q[\vec{q}] &:= \left(\langle \tilde{\psi}_n | - \sum_{\alpha} q_{\alpha}^* \langle \tilde{\chi}_{\alpha} | \right) \left[\sum_{\delta, \gamma} |\tilde{p}_{\delta}\rangle \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \langle \tilde{p}_{\gamma} | \right] \left(|\tilde{\psi}_n\rangle - \sum_{\beta} |\tilde{\chi}_{\beta}\rangle q_{\beta} \right) \\ &= \sum_{\gamma} \left[\sum_{\delta} \left(\langle \tilde{\psi}_n | \tilde{p}_{\delta} \rangle - \sum_{\alpha} q_{\alpha}^* \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta} \rangle \right) \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \right] \left(\langle \tilde{p}_{\gamma} | \tilde{\psi}_n \rangle - \sum_{\beta} \langle \tilde{p}_{\gamma} | \tilde{\chi}_{\beta} \rangle q_{\beta} \right) \end{aligned} \quad (2.33)$$

where $\theta_{\Omega_{R_{\delta}}}$ is a step function that vanishes outside the augmentation sphere at R_{δ} .

Minimization yields

$$\begin{aligned} \frac{\partial Q}{\partial q_{\alpha}^*} &= - \sum_{\gamma} \left[\sum_{\delta} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta} \rangle \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \right] \left(\langle \tilde{p}_{\gamma} | \tilde{\psi}_n \rangle - \sum_{\beta} \langle \tilde{p}_{\gamma} | \tilde{\chi}_{\beta} \rangle q_{\beta} \right) \stackrel{!}{=} 0 \\ \Rightarrow \quad \sum_{\gamma} \left[\sum_{\delta} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta} \rangle \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \right] \langle \tilde{p}_{\gamma} | \tilde{\psi}_n \rangle &= \sum_{\gamma, \beta} \left[\sum_{\delta} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta} \rangle \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \right] \langle \tilde{p}_{\gamma} | \tilde{\chi}_{\beta} \rangle q_{\beta} \\ \Rightarrow \quad q_{\beta} &= \sum_{\gamma} \left[\sum_{\gamma', \delta'} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta'} \rangle \langle \phi_{\delta'} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma'} \rangle \langle \tilde{p}_{\gamma'} | \tilde{\chi}_{\beta} \rangle \right]^{-1} \left[\sum_{\gamma, \delta} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta} \rangle \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \right] \langle \tilde{p}_{\gamma} | \tilde{\psi}_n \rangle \end{aligned} \quad (2.34)$$

This allows one to write the wave function in the form

$$|\psi_n\rangle \approx \sum_{\alpha} |\chi_{\alpha}\rangle \langle \tilde{\pi}_{\alpha} | \tilde{\psi}_n \rangle \quad (2.35)$$

with

$$\langle \tilde{\pi}_{\alpha} | = \sum_{\gamma} \left[\sum_{\gamma', \delta'} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta'} \rangle \langle \phi_{\delta'} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma'} \rangle \langle \tilde{p}_{\gamma'} | \tilde{\chi}_{\beta} \rangle \right]^{-1} \left[\sum_{\delta} \langle \tilde{\chi}_{\alpha} | \tilde{p}_{\delta} \rangle \langle \phi_{\delta} | \theta_{\Omega_{R_{\delta}}} | \phi_{\gamma} \rangle \right] \langle \tilde{p}_{\gamma} | \quad (2.36)$$

This expression works also if the number of local orbitals $|\chi_{\alpha}\rangle$ is smaller than the number of projector functions $\langle p_{\gamma} |$. Because of the inversion, The multicenter expansion for the projector function is long-ranged so that this expression needs to be evaluated in a Bloch representation.

2.6.2 Transformation between local-orbital and partial-wave projections

In the previous section, I derived in Eq. 2.36 a relation between orbital and partial wave projector functions.

$$\langle \tilde{\pi}_{\alpha} | \tilde{\psi}_n \rangle = \sum_{\beta} M_{\alpha, \beta} \langle \tilde{p}_{\alpha} | \tilde{\psi}_n \rangle \quad (2.37)$$

This operation is performed in `lmtotproj` with ID='FWRD'

The derivatives are correspondingly derived as

$$\begin{aligned}
 dE &= \sum_{\alpha,\beta} \underbrace{\frac{dE}{d\rho_{\alpha,\beta}}}_{=:h_{\beta,\alpha}} d\rho_{\alpha,\beta} \\
 &= \sum_{\alpha,\beta} h_{\beta,\alpha} \left[\sum_n \langle \pi_\alpha | d\psi_n \rangle f_n \langle \psi_n | \pi_\beta \rangle + \sum_n \langle \pi_\alpha | \psi_n \rangle df_n \langle \psi_n | \pi_\beta \rangle + \sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle d\psi_n | \pi_\beta \rangle \right] \\
 &= \sum_n \sum_\alpha f_n \underbrace{\sum_\beta \langle \psi_n | \pi_\beta \rangle h_{\beta,\alpha} \langle \pi_\alpha | d\psi_n \rangle}_{HTBC_{n,\alpha}^\dagger} + \sum_n \sum_\beta \langle d\psi_n | \pi_\beta \rangle \underbrace{\sum_\alpha h_{\beta,\alpha} \langle \pi_\alpha | \psi_n \rangle f_n}_{HTBC_{\beta,n}} \\
 &\quad + \sum_n \sum_\beta \langle \psi_n | \pi_\beta \rangle \underbrace{\sum_\alpha h_{\beta,\alpha} \langle \pi_\alpha | \psi_n \rangle df_n}_{HTBC_{\beta,n}} \\
 &= \sum_n \sum_\gamma f_n \underbrace{\sum_\alpha \sum_\beta \langle \psi_n | \pi_\beta \rangle h_{\beta,\alpha} M_{\alpha,\gamma} \langle \tilde{\rho}_\gamma | d\tilde{\psi}_n \rangle}_{\underbrace{HTBC_{n,\alpha}^\dagger}_{HPROJ_{n,\gamma}^\dagger}} + \sum_n \sum_\gamma \langle d\tilde{\psi}_n | \tilde{\rho}_\gamma \rangle \underbrace{\sum_\beta M_{\gamma,\beta}^\dagger \sum_\alpha h_{\beta,\alpha} \langle \pi_\alpha | \psi_n \rangle f_n}_{\underbrace{HTBC_{\beta,n}}_{HPROJ_{\gamma,n}}} \\
 &\quad + \sum_n df_n \sum_\gamma \langle \tilde{\psi}_n | \tilde{\rho}_\gamma \rangle \underbrace{\sum_\beta M_{\gamma,\beta}^\dagger \sum_\alpha h_{\beta,\alpha} \langle \pi_\alpha | \psi_n \rangle}_{\underbrace{HTBC_{\beta,n}}_{HPROJ_{\gamma,n}}} \tag{2.38}
 \end{aligned}$$

Thus, we first define the Hamiltonian \mathbf{h} (HAMIL)

$$\begin{aligned}
 \underbrace{h_{\alpha,\beta}}_{\text{HAMIL}} &= \frac{dE}{d\rho_{\beta,\alpha}} \\
 HTBC_{\beta,n} &= \sum_\alpha \underbrace{h_{\beta,\alpha}}_{\text{HAMIL}} \underbrace{\langle \tilde{\pi}_\alpha | \tilde{\psi}_n \rangle}_{TBC_{\alpha,n}} \\
 HPROJ_{\gamma,n} &= \sum_\beta M_{\gamma,\beta}^\dagger \cdot HTBC_{\beta,n} \tag{2.39}
 \end{aligned}$$

This operation is performed in `lmtotprojto` with `ID='BACK'`.

Then the forces on the wave functions

$$\sum_\gamma \frac{1}{f_n} \frac{\delta \Delta E}{\delta \langle \psi_n |} = |\tilde{\rho}_\gamma \rangle HPROJ_{\gamma,n} \tag{2.40}$$

The forces acting on the occupations are, as usual, determined as

$$\frac{\partial E}{\partial f_n} = \langle \psi_n | \left[\frac{1}{f_n} \frac{\delta E}{\delta \langle \psi_n |} \right] \tag{2.41}$$

2.7 Tailed representation of the natural tight-binding orbitals

The tailed representation replaces the multi-center expansion of the natural tight-binding orbitals by a one-center expansion. The underlying idea the short-ranged tails of the screened orbitals can be approximated by exponential tails, that are matched to the one-center expansion of the orbitals at the central site. Thus both the head and the tail functions centered on the central site are extended by a superposition of exponential tails.

2.8 How to choose the parameters

```
!CONTROL!DFT!NTBO
  MODUS='HYBRID' OFFSITE=F K2=-0.25 SCALERCUT=2. !END
!END!END!END
!STRUCTURE!SPECIES!NTBO
  NOFL=1 1 1 1 CV=T LHFWEIGHT=0.15
  TAILLAMBDA=4.0 2.0 RAUG/RCOV=0.9 RTAIL/RCOV=1.2
!END!END!END
```

2.8.1 Augmentation radius must be large for semi-core states

Observation: We had the problem that the total charge for core states of Ca has been much larger than one.

Explanation: This was apparently due to a augmentation radius that was chosen too small. The tail, represented by Hankel and Bessel functions decayed much slower than the real core state, so that the norm of the corresponding state was overestimated dramatically.

The augmentation radius specifies the matching radius of the Bessel and Hankel functions to the nodeless partial waves.¹⁰ The kinetic energy of the Hankel and Bessel function is set by the parameter `k2`. Ideally it would approximate the kinetic energy of the partial waves at the augmentation radius.

Remedy: the augmentation radius must be chosen sufficiently large so that semi-core states are well represented by their partial wave alone, while the tail represented by Hankel and Bessel functions is negligible.¹¹

¹⁰It is not related to the matching radius of all-electron, nodeless and pseudo partial waves, which defines the augmentation. The parameter here defines the shape of the natural tight-binding orbital.

¹¹Special thanks to Robert Schade.

Chapter 3

Contributions to the Hamiltonian

3.1 Core-valence exchange

The exchange term between core and valence electrons acts like a fixed, nonlocal potential acting on the electrons, of the form¹

$$\hat{v}_{x,cv} = \sum_{\alpha,\beta} |\tilde{p}_\alpha\rangle M_{\alpha,\beta} \langle \tilde{p}_\beta| \quad (3.1)$$

The core-valence exchange is furthermore diagonal in the site indices.

$$\begin{aligned} \langle \chi_\alpha | \hat{v}_{x,cv} | \chi_\beta \rangle &= \sum_{\gamma,\delta} \langle \chi_\alpha | p_\gamma \rangle M_{\gamma,\delta} \langle p_\delta | \chi_\beta \rangle \\ &= \sum_{\gamma,\delta} \langle \tilde{\phi}_\alpha^K | \tilde{p}_\gamma \rangle M_{\gamma,\delta} \langle \tilde{p}_\delta | \tilde{\phi}_\beta^K \rangle \\ &\quad - \sum_{\gamma,\delta,\beta'} \langle \tilde{\phi}_\alpha^K | \tilde{p}_\gamma \rangle M_{\gamma,\delta} \langle \tilde{p}_\delta | \tilde{\phi}_{\beta'}^J \rangle \bar{S}_{\beta',\beta}^\dagger \\ &\quad - \sum_{\gamma,\delta,\alpha',\alpha} \bar{S}_{\alpha,\alpha'} \langle \tilde{\phi}_{\alpha'}^J | \tilde{p}_\gamma \rangle M_{\gamma,\delta} \langle \tilde{p}_\delta | \tilde{\phi}_\beta^K \rangle \\ &\quad + \sum_{\gamma,\delta,\alpha',\alpha} \bar{S}_{\alpha,\alpha'} \langle \tilde{\phi}_{\alpha'}^J | \tilde{p}_\gamma \rangle M_{\gamma,\delta} \langle \tilde{p}_\delta | \tilde{\phi}_{\beta'}^J \rangle \bar{S}_{\beta',\beta}^\dagger \end{aligned} \quad (3.2)$$

Here we used the augmented Hankel and screened Bessel functions, respectively their pseudo versions.

As usual we build the expanded density matrix

$$\begin{pmatrix} \rho & -\rho \bar{S}^\dagger \\ -\bar{S} \rho & \bar{S} \rho \bar{S}^\dagger \end{pmatrix} \quad (3.3)$$

The matrix

$$\begin{pmatrix} \langle \tilde{\phi}^K | \tilde{p} \rangle \mathbf{M} \langle \tilde{p} | \tilde{\phi}^K \rangle & \langle \tilde{\phi}^K | \tilde{p} \rangle \mathbf{M} \langle \tilde{p} | \tilde{\phi}^J \rangle \\ \langle \tilde{\phi}^J | \tilde{p} \rangle \mathbf{M} \langle \tilde{p} | \tilde{\phi}^K \rangle & \langle \tilde{\phi}^J | \tilde{p} \rangle \mathbf{M} \langle \tilde{p} | \tilde{\phi}^J \rangle \end{pmatrix} \quad (3.4)$$

¹Note that this matrix \mathbf{M} differs from the one with the same symbol in the previous section.

is calculated first using `potpar1(isp)%prok` and `potpar1(isp)%projbar`.²

3.2 U-tensor

3.3 Double-counting correction

3.3.1 HSE-like

In the HSE functional[?], the double counting term exploits that the exchange-hole of the PBE functional is known.[?]. The procedure is described in the appendix of the HSE03 paper.[?]. A subroutine for the range separated exchange derived from the exchange hole is given in the Thesis of J. Heyd. ([heyd04_thesis](#)) An improved exchange-correlation hole has been determined by Bahmann and Ernzerhof[?].

See Kevin E. Riley et. al, Critical Assesment of the Performance of Density Functional Methods for Several Atomic and Molecular Properties, J. Chem. Theory Comput. 2007; 3(2): 407-433

The following does not belong here. It is obtained from Joost VandeVondele "Hybrid functionals in CP2K, A tutorial, Feb.10, 2011.https://www.cecam.org/upload/talk/presentation_5766.pdf Least square fit to the correct wave function under the constraint of orthonormality. They use the term Auxiliary density matrix method (ADMM): Guidon, Hutter and VandeVondele, J. Chem. Theory Comput., 6, 2348 (2010)

$$I = \min \left[\sum_n f_n \left(\langle \psi_n | - \sum_{\alpha} C_{n,\alpha}^{\dagger} \langle \chi_{\alpha} | \right) \left(\sum_{\gamma, \gamma'} |p_{\gamma}\rangle W_{\gamma, \gamma'} \langle p_{\gamma'} | \right) \left(| \psi_n \rangle - \sum_{\beta} | \chi_{\beta} \rangle C_{\beta, n} \right) \right. \\ \left. + \sum_{m, n} \Lambda_{n, m} \left(\left(\sum_{\alpha, \beta} C_{m, \alpha}^{\dagger} \langle \chi_{\alpha} | \chi_{\beta} \rangle C_{\beta, n} \right) - \delta_{m, n} \right) \right] \quad (3.5)$$

$$\frac{dI}{dC_{n,\alpha}^{\dagger}} = \sum_{\gamma} \left(\sum_{\gamma'} \langle \chi_{\alpha} | p_{\gamma'} \rangle W_{\gamma', \gamma} \right) \left(\langle p_{\gamma} | \psi_n \rangle - \sum_{\beta} \langle p_{\gamma} | \chi_{\beta} \rangle C_{\beta, n} \right) f_n \\ + \sum_m \sum_{\beta} \langle \chi_{\alpha} | \chi_{\beta} \rangle C_{\beta, m} \Lambda_{m, n} \stackrel{!}{=} 0 \quad (3.6)$$

$$\Rightarrow \sum_{\beta} \left(\sum_{\gamma, \gamma'} \langle \chi_{\alpha} | p_{\gamma'} \rangle W_{\gamma', \gamma} \langle p_{\gamma} | \chi_{\beta} \rangle \right) C_{\beta, n} - \sum_m \sum_{\beta} \langle \chi_{\alpha} | \chi_{\beta} \rangle C_{\beta, m} \Lambda_{m, n} \frac{1}{f_n} \\ \stackrel{!}{=} \sum_{\gamma} \left(\sum_{\gamma'} \langle \chi_{\alpha} | p_{\gamma'} \rangle W_{\gamma', \gamma} \right) \langle p_{\gamma} | \psi_n \rangle \quad (3.7)$$

² In the earlier version the contribution from the $\dot{\phi}$ has been ignored!!! It has been verified by temporarily switching off the jbar contributiun to `potpar1(isp)%prok` and `potpar1(isp)%projbar`. In this old version only `potpar(isp)%ktophi` is used to extract the ϕ contribution.

3.3.2 Blöchl, Walther, Pruschke

The double-counting correction is that of Eq. 36 of the paper by Blöchl, Walther, Pruschke[?].

It is based on a partitioning of the exchange correlation energy in the form using the functions

$$g_R(\vec{r}) = \frac{n_R^{\chi,s}(\vec{r})}{\sum_{R'} n_{R'}^{\chi,s}(\vec{r})} = \left[1 + \frac{\sum_{R'; R' \neq R} n_{R'}(\vec{r})}{n_R(\vec{r})} \right]^{-1} \quad (3.8)$$

where $n_R^{\chi,s}(\vec{r})$ is the spherical part of the density from the local orbitals $|\chi\rangle$. The partitioning functions obey $\sum_R g_R(\vec{r}) = 1$, which allows to rewrite the exchange correlation energy in the form

$$E_{xc} = \sum_{R,R'} \int d^3r n(\vec{r}) \epsilon_{xc}[n(\vec{r})] g_R(\vec{r}) g_{R'}(\vec{r}) \quad (3.9)$$

The resulting onsite terms are used for our double-counting term for an on-site interaction.

$$E_{xc}^{\hat{W}_R} = \int d^3r \left(n(\vec{r}) \epsilon_{xc}[n] \right) g_R^2(\vec{r}) \quad (3.10)$$

The exchange correlation density is calculated with the partial wave expansion, and it is cut off by the squared partitioning function.

Other double-counting schemes are discussed in appendix D.

- For hybrid functionals only the scaled Hartree-Fock energy is added, so that the double counting is weighted as well and it is furthermore limited to the exchange term only. **This will be different, if also corrections for the correlation energies are added.**
- Let us simplify the expression by introducing the symbols $n_R = n_R^{\chi}(\vec{r})$ and $n_t = n_{\sigma,\sigma'}^{\chi}(\vec{r})$. With n_R^s and n_t^s we denote the spherical parts of the respective densities.
- If the frozen-core density is employed, the contribution of the core must be excluded in the double-counting term.
- In the denominator of the cutoff function $g_R(\vec{r})$, I replace the multicenter expansion of the density in terms of local orbitals, the one-center partial-wave expansion.

The expression above can be written in the following form:

$$\begin{aligned} E_{xc}^{\hat{W}_R} &= \int d^3r \left[n_t(\vec{r}) \epsilon_{xc}[n_t(\vec{r})] - n^{core}(\vec{r}) \epsilon_{xc}[n^{core}(\vec{r})] \right] \left(\frac{n_R^s(\vec{r})}{n_t^s(\vec{r})} \right)^2 \\ &= \int d^3r f_{xc}(\vec{r}) \left(\frac{n_R^s(\vec{r})}{n_t^s(\vec{r})} \right)^2 \end{aligned} \quad (3.11)$$

With $f_{xc} := n_t \epsilon_{xc}[n_t] - n^{core} \epsilon_{xc}[n^{core}]$ and $\mu_{xc} := \frac{df_{xc}}{dn_t}$, we obtain

$$\begin{aligned} dE_{xc}^{\hat{W}_R} &= \int d^3r \left(\frac{n_R^s}{n_t^s} \right)^2 \mu_{xc} dn_t + f_{xc} 2 \left(\frac{n_R^s}{n_t^s} \right)^2 \frac{dn_R^s}{n_R^s} - f_{xc} 2 \left(\frac{n_R^s}{n_t^s} \right)^2 \frac{dn_t^s}{n_t^s} \\ &= \int d^3r \left[\left(\frac{n_R^s}{n_t^s} \right)^2 \mu_{xc} - 2f_{xc}^s \left(\frac{n_R^s}{n_t^s} \right)^2 \frac{1}{n_t^s} \right] dn_t + \left[2f_{xc}^s \left(\frac{n_R^s}{n_t^s} \right)^2 \frac{1}{n_R^s} \right] dn_R^s \end{aligned} \quad (3.12)$$

which yields the two potentials

$$\begin{aligned} v_t &\stackrel{\text{def}}{=} \left(\frac{n_R^s}{n_t^s} \right)^2 \left[\mu_{xc} - \frac{2f_{xc}^s}{n_t^s} \right] \\ v_R &\stackrel{\text{def}}{=} \left(\frac{n_R^s}{n_t^s} \right)^2 \frac{2f_{xc}^s}{n_R^s} \end{aligned} \quad (3.13)$$

For the cutoff function $(n_R^s/n_t^s)^2$ we consider only the spherical contributions of the density and we ignore the spin contributions. This is accounted for in the derivations by only considering the spherical part f_{xc}^s of f_{xc} , while maintaining the non-spherical contributions to μ_{xc} .

The total density n_t contains also the core density. If the core valence Fock term is included, they are part of the correlated electrons and need to be considered in the density n_R .

The double-counting correction has the negative sign, because the DFT-expression needs to be subtracted.³ Its energy and the corresponding contributions to the auxiliary Hamiltonian are

$$\begin{aligned} E_{dc} &= - \int d^3r \left(\frac{n_R^s}{n_t^s} \right)^2 f_{xc}^s \\ d\hat{H}_{dc} &= -|\tilde{\pi}_\alpha\rangle\langle\chi_\alpha|\hat{v}_t|\chi_\beta\rangle\langle\tilde{\pi}_\beta| - |\tilde{\rho}_\alpha\rangle\langle\phi_\alpha|\hat{v}_R|\phi_\beta\rangle\langle\tilde{\rho}_\beta| \end{aligned} \quad (3.14)$$

We start from two density matrices, which are the same at the moment.

see `lmto_simplifiedc_new`.

Idea

$$E_{xc}^{\hat{W}_R} = \sum_{a,b,c,d} \int d^3r \chi_a^*(r) \chi_b(r) \chi_d^*(r) \chi_c(r) \frac{\epsilon_{xc}[n_{\sigma,\sigma'}^x(\vec{r})]}{n^x(\vec{r})} \quad (3.15)$$

³This subtraction is done outside the routine calculating the double-counting term.

Chapter 4

Description of Subroutines

4.1 Workflow

```
---initialization-----
POTPAR = potential parameters
SBAR = screened structureconstants
<ptilde|chitilde> tailed partial waves overlap (Onsite)
utensor (Onsite)
utensor (offsite)
...
----cycle-----
TBC=<pi-tilde|psi> from PROJ=<ptilde|psitide>
DENMAT density matrix in local orbitals
...
total energy and derivatives
HAMIL hamiltonian matrix in tight-binding orbitals
...
HTBC = de/dtbc * 1/f
HPROJ = de/dproj * 1/f
```

4.2 LMTO_POTPAR

All information that depends directly on the partial waves is stored in the structure POTPAR.

POTPAR Structure	
RAD	augmentation radius
Quantities connected to head functions	
NHEAD	number of head functions
LOFH(NHEAD)	angular momentum
ITAIL(NHEAD)	pointer to tail function
LNOFH(NHEAD)	pointer to partial wave $ \phi\rangle$ and projector $\langle p $
KTOPHI(NHEAD)	$ K^\Omega\rangle \rightarrow \phi\rangle_{C_{KTOPHI}} + \dot{\phi}\rangle_{C_{KTOPHIDOT}}$
KTOPHIDOT(NHEAD)	
Quantities connected to tail functions	
NTAIL(NTAIL)	number of tail functions
LOFT(NTAIL)	angular momentum
LNOFT(NTAIL)	pointer to partial wave $ \phi\rangle$ and projector $\langle p $
QBAR(NTAIL)	screening charge $ \bar{J}\rangle = J\rangle - K\rangle\bar{Q}$
JBARTOPHIDOT(NTAIL)	$ \bar{J}^\Omega\rangle \rightarrow \dot{\phi}\rangle_{C_{JBARTOPHIDOT}}$
Other stuff	
PROK(LNX,NHEAD)	$\langle \tilde{p} \phi_{C_{ktopphi}} + \dot{\phi}_{C_{ktophidot}} \rangle$
PROJBAR(LNX,NTAIL)	$\langle \tilde{p} \dot{\phi}_{C_{jbartophidot}} \rangle$
PHIOV(LNX,LNX)	$\langle \phi_\alpha \theta_\Omega \phi_\beta \rangle$
Tailed representation	
TAILED%GID	grid id for the radial grid
TAILED%LNX	
TAILED%LMNX	
TAILED%LOX(LNX)	
TAILED%AEF(NR,LNX)	
TAILED%PSF(NR,LNX)	
TAILED%NLF(NR,LNX)	
TAILED%U(LMNX,LMNX,LMNX,LMNX)	
TAILED%OVERLAP(LMNX,LMNX)	
TAILED%QLN(2,LNX,LNX)	

The variable lnx and lmnx in the substructure TAILED differ from the corresponding functions from the partial wave expansion.

4.3 LMT0\$CLUSTERSTRUCTURECONSTANTS

LMT0\$CLUSTERSTRUCTURECONSTANTS calculates the screened structure constants SBAR (\bar{S}) for a cluster of NAT atomic sites RPOS, of which the first site is called the central site of the cluster. The number of angular momenta on each site is defined by LX. The screening is defined by the vector QBAR (\bar{Q}). K2 ($\bar{k}^2 = -\kappa^2$) is the squared wave vector. (For envelope functions that fall off exponentially, this parameter is negative.)

```

SUBROUTINE LMT0$CLUSTERSTRUCTURECONSTANTS(K2,NAT,RPOS,LX,QBAR,NORB,N,SBAR)
REAL(8)      ,INTENT(IN)  :: K2
INTEGER(4)   ,INTENT(IN)  :: NAT          ! NUMBER OF ATOMS ON THE CLUSTER
REAL(8)      ,INTENT(IN)  :: RPOS(3,NAT) ! ATOMIC POSITIONS ON THE CLUSTER
INTEGER(4)   ,INTENT(IN)  :: LX(NAT)      ! X(ANGULAR MOMENTUM ON EACH CLUSTER)
INTEGER(4)   ,INTENT(IN)  :: N
REAL(8)      ,INTENT(IN)  :: QBAR(N)
INTEGER(4)   ,INTENT(IN)  :: NORB
REAL(8)      ,INTENT(INOUT):: SBAR(NORB,N)

```

First, the bare structure constants are evaluated on the cluster using LMT0\$\\$STRUCTURECONSTANTS and then the structure constants are screened using LMT0\$\\$SCREEN.

4.3.1 LMT0\$\\$STRUCTURECONSTANTS

LMT0\$\\$STRUCTURECONSTANTS calculates the bare structure constants for a pair of sites. The first site is at the origin, where the Hankel function is centered, and the second site at \vec{R} specified by R21, is the center of the expansion into solid Bessel functions.

```

subroutine lmt0$structureconstants(r21,K2,L1x,L2x,S)
REAL(8)      ,INTENT(IN)  :: R21(3) ! EXPANSION CENTER
INTEGER(4)   ,INTENT(IN)  :: L1X
INTEGER(4)   ,INTENT(IN)  :: L2X
REAL(8)      ,INTENT(IN)  :: K2 ! 2ME/HBAR**2
REAL(8)      ,INTENT(OUT):: S((L1X+1)**2,(L2X+1)**2)

```

The bare structure constants are evaluated in LMT0\$\\$STRUCTURECONSTANTS as

$$S_{RL,R'L'} \stackrel{\text{Eq. 2.7}}{=} (-1)^{\ell'+1} 4\pi \sum_{L''} C_{L,L',L''} H_{L''}(\vec{R}' - \vec{R}) \begin{cases} (-ik)^{\ell+\ell'-\ell''} & \text{for } k^2 > 0 \\ \delta_{\ell+\ell'-\ell''} & \text{for } k^2 = 0 \\ \kappa^{\ell+\ell'-\ell''} & \text{for } k^2 = -\kappa^2 < 0 \end{cases} \quad (4.1)$$

where $H_L(k^2, \vec{R})$ is the solid Hankel function calculated in LMT0\$\\$SOLIDHANKEL. The solid Hankel function is the solution of the Helmholtz equation, Eq. 2.3.¹

More information on the solid Hankel function can be found in appendix A.

Remark: Because the Gaunt coefficients vanish for odd $\ell + \ell' - \ell''$, the structure constants are real even for $k^2 > 0$.

4.3.2 LMT0\$\\$SCREEN

I describe here what has been implemented as “version 3”.

LMT0\$\\$SCREEN takes the bare structure constants $S_{RL,R'L'}$ connecting all orbitals on a specific cluster with each other and the screening constants \bar{Q} for all orbitals on the cluster. It returns

¹The factors and signs of the inhomogeneity need to be confirmed. The equation has been taken from the methods book, chapter “Working with spherical Hankel and Bessel functions.”

the screened structure constants connecting the orbitals on the central (first) site (1st index) with all orbitals (2nd index).

The structure constants are defined so that

$$\langle K_{RL} | = - \sum_{L'} S_{RL,R'L'} \langle J_{R'L'} | \quad \text{for } R' \neq R \quad (4.2)$$

First we evaluate

$$\mathbf{A} = \mathbf{1} - \bar{\mathbf{Q}}\mathbf{S}^\dagger \quad (4.3)$$

and the vectors \vec{e}_α defined by $(\vec{e}_\alpha)_\beta = \delta_{\beta,\alpha}$. Note that the number of vectors corresponds to the number of orbitals on the central site only. Therefore, these vectors do not build up a complete unit matrix.

Then we solve the equation system

$$\mathbf{A}\vec{c}_\alpha \stackrel{\text{Eq. 2.23}}{=} \vec{e}_\alpha \quad (4.4)$$

for \vec{c}_α and

$$\vec{s}_\alpha \stackrel{\text{Eq. 2.24}}{=} \mathbf{S}^\dagger \vec{c}_\alpha \quad (4.5)$$

$(\vec{s}_\alpha)_\beta = \bar{S}_{\beta,\alpha}^\dagger$ contains the transposed screened structure constants. After transposition, \bar{S} is returned.

4.4 Waves object

The data exchange between the waves object and the lmtot object is determined by the local-orbital projections $\langle \tilde{\pi}_\alpha | \tilde{\psi}_n \rangle$ specified by the array THIS%TBC, which in turn is obtained from the partial-wave projections $\langle \tilde{\rho} | \tilde{\psi}_n \rangle$.

In waves\$etot

```
CALL WAVES$TONTBO
-> CALL LMTO$PROJTONTBO('FWRD'...)
..
..
CALL LMTO$ETOT(LMNXX,NDIMD,NAT,DENMAT)
..
..
CALL WAVES$FROMNTBO()
-> CALL LMTO$PROJTONTBO('BACK'...)
..
..
CALL WAVES$FORCE
-> CALL WAVES_FORCE_ADDHTBC
...
CALL WAVES$HPSI
```

$$\begin{aligned}
\vec{F} &= - \sum_{\alpha} \frac{dE}{d\langle \tilde{\rho}_{\alpha} | \psi_n \rangle} \langle \vec{\nabla}_R \tilde{\rho}_{\alpha} | \psi_n \rangle + \text{c.c.} \\
&= - \sum_{\alpha, \beta} \frac{dE}{d\langle \tilde{\pi}_{\beta} | \psi_n \rangle} \frac{d\langle \tilde{\pi}_{\beta} | \psi_n \rangle}{d\langle \tilde{\rho}_{\alpha} | \psi_n \rangle} \langle \vec{\nabla}_R \tilde{\rho}_{\alpha} | \psi_n \rangle + \text{c.c.} \\
&= - \sum_{\alpha, \beta} \frac{dE}{d\langle \tilde{\pi}_{\beta} | \psi_n \rangle} \frac{d\langle \tilde{\pi}_{\beta} | \psi_n \rangle}{d\langle \tilde{\rho}_{\alpha} | \psi_n \rangle} \left[- \langle \vec{\nabla}_r \tilde{\rho}_{\alpha} | \psi_n \rangle \right] + \text{c.c.}
\end{aligned}$$

4.5 Offsite matrix elements

The offsite matrix elements are kept in the data type

```

TYPE OFFSITEX_TYPE
  INTEGER(4)      :: NDIS
  INTEGER(4)      :: NF
  REAL(8) , POINTER :: OVERLAP(:, :) ! OVERLAP MATRIX ELEMENTS
  REAL(8) , POINTER :: X22(:, :)    !
  REAL(8) , POINTER :: X31(:, :)
  REAL(8) , POINTER :: BONDU(:, :)
  REAL(8) , POINTER :: DIS(:)
  REAL(8) , POINTER :: LAMBDA(:)
END TYPE OFFSITEX_TYPE

```

The matrix elements are initialized in LMT0_initialize

```

LMT0_TAILEDGAUSSFIT()
  GAUSSIAN_FITGAUSS(GID, NR, W, L, AUX, NE, NPOW2, E, C(:NPOW2, :, LN))
  LMT0_TAILEDGAUSSORBTOLM()
LMT0_OFFXINT
  LMT0_OFFSITEOVERLAPSETUP !O(AB) ->OFFSITEX%OVERLAP
    LMT0_TWOCENTER
  LMT0_OFFSITEX22SETUP      !U(AABB) ->OFFSITEX%X22
    LMT0_TWOCENTER
  LMT0_OFFSITEX31SETUP      !U(AAAB) ->OFFSITEX%X31
    LMT0_TWOCENTER
  LMT0_TAILEDGAUSSOFFSITEU !U(ABAB) ->OFFSITEX%BONDU
    GAUSSIAN$ZDIRECTION_FOURCENTER(NIJKA, NEA, EA, LMNXA, ORBA &
    LMT0_OFFSITEXCONVERT()

```

The energy contibution is then calculated using offsitex as follows

```

LMT0_OFFSITEXEVAL_NEW(EX)
  LMT0_EXPANDNONLOCAL
  LMT0_EXPANDLOCAL
  SPHERICAL$ROTATEYLM(LMX, ROT, YLMROT)
  LMT0_OFFSITEX22U(ISPA, ISPB, DIS, LMNXTA, LMNXTB, U22, DU22)

```

```

      lmto_offsitexvalue
      LMTO_OFFSITEEX31U(ISPA,ISPB, DIS,LMNXTA,LMNXTB,U3A1B,DU3A1B)
      lmto_offsitexvalue
      LMTO_OFFSITEEX31U(ISPB,ISPA,-DIS,LMNXTB,LMNXTA,U3B1A,DU3B1A)
      lmto_offsitexvalue
      LMTO_OFFSITEEXBONDU(ISPA,ISPB,DIS,LMNXTA,LMNXTB,BONDU,DBONDU)
      lmto_offsitexvalue

```

4.6 Matrix elements using Gaussians

LMTO_TAILEDGAUSSOFFSITEU uses the gauss decomposition of the tailed orbitals in potpar%tailed%gaussnlf. In tailedgaussfit the following data structure is prepared.

```

POTPAR%TAILED%GAUSSNLF%NIJK
POTPAR%TAILED%GAUSSNLF%NORB
POTPAR%TAILED%GAUSSNLF%NPOW
POTPAR%TAILED%GAUSSNLF%NE
POTPAR%TAILED%GAUSSNLF%E
POTPAR%TAILED%GAUSSNLF%C

```

4.7 Matrix elements on an adaptive grid

```

lmto_twocenter

MODULE LMTO_TWOCENTER_MODULE
LMTO_TWOCENTER
  ADAPT$EVALUATE
    ADAPTINI
    ADAPT_BASICRULE
    ADAPT_INTEGRAND
    LMTO_TWOCENTER_MYFUNC

```

4.8 Routines for reporting

```

LMTO$REPORT(NFIL)
...
LMTO$REPORTOVERLAP(NFIL)
LMTO$REPORTSBAR(NFIL)
LMTO$REPORTDENMAT(NFIL)
LMTO$REPORTHAMIL(NFIL)
LMTO$REPORTPERIODICMAT(NFIL,NAME,NNS,SBAR)
...
LMTO$WRITEPHI(FILE,GID,NR,NPHI,PHI)

```

4.9 Routines for plotting orbitals

There are three routines that calculate the orbitals either in a spherical-harmonics expansion on radial grids or directly on an array of real-space points.

```

LMTO_TAILEDORBLM(IAT,IORB,NR,LMX,ORB)
LMTO_TAILED_NTBOOFR(IAT,iORB,NP,P,chi)
  LMTO_TAILEDORBLM(IAT,IORB,NR,LMX,ORB)
LMTO_NTBOOFR(IAT,iORB,NP,P,chi)

LMTO_PLOTTAILED() [OK]
  LMTO_TAILEDORBLM(IAT,IORB,NR,LMX,ORB)
LMTO_GRIDPLOT(type,iat) [ok]
  LMTO_GRIDORB_CUBEGRID(RO(: , IATO), RANGE, N1, N2, N3, ORIGIN, TVEC, P)
  LMTO_GRIDORB_STARGRID(RO(: , IATO), RANGE, NDIR, DIR, NRAD, X1D, P)
  LMTO_TAILED_NTBOOFR(TYPE, IAT, iORB, NP, P, chi)
    LMTO_TAILEDORBLM(IAT, IORB, NR, LMX, ORB)
    LMTO_NTBOOFR(IAT, iORB, NP, P, chi)
  LMTO_WRITECUBEFILE
..
LMTO_GRIDPLOT_UNTAILED(IATO)
  LMTO_GRIDORB_CUBEGRID(RO(: , IATO), RANGE, N1, N2, N3, ORIGIN, TVEC, P)
  LMTO_GRIDORB_STARGRID(RO(: , IATO), RANGE, NDIR, DIR, NRAD, X1D, P)
  LMTO_GRIDENVELOPE(RBAS, NAT, RO, IATO, LMX, NP, P, ENV, ENV1)
  LMTO_GRIDAUGMENT(RBAS, NAT, RO, IATO, LMX, NP, P, ORB1, ENV1)
  LMTO_GRIDGAUSS(RBAS, NAT, RO, IATO, LMX, NP, P, ORBG)
  LMTO_WRITECUBEFILE(NFIL, NATCLUSTER, ZCLUSTER, RCLUSTER &
LMTO_PLOTLOCORB(IATO)
  LMTO_GRIDENVELOPE(RBAS, NAT, RO, IATO, LM1X, NP, P, ENV, ENV1)
  LMTO_GRIDAUGMENT(RBAS, NAT, RO, IATO, LM1X, NP, P, ORB1, ENV1)
  LMTO_GRIDGAUSS(RBAS, NAT, RO, IATO, LM1X, NP, P, ORBG)
  LMTO_WRITECUBEFILE
LMTO_PLOTNTBO(TYPE, IATORB, LMNORB)
  LMTO_GRIDORB_CUBEGRID(CENTER, RADIUS, N1, N2, N3, ORIGIN, TVEC, P)
  LMTO_GRIDORB_STARGRID(CENTER, RADIUS, NDIR, DIR, NR, X1D, P)
  LMTO_TAILED_NTBOOFR(TYPE, IATORB, LMNORB, NP, P, ORB)
  LMTO_NTBOOFR(TYPE, IATORB, LMNORB, NP, P, ORB)
..
LMTO$PLOTWAVE(NFIL, IDIMO, IBO, IKPTO, ISPIN0, NR1, NR2, NR3)
  LMTO$PLOTWAVE_TAILED(NFIL, IDIMO, IBO, IKPTO, ISPIN0, NR1, NR2, NR3)
  WRITEWAVEPLOT(NFIL, TITLE, RBAS, NAT, RO, ZAT, Q, NAME, XK, NR1, NR2, NR3, WAVE)

```

- LMTO_TAILEDORBLM(IAT, IORB, NR, LMX, ORB) calculates a specific orbital on radial grids in a spherical harmonics representation.
- LMTO_PLOTTAILED() writes the tailed local orbitals in a spherical harmonics expansion to file, so that the componenst can be viewed by xmgrace. Each orbital is written to a file CHI5_3.DAT where 5 is the atom index and 3 is the orbital index.

- `LMT0_TAILED_NTBOOFR(IAT,iORB,NP,P,chi)` calculates a specific tailed orbital on a set of real space points.
- `LMT0_GRIDPLOT_TAILED(IAT0)` writes the orbitals for the specified site to file. It supports a 3D representation with cube files, and one-dimensional representation on a star grid.

Multicenter expansion

First the interstitial orbital is determined

$$\begin{aligned}
 |\bar{K}_\alpha^I\rangle &= |\bar{K}_\alpha^\infty\rangle - \left[|K_\alpha^\Omega\rangle - \sum_\beta |J_\beta^\Omega\rangle \bar{S}_{\beta,\alpha}^\dagger \right] \\
 &= \sum_\beta |K_\beta^\infty\rangle \left(\delta_{\alpha,\beta} + \bar{Q}_\beta \bar{S}_{\beta,\alpha}^\dagger \right) - \left[|K_\alpha^\Omega\rangle - \sum_\beta |J_\beta^\Omega\rangle \bar{S}_{\beta,\alpha}^\dagger \right] \\
 &= \sum_\beta |K_\beta^\infty\rangle \left(\delta_{\alpha,\beta} + \bar{Q}_\beta \bar{S}_{\beta,\alpha}^\dagger \right) - \left[|K_\alpha^\Omega\rangle - \sum_\beta \left(|J_\beta^\Omega\rangle - |K_\beta^\Omega\rangle \bar{Q}_\beta \right) \bar{S}_{\beta,\alpha}^\dagger \right] \\
 &= \sum_\beta \left(|K_\beta^\infty\rangle - |K_\beta^\Omega\rangle \right) \left(\delta_{\alpha,\beta} + \bar{Q}_\beta \bar{S}_{\beta,\alpha}^\dagger \right) + \sum_\beta |J_\beta^\Omega\rangle \bar{S}_{\beta,\alpha}^\dagger
 \end{aligned} \tag{4.6}$$

This implies that within the sphere centered at R_β only the expansion into bare bessel functions survive, while outside only the bare Hankel function is considered.

Chapter 5

Benchmarks

5.1 Silicon

The files are in [~/Tree/Projects/SetupTests/Si](#).

I performed calculation for a silicon crystal. We used a mixing of the local exchange of $\alpha_X = 0.1$. The local basisset included one s-type and one p-type tight-binding function.

The following dependencies have been explored:

- kinetic energy of the Hankel function: $K^2 = 0, \dots, -0.5$




Fig. 5.1: Energy in Hartree and band gap in eV of the silicon crystal as function of the kinetic energy (times -0.01) of the envelope function for the PBE functional (black; none), local Hartree-Fock (red; onsite or local) NDDO-type exchange, i.e density of one site with the density on a bond partner (blue, NDDO), terms with three orbitals on one site and one on the bond partner (green NDDO+31) and the exchange of the bond density with itself (orange, all bond). The full line is the result for one s-type and one p-type orbital. For the dashed lines also d-type orbitals are included. The dotted lines are calculated with double orbitals for s,p and single orbitals for d.

- Augmentation radius. The augmentation radius is the radius at which the partial waves

are matched to the envelope function. It is also the radius for the sphere used to derive the projector function onto the local orbitals. The procedure apparently breaks down






Fig. 5.2: Energy in Hartree and band gap in eV of the silicon crystal as function of the augmentation radius for the PBE functional (black; none), local Hartree-Fock (red; onsite or local) NDDO-type exchange, i.e. density of one site with the density on a bond partner (blue, NDDO), terms with three orbitals on one site and one on the bond partner (green NDDO+31) and the exchange of the bond density with itself (orange, all bond). The full line is the result for one s-type and one p-type orbital. For the dashed lines also d-type orbitals are included. The dotted lines are calculated with double orbitals for s,p and single orbitals for d.

completely, if the augmentation radius is too large. The dependency becomes stronger, if more non-local terms are included. This may be due to the fact that the double-counting term is only included for the local terms.

- tail matching radius





5.1.1 Summary

- The augmentation radius has the largest effects on the results both for the gap and for the total energy. Beyond a certain radius ($> 1.2 r_{cov}$) the calculation becomes even unstable. For an “overcomplete TB-basisset $2s + 2p + 1d$ ” the calculation also fails for ($< 1.1 r_{cov}$).
- It becomes evident that the energy is rather insensitive to the parameters describing the local orbitals for local exchange and the NDDO terms. Additional terms such as “31” and “bondx”, which include the bond overlap density $\chi_R(r)\chi_{R'}(r)$ lead to a very strong dependency on the choice of orbitals. Note that $0.05 \text{ H} \approx 1.4 \text{ eV}$!

This may be due to the poor description of the bond density by the exponential tails of the tailed orbitals.

It may also be due to the fact these terms are not compensated by the double-counting term.

- It should be noted that the double-counting term is included only for local exchange.
- Choose value of $-0.2 < k^2 < -0.5 \text{ H}$
- Choose value of $r_{tail} = 1.2 r_{cov}$

Chapter 6

Implementation of simpler orbitals

This is a description of a more recent development using simpler orbitals. The basic idea is to simplify the orbitals so that they do not change their shape but that are directly tied to the central atom. This can be accomplished by using envelope functions that fall off so rapidly, that they are almost zero at the nuclear site of the neighboring atoms.

6.1 Local orbitals

Local orbitals are augmented Hankelfunctions

$$\begin{aligned}
 |\chi_\gamma\rangle &= \sum_\alpha \left\{ \underbrace{|\phi_\alpha^K\rangle + |K_\alpha^{\notin\Omega_\alpha}\rangle}_{\text{R-independent}} - \underbrace{\sum_\beta \left(|\phi_\beta^J\rangle - |J_\beta\rangle \right) S_{R_\beta, L_\beta, R_\alpha, L_\alpha}^\dagger}_{\text{R-dependent and small}} \right\} c_{\alpha, \gamma} \\
 &= \sum_\alpha \left\{ |\phi_\alpha^K\rangle - \sum_\beta |\phi_\beta^J\rangle S_{R_\beta, L_\beta, R_\alpha, L_\alpha}^\dagger + |K_\alpha^I\rangle \right\} c_{\alpha, \gamma}
 \end{aligned} \tag{6.1}$$

Here, $|\phi_\alpha^K\rangle$ is a superposition of partial waves, which match at the augmentation radius to $|K_\alpha\rangle$, and which are set to zero outside the augmentation region. $|K_\alpha^{\notin\Omega_\alpha}\rangle$ is the Hankelfunction at site R_α and angular momentum L_α , which is nonzero only outside its central augmentation region.

A suitable definition for \mathbf{c} is a matrix that produces orthonormal states for an isolated atom. Thus, it is an onsite matrix.

We introduce head and tail functions defined as

$$\begin{aligned}
 |\phi_\alpha^H\rangle &= \sum_{\beta; R_\beta=R_\alpha} |\phi_\beta^K\rangle c_{\beta, \alpha} \\
 |\phi_\alpha^T\rangle &= |\phi_\alpha^J\rangle
 \end{aligned} \tag{6.2}$$

so that

$$|\chi_\gamma\rangle = |\phi_\gamma^H\rangle - \sum_{\alpha, \beta} |\phi_\beta^J\rangle S_{R_\beta, L_\beta, R_\alpha, L_\alpha}^\dagger c_{\alpha, \gamma} + \sum_\alpha |K_\alpha^I\rangle c_{\alpha, \gamma} \tag{6.3}$$

6.2 Projector functions

6.2.1 Fitting

The projections are determined such that the square deviation between the original wave functions and the local orbital representation is minimized. The mean square deviation is approximated by the partial wave expansions in the augmentation regions.

$$F(\vec{q}) \stackrel{\text{def}}{=} \left(\langle \psi | - q \langle \chi | \right) \left(| p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \right) \left(| \psi \rangle - | \chi \rangle q \right) \stackrel{!}{=} \min \quad (6.4)$$

$$\begin{aligned} \frac{dF}{dq^*} &= -\langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \left(\langle p | \psi \rangle - \langle p | \chi \rangle q \right) \stackrel{!}{=} 0 \\ \langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \psi \rangle &= \langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \chi \rangle q \\ q &= \left(\langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \chi \rangle \right)^{-1} \langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \psi \rangle \\ \langle \pi | &= \left(\langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \chi \rangle \right)^{-1} \langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \end{aligned} \quad (6.5)$$

Projection

Here I show that the fitting procedure described above is identical to a projection, with the integration limited to the partial wave expansion in the augmentation regions.

$$\begin{aligned} \hat{P}^\chi | \psi \rangle &= | \chi \rangle \left(\langle \chi | \chi \rangle \right)^{-1} \langle \chi | \psi \rangle \\ &\approx | \chi \rangle \left(\langle \chi | \left(\sum_R \theta_{\Omega_R} \right) | \chi \rangle \right)^{-1} \langle \chi | \left(\sum_R \theta_{\Omega_R} \right) | \psi \rangle \\ &\approx | \chi \rangle \left(\langle \chi | \left(| p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \right) | \chi \rangle \right)^{-1} \langle \chi | \left(| p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \right) | \psi \rangle \end{aligned} \quad (6.6)$$

Thus, I obtain the identical form for the projector function as found above in Eq. 6.5 for the fitting.

$$\langle \pi | = \left(\langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \chi \rangle \right)^{-1} \langle \chi | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \quad (6.7)$$

6.3 Expressions

$$\begin{aligned} \langle p_\alpha | \chi_\beta \rangle &= \langle p_\alpha | \phi_\beta^H \rangle - \sum_{\gamma, \delta} \langle p_\alpha | \phi_\gamma^T \rangle \underbrace{S_{R_\gamma, L_\gamma, R_\delta, L_\delta}^\dagger}_{S_{R_\delta, L_\delta, R_\gamma, L_\gamma}^*} c_{\delta, \beta} \\ \langle \chi_\alpha | p_\beta \rangle &= \langle \phi_\alpha^H | p_\beta \rangle - \sum_{\gamma, \delta} c_{\alpha, \delta}^\dagger S_{R_\delta, L_\delta, R_\gamma, L_\gamma} \langle \phi_\gamma^T | p_\beta \rangle \end{aligned} \quad (6.8)$$

$$\begin{aligned} \langle \pi | &= \left\{ \langle \phi^H | \theta_\Omega | \phi^H \rangle - \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | \theta_\Omega | \phi^H \rangle - \langle \phi^H | \theta_\Omega | \phi^J \rangle \mathbf{S}^\dagger \mathbf{c} + \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | \theta_\Omega | \phi^J \rangle \mathbf{S}^\dagger \mathbf{c} \right\}^{-1} \\ &\times \left(\langle \phi^H | p \rangle - \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | p \rangle \right) \langle \phi | \theta_\Omega | \phi \rangle \langle p | \end{aligned} \quad (6.9)$$

where we used the short-hand notation

$$\begin{aligned}\langle \phi^H | \theta_\Omega | \phi^H \rangle &= \langle \phi^H | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \phi^H \rangle \\ \langle \phi^H | \theta_\Omega | \phi^J \rangle &= \langle \phi^H | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \phi^J \rangle \\ \langle \phi^J | \theta_\Omega | \phi^J \rangle &= \langle \phi^J | p \rangle \langle \phi | \theta_\Omega | \phi \rangle \langle p | \phi^J \rangle\end{aligned}\tag{6.10}$$

6.3.1 Linear approximation in structure constants

To linear order in the structure constants I obtain the projector as

$$\begin{aligned}\langle \pi | &= \left\{ \langle \phi^H | \theta_\Omega | \phi^H \rangle - \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | \theta_\Omega | \phi^H \rangle - \langle \phi^H | \theta_\Omega | \phi^J \rangle \mathbf{S}^\dagger \mathbf{c} + \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | \theta_\Omega | \phi^J \rangle \mathbf{S}^\dagger \mathbf{c} \right\}^{-1} \\ &\quad \times \left(\langle \phi^H | p \rangle - \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | p \rangle \right) \langle \phi | \theta_\Omega | \phi \rangle \langle p | \\ &\approx \left\{ \left(\langle \phi^H | \theta_\Omega | \phi^H \rangle \right)^{-1} - \left(\langle \phi^H | \theta_\Omega | \phi^H \rangle \right)^{-1} \left(-\mathbf{c}^\dagger \mathbf{S} \langle \phi^J | \theta_\Omega | \phi^H \rangle - \langle \phi^H | \theta_\Omega | \phi^J \rangle \mathbf{S}^\dagger \mathbf{c} \right) \left(\langle \phi^H | \theta_\Omega | \phi^H \rangle \right)^{-1} \right\} \\ &\quad \times \left(\langle \phi^H | p \rangle - \mathbf{c}^\dagger \mathbf{S} \langle \phi^J | p \rangle \right) \langle \phi | \theta_\Omega | \phi \rangle \langle p | \\ &\approx \left(\langle \phi^H | \theta_\Omega | \phi^H \rangle \right)^{-1} \left\{ \langle \phi^H | p \rangle - \mathbf{c}^\dagger \mathbf{S} \left[\langle \phi^J | p \rangle - \langle \phi^J | \theta_\Omega | \phi^H \rangle \left(\langle \phi^H | \theta_\Omega | \phi^H \rangle \right)^{-1} \langle \phi^H | p \rangle \right] \right. \\ &\quad \left. + \langle \phi^H | \theta_\Omega | \phi^J \rangle \mathbf{S}^\dagger \mathbf{c} \left(\langle \phi^H | \theta_\Omega | \phi^H \rangle \right)^{-1} \langle \phi^H | p \rangle \right\} \langle \phi | \theta_\Omega | \phi \rangle \langle p |\end{aligned}\tag{6.11}$$

We define the quantity

$$\langle \pi | = \langle \pi | \phi \rangle \langle p | \tag{6.12}$$

Density matrix

$$\rho^X = \langle \pi | \phi \rangle \rho^\phi \langle \phi | \pi \rangle \langle \pi | \phi \rangle \rho^\phi \langle \phi | \pi \rangle$$

6.4 Failed procedure

The procedure described here failed, probably due to inverting a singular matrix.

The projector functions are determined so that

$$\langle p_\zeta | \psi \rangle = \sum_\gamma \langle p_\zeta | \chi_\alpha \rangle \langle \pi_\alpha | \psi \rangle \Rightarrow \langle \pi_\alpha | = \sum_\gamma \left(\langle p | \chi \rangle \right)_{\alpha, \gamma}^{-1} \langle p_\gamma | \tag{6.13}$$

$$\begin{aligned}
\langle p_\zeta | \chi_\gamma \rangle &= \sum_\alpha \left\{ \langle p_\zeta | \phi_\alpha^H \rangle - \sum_\beta \langle p_\zeta | \phi_\beta^T \rangle S_{R_\beta, L_\beta, R_\alpha, L_\alpha}^\dagger c_{\alpha, \gamma} \right\} \\
&= \sum_\eta \left\{ \delta_{\zeta, \eta} - \underbrace{\sum_{\beta, \alpha, \mu} \langle p_\zeta | \phi_\beta^T \rangle S_{R_\beta, L_\beta, R_\alpha, L_\alpha}^\dagger c_{\alpha, \mu} \left(\langle p | \phi^H \rangle \right)_{\mu, \eta}^{-1}}_{=: Z_{\zeta, \eta}^\dagger} \right\} \langle p_\eta | \phi_\gamma^H \rangle \\
\langle p | \chi \rangle &= (\mathbf{1} - \mathbf{Z}^\dagger) \langle p | \phi^H \rangle
\end{aligned} \tag{6.14}$$

with

$$Z_{\zeta, \eta} \stackrel{\text{def}}{=} \left(\langle p | \phi^H \rangle \right)_{\zeta, \mu}^{-1, \dagger} c_{\mu, \alpha}^\dagger S_{\alpha, \beta} \left(\langle p | \phi^J \rangle \right)^\dagger. \tag{6.15}$$

Thus, we obtain the projector functions for the local orbitals as

$$\begin{aligned}
\langle p | \chi \rangle \langle \pi | &= \langle p | \stackrel{\text{Eq. 6.14}}{\Leftrightarrow} \sum_{\beta, \gamma} \left[\delta_{\alpha, \beta} - (Z^\dagger)_{\alpha, \beta} \right] \langle p_\beta | \phi_\gamma^H \rangle \langle \pi_\gamma | = \langle p_\alpha | \\
\Rightarrow \langle \pi_\gamma | &= \sum_{\alpha, \beta} \left(\langle p | \phi^H \rangle \right)_{\gamma, \beta}^{-1} \left[\mathbf{1} - \mathbf{Z}^\dagger \right]_{\beta, \alpha}^{-1} \langle p_\alpha |
\end{aligned} \tag{6.16}$$

6.5 Code structure

Initialization

```
simplelmtomakepotpar1
  simplelmtomakechi1
  simplelmtomatrixelements
  simplelmtomoffxint
```

Iteration:

```
simplelmtomhybrid
  waves$getrspacemata('denmat')
  simplelmtomonsitedenmatextractadd('extract' denmat ->donsite)
  simplelmtomstructureconstants
  simplelmtomzmat1
  SIMPLELMTO_DENMATPHITOCHE1('FWRD',
  !
  SIMPLELMTO_HYBRIDENERGY:(denmat,donsite) ->(etot,hamil,honsite,F)
  !
  SIMPLELMTO_DENMATPHITOCHE1('BACK')
  SIMPLELMTO_ONSITEDENMATRETRACTADD('ADDBACK',hamil+honsite->hamil
  WAVES$SETRSPACEMATA('HAMIL',NND,HAMIL)
```

6.5.1 Local orbitals

In `simplelmtomakechi1` the local orbitals are constructed.

First the head and tail functions are constructed such that

- the nodeless head function matches to the solid Hankel function at the radius `RAUG` with value and derivative.
- the nodeless tail function matches to the solid Bessel function function at the radius `RAUG` with value and derivative.

With the same coefficients we determine the all-electron and pseudo head and tail functions. This implies that the all-electron and pseudo head and tail functions do not obey the matching conditions.

The local orbitals are composed from nodeless head function within the augmentation sphere and solid Hankel function beyond. In a second step we add the difference of the all-electron and pseudo head function to obtain the corresponding local orbitals.

An on-site Gram-Schmidt-type orthonormalization is performed on the local orbitals. The matrix that changes the augmented Hankel functions to the normalized local orbitals is `CMAT`. It is required later to translate the structure constants for Hankel functions to those of local orbitals.

Alongside with the orbitals also the head functions are transformed.

6.5.2 Projections

In `simplelmtomakechi1` we also determine

$$\begin{aligned}\langle \tilde{p}_\alpha | \tilde{\phi}_\beta^H \rangle &= \text{PROPHIH} \\ \langle \tilde{p}_\alpha | \tilde{\phi}_\beta^T \rangle &= \text{PROPHIT}\end{aligned}\tag{6.17}$$

Later we will need the inverse of $\langle p | \phi^H \rangle$. $\langle p | \phi^H \rangle$ is a $N^\phi \times N^\chi$ matrix and \mathbf{c} is a $N^\chi \times N^\chi$ matrix. The inversion shall be done in the least-square sense, that is

$$\left(\langle p | \phi^H \rangle \right)^{-1} = \left(\langle \phi^H | p \rangle \langle \phi | \phi \rangle \langle p | \phi^H \rangle \right)^{-1} \langle \phi^H | p \rangle \langle \phi | \phi \rangle = \text{PROPHIINV}\tag{6.18}$$

ZMAT

First the structure constants are calculated. The calculated structure constants are defined

$$|K_\alpha^\infty\rangle = - \sum_{\beta \in R'} |J_\beta\rangle S_{\alpha,\beta} \text{ for } R' \neq R_\alpha\tag{6.19}$$

The object `ZMAT` is calculated in `SIMPLELMT0ZMAT1`. It is constructed in each iteration because it takes the structure constants, which depend on the atomic structure.

WARNING!! In the code, the variable `ZMAT` is overloaded.

- the off-site terms contain

$$\mathbf{Zmat} = \left(\langle p | \phi^H \rangle \right)^{-1, \dagger} \mathbf{c}^\dagger \mathbf{S} \left(\langle p | \phi^J \rangle \right)^\dagger\tag{6.20}$$

- the on-site terms contain

$$\mathbf{Z}_{\text{mat}} = \left(\mathbf{1} - \sum_{R'} \mathbf{Z}_{R,R'} \mathbf{Z}_{R',R} \right)^{-1} \quad (6.21)$$

The variable \mathbf{Z} does not have non-zero on-site elements, because the on-site matrix elements of the bare structure constants vanish.

6.6 Transformation of density matrix and Hamiltonian

6.6.1 Formally correct derivation

Let me introduce symbols for the density matrix ρ^ϕ in terms of partial waves and the one ρ^χ in terms of local orbitals.

$$\begin{aligned} \rho_{\alpha,\beta}^\phi &:= \langle \tilde{p}_\alpha | \tilde{\psi}_n \rangle f_n \langle \tilde{\psi}_n | \tilde{p}_\beta \rangle \\ \rho_{\alpha,\beta}^\chi &:= \langle \pi_\alpha | \hat{\rho} | \pi_\beta \rangle \end{aligned} \quad (6.22)$$

We use the definition of the orbital projector functions Eq. 6.16 to transform the density matrix from a partial-wave to a local-orbital representation.

$$\rho^\chi = \left(\langle p | \phi^H \rangle \right)^{-1} (\mathbf{1} - \mathbf{Z})^{-1, \dagger} \rho^\phi (\mathbf{1} - \mathbf{Z})^{-1} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1} \quad (6.23)$$

The Hamiltonian is then obtained from the derivatives.

$$\begin{aligned} dE &= \sum_{\alpha,\beta} \frac{dE}{d\rho_{\alpha,\beta}^\chi} d\rho_{\alpha,\beta}^\chi = \sum_{\alpha,\beta} H_{\beta,\alpha}^\chi d\rho_{\alpha,\beta}^\chi = \sum_{\alpha,\beta} H_{\beta,\alpha}^\chi d \left(\sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle \psi_n | \pi_\alpha \rangle \right) \\ &= \sum_{\alpha,\beta} H_{\beta,\alpha}^\chi d \left(\sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle \psi_n | \pi_\alpha \rangle \right) \\ &= \text{Tr} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1} \mathbf{H}^\chi \left(\langle p | \phi^H \rangle \right)^{-1} d \left[(\mathbf{1} - \mathbf{Z})^{\dagger, -1} \left(\sum_n \langle p | \psi_n \rangle f_n \langle \psi_n | p \rangle \right) (\mathbf{1} - \mathbf{Z})^{-1} \right] \\ &= \text{Tr} \underbrace{(\mathbf{1} - \mathbf{Z})^{-1} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1} \mathbf{H}^\chi \left(\langle p | \phi^H \rangle \right)^{-1} (\mathbf{1} - \mathbf{Z})^{\dagger, -1}}_{\mathbf{H}^\phi} \\ &\quad \times \left[d\mathbf{Z}^\dagger (\mathbf{1} - \mathbf{Z})^{\dagger, -1} \rho^\phi + d\rho^\phi + \rho^\phi (\mathbf{1} - \mathbf{Z})^{-1} d\mathbf{Z} \right] \\ &= \text{Tr} [\mathbf{H}^\phi d\rho^\phi] + 2\text{Tr} [\mathbf{H}^\phi \rho^\phi (\mathbf{1} - \mathbf{Z})^{-1} d\mathbf{Z}] \end{aligned} \quad (6.24)$$

The Hamiltonian \mathbf{H}^ϕ describes the correction to the Hamiltonian

$$d\hat{H} = \sum_{\alpha,\beta} |p_\alpha\rangle H_{\alpha,\beta}^\phi \langle p_\beta| \quad (6.25)$$

The second term depends on the atomic positions via the bare structure constants. With Eq. 6.15, we obtain

$$F_j = -\text{Tr} \left\{ \left(\langle p | \phi^T \rangle \right)^\dagger \mathbf{H}^\phi \rho^\phi (\mathbf{1} - \mathbf{Z})^{-1} \left(\langle p | \phi^H \rangle \right)^{-1, \dagger} \mathbf{c}^\dagger \frac{d\mathbf{S}}{dR_j} \right\} \quad (6.26)$$

6.6.2 Coding of direct inversion

First, I convert

$$\sum_{\vec{t}} (\mathbf{1} - \mathbf{Z})_{\alpha, \vec{0}, \beta, \vec{t}} e^{-i\vec{k}\vec{t}} \quad (6.27)$$

then I invert the \mathbf{k} -dependent matrix $\mathbf{X}_{\alpha, \beta}(\vec{k})$ and transform it back

$$X_{\beta, \vec{0}, \alpha, \vec{t}} = \frac{1}{N_k} \sum_{\vec{k}} e^{-i\vec{k}\vec{t}} \left[\sum_{\vec{t}'} (\mathbf{1} - \mathbf{Z})_{\alpha, \vec{0}, \beta, \vec{t}'} e^{-i\vec{k}\vec{t}'} \right]_{\beta, \alpha}^{-1} \quad (6.28)$$

6.6.3 Approximate inversion

We start with Eq. 6.16

$$\langle \pi | = \left(\langle p | \phi^H \rangle \right)^{-1} \left\{ \mathbf{1} - \mathbf{Z} \right\}^{-1, \dagger} \langle p | \quad (6.29)$$

The matrix \mathbf{Z} is not hermitean, nor does it have the same dimensions for the two dimensions. According to the derivation Eq. 6.14, we need the left-handed inverse of $\mathbf{1} - \mathbf{Z}^\dagger$, respectively the transpose of the right-handed inverse of $\mathbf{1} - \mathbf{Z}$.

Instead of the complete inversion of $\mathbf{1} - \mathbf{Z}$, I approximate it by downfolding: For this purpose, I divide the system in an on-site term and the surrounding, which is then treated as bath.

$$\begin{pmatrix} \mathbf{1} & -\mathbf{Z}_{AB} \\ -\mathbf{Z}_{BA} & \mathbf{Y} \end{pmatrix} \begin{pmatrix} \mathbf{A}_{AA} & \mathbf{A}_{AB} \\ \mathbf{A}_{BA} & \mathbf{A}_{BB} \end{pmatrix} = \begin{pmatrix} \mathbf{1}_{AA} & \mathbf{0}_{AB} \\ \mathbf{0}_{BA} & \mathbf{1}_{BB} \end{pmatrix} \quad (6.30)$$

The matrix \mathbf{Z} is not hermitean. The matrix \mathbf{Y} is defined as

$$\mathbf{Y} := \mathbf{1}_{BB} - \mathbf{Z}_{BB} \quad (6.31)$$

Below, the approximations will be introduced by “tampering” with \mathbf{Y} . The onsite terms of \mathbf{Z} are zero.

Before I work out \mathbf{A} , let me investigate which parts of \mathbf{A} are actually required:

$$\begin{aligned} |p\rangle &= |\pi\rangle \left(\langle p | \phi^H \rangle \right)^\dagger (\mathbf{1} - \mathbf{Z}) \\ |p\rangle \mathbf{A} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1} &= |\pi\rangle \underbrace{\left(\langle p | \phi^H \rangle \right)^\dagger (\mathbf{1} - \mathbf{Z}) \mathbf{A} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1}}_1 \\ \langle \pi | &= \left(\langle p | \phi^H \rangle \right)^{-1} \mathbf{A}^\dagger \langle p | \\ \langle \pi_A | &= \left(\langle p_A | \phi_A^H \rangle \right)^{-1} \left(\mathbf{A}_{AA} \right)^\dagger \langle p_A | + \left(\langle p_A | \phi_A^H \rangle \right)^{-1} \left(\mathbf{A}_{BA} \right)^\dagger \langle p_B | \end{aligned} \quad (6.32)$$

Now I work out \mathbf{A} , the inverse of $\mathbf{1} - \mathbf{Z}$ by resolving Eq. 6.30.

$$\begin{aligned} \mathbf{A}_{AA} - \mathbf{Z}_{AB} \mathbf{A}_{BA} &= \mathbf{1}_{AA} \\ \mathbf{A}_{AB} - \mathbf{Z}_{AB} \mathbf{A}_{BB} &= \mathbf{0}_{AB} \\ -\mathbf{Z}_{BA} \mathbf{A}_{AA} + \mathbf{Y}_{BB} \mathbf{A}_{BA} &= \mathbf{0}_{BA} \\ -\mathbf{Z}_{BA} \mathbf{A}_{AB} + \mathbf{Y}_{BB} \mathbf{A}_{BB} &= \mathbf{1}_{BB} \end{aligned} \quad (6.33)$$

The third equation links \mathbf{A}_{BA} to \mathbf{A}_{AA}

$$\mathbf{A}_{BA} = \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA} \mathbf{A}_{AA} \quad (6.34)$$

Insertion into the first equation produces an equation for \mathbf{A}_{AA}

$$\begin{aligned} (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA}) \mathbf{A}_{AA} &= \mathbf{1}_{AA} \\ \Rightarrow \mathbf{A}_{AA} &= (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1} \end{aligned} \quad (6.35)$$

and \mathbf{A}_{BA}

$$\mathbf{A}_{BA} = \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA} (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1} \quad (6.36)$$

$$\begin{aligned} \langle \pi_A | &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{A}_{AA})^\dagger \langle p_A | + (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{A}_{BA})^\dagger \langle p_B | \\ &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} \langle p_A | \\ &\quad + (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} \langle p_B | \\ &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} [\langle p_A | + (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} \langle p_B |] \end{aligned} \quad (6.37)$$

Test: I test the biorthogonality condition $\langle \pi_\alpha | \chi_\beta \rangle = \delta_{\alpha, \beta}$ using

$$|\chi\rangle = (|\phi^K\rangle - |\phi^J\rangle \mathbf{S}^\dagger) \mathbf{c} = |\phi^H\rangle - |\phi^J\rangle \mathbf{S}^\dagger \mathbf{c} \quad (6.38)$$

$$\begin{aligned} \langle \pi_A | \chi \rangle &= \left\{ (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} \right. \\ &\quad \times \left. [\langle p_A | + (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} \langle p_B |] \right\} \left\{ |\phi^H\rangle - |\phi^J\rangle \mathbf{S}^\dagger \mathbf{c} \right\} \\ &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} \\ &\quad \times \begin{cases} \langle p_A | \phi_A^H \rangle - (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} \langle p_B | \phi_B^J \rangle (\mathbf{S}_{AB})^\dagger \mathbf{c}_{AA} & \text{for } \chi \in A \\ -\langle p_A | \phi_A^J \rangle (\mathbf{S}_{B,A})^\dagger \mathbf{c}_{BB} + (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} [\langle p_B | \phi_B^H \rangle - \langle p_B | \phi_B^J \rangle (\mathbf{S}_{BB})^\dagger \mathbf{c}_{BB}] & \text{for } \chi \in B \end{cases} \\ &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} \\ &\quad \times \begin{cases} [\mathbf{1}_{AA} - (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} (\mathbf{Z}_{AB})^\dagger] \langle p_A | \phi_A^H \rangle & \text{for } \chi \in A \\ -(\mathbf{Z}_{BA})^\dagger (\langle p_B | \phi_B^H \rangle + (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} [\langle p_B | \phi_B^H \rangle - (\mathbf{Z}_{BB})^\dagger (\langle p_B | \phi_B^H \rangle)]) & \text{for } \chi \in B \end{cases} \\ &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Y}_{BB}^{-1} \mathbf{Z}_{BA})^{-1, \dagger} \\ &\quad \times \begin{cases} [\mathbf{1}_{AA} - (\mathbf{Z}_{BA})^\dagger (\mathbf{Y}_{BB})^{-1, \dagger} (\mathbf{Z}_{AB})^\dagger] \langle p_A | \phi_A^H \rangle & \text{for } \chi \in A \\ (\mathbf{Z}_{BA})^\dagger \left\{ -\mathbf{1} + (\mathbf{Y}_{BB})^{-1, \dagger} [\mathbf{1} - (\mathbf{Z}_{BB})^\dagger] \right\} (\langle p_B | \phi_B^H \rangle) & \text{for } \chi \in B \end{cases} \end{aligned} \quad (6.39)$$

We have purposely not used the identity $\mathbf{Y} = (\mathbf{1}_{BB} - \mathbf{Z}_{BB})^{-1}$, because this identity will be violated when we approximate \mathbf{Y} .

If the identity holds one can show in a few additional steps that $\langle \pi_A | \chi_A \rangle = \mathbf{1}_{AA}$ and $\langle \pi_A | \chi_B \rangle = \mathbf{0}_{AB}$.

Approximating \mathbf{Y} by the unit matrix

In the limit of vanishing orbital overlap, the off-site terms of \mathbf{Z} vanish and $\mathbf{Y}_{22} = \mathbf{1}$.

$$\begin{aligned} \langle \pi_A | &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1}_{AA} - \mathbf{Z}_{AB} \mathbf{Z}_{BA})^{-1, \dagger} [\langle p_A | + (\mathbf{Z}_{BA})^\dagger \langle p_B |] \\ \Leftrightarrow \quad \langle \pi | &= (\langle p | \phi^H \rangle)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA})^{-1, \dagger} (\mathbf{1} + \mathbf{Z})^\dagger \langle p | \end{aligned} \quad (6.40)$$

When the matrix \mathbf{Y}_{22} is set equal to the unit matrix, we obtain

$$\begin{aligned} \langle \pi_A | \chi \rangle &= (\langle p_A | \phi_A^H \rangle)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA})^{-1, \dagger} \\ &\times \begin{cases} [\mathbf{1}_{AA} - (\mathbf{Z}_{BA})^\dagger (\mathbf{Z}_{AB})^\dagger] \langle p_A | \phi_A^H \rangle & \text{for } \chi \in A \\ (\mathbf{Z}_{BA})^\dagger \left\{ -\mathbf{1} + [\mathbf{1} - (\mathbf{Z}_{BB})^\dagger] \right\} \langle p_B | \phi_B^H \rangle & \text{for } \chi \in B \end{cases} \\ &= \begin{cases} \mathbf{1}_{AA} & \text{for } \chi \in A \\ (\langle p_A | \phi_A^H \rangle)^{-1} \left([\mathbf{0}_{BA} - \mathbf{Z}_{BB} \mathbf{Z}_{BA}] (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA})^{-1} \right)^\dagger \langle p_B | \phi_B^H \rangle & \text{for } \chi \in B \end{cases} \end{aligned} \quad (6.41)$$

Approximating the conversion of the density matrix

As long as we do not approximate \mathbf{Y} , the bi-orthogonality is exactly obeyed. However, we need to evaluate the inverse of $\mathbf{Y} := \mathbf{1} - \mathbf{Z}_{BB}$. Secondly the conversion of the density matrix involves complicated multicenter sums, which produce long-ranged contributions. The goal is to truncate the multicenter sum, but in way that is as consistent as possible. This is probably better than just truncating the sum so that the neighborlist is filled up.

- We approximate $\mathbf{Y} \approx \mathbf{1}_{BB}$.
- Ontop of this approximation, when we convert the density matrix from a partial-wave representation into a local-orbital representation, we ignore
 - all onsite terms, that contain more than two off-site terms of ρ or \mathbf{Z} in a product and we ignore
 - all off-site terms with more than one off-site term.

This approximation limits all products to terms that exist only on one atom pair at a time. This seems to be consistent with the first approximation setting $\mathbf{Y} = \mathbf{1}$.

$$(\mathbf{1} + \mathbf{Z})^\dagger \rho^\phi (\mathbf{1} + \mathbf{Z}) \approx \begin{cases} \rho_{\text{onsite}}^\phi + \mathbf{Z}^\dagger \rho_{\text{offsite}}^\phi + \rho_{\text{offsite}}^\phi \mathbf{Z} + \mathbf{Z}^\dagger \rho_{\text{onsite}}^\phi \mathbf{Z} & \text{onsite} \\ \rho_{\text{offsite}}^\phi + \mathbf{Z}^\dagger \rho_{\text{onsite}}^\phi + \rho_{\text{on-site}}^\phi \mathbf{Z} & \text{offsite} \end{cases} \quad (6.42)$$

Note that the onsite terms of \mathbf{Z} are zero.

The error terms have the form

$$\Delta \approx \begin{cases} \mathbf{Z} \boldsymbol{\rho}_{\text{offsite}}^\phi \mathbf{Z}^\dagger & \text{onsite} \\ \mathbf{Z} \left(\boldsymbol{\rho}_{\text{onsite}}^\phi + \boldsymbol{\rho}_{\text{offsite}}^\phi \right) \mathbf{Z}^\dagger & \text{offsite} \end{cases} \quad (6.43)$$

Transformation of density matrix and Hamiltonian

$$\begin{aligned} \boldsymbol{\rho}^\chi &= \left(\langle p | \phi^H \rangle \right)^{-1} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA})^{-1, \dagger} \underbrace{\left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right)}_{\text{special}} (\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA})^{-1} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1} \\ dE &= \sum_{\alpha, \beta} \frac{dE}{d\rho_{\alpha, \beta}^\chi} d\rho_{\alpha, \beta}^\chi = \sum_{\alpha, \beta} H_{\beta, \alpha}^\chi d\rho_{\alpha, \beta}^\chi = \sum_{\alpha, \beta} H_{\beta, \alpha}^\chi d \left(\sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle \psi_n | \pi_\alpha \rangle \right) \\ &= \sum_{\alpha, \beta} H_{\beta, \alpha}^\chi d \left(\sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle \psi_n | \pi_\alpha \rangle \right) \\ &= \text{Tr} \left[\left(\langle p | \phi^H \rangle \right)^{\dagger, -1} \mathbf{H}^\chi \left(\langle p | \phi^H \rangle \right)^{-1} d \left[\left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1, \dagger} \underbrace{\left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right)}_{\text{special}} \left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1} \right] \right] \\ &= \text{Tr} \left[\underbrace{\left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1} \left(\langle p | \phi^H \rangle \right)^{\dagger, -1} \mathbf{H}^\chi \left(\langle p | \phi^H \rangle \right)^{-1} \left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1, \dagger}}_{\tilde{H}} \right. \\ &\quad \times \left[d \left(\mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^\dagger \left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1, \dagger} \underbrace{\left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right)}_{\text{special}} + \underbrace{d \mathbf{Z}^\dagger \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right)}_{\text{special}} \right. \\ &\quad \left. \left. + \underbrace{\left(\mathbf{1} + \mathbf{Z} \right)^\dagger d \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right)}_{\text{special}} \right. \right. \\ &\quad \left. \left. + \underbrace{\left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi d \mathbf{Z}}_{\text{special}} + \underbrace{\left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right)}_{\text{special}} \left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1} d \left(\mathbf{Z}_{AB} \mathbf{Z}_{BA} \right) \right] \right] \\ &= \text{Tr} \left[\left(\mathbf{1} + \mathbf{Z} \right) \tilde{H} \left(\mathbf{1} + \mathbf{Z} \right)^\dagger d \boldsymbol{\rho}^\phi \right] + 2 \text{Tr} \left[\tilde{H} \left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi d \mathbf{Z} \right] \\ &\quad + 2 \text{Tr} \left[\tilde{H} \left(\mathbf{1} + \mathbf{Z} \right)^\dagger \boldsymbol{\rho}^\phi \left(\mathbf{1} + \mathbf{Z} \right) \left(\mathbf{1} - \mathbf{Z}_{AB} \mathbf{Z}_{BA} \right)^{-1} d \left(\mathbf{Z}_{AB} \mathbf{Z}_{BA} \right) \right] \end{aligned} \quad (6.44)$$

6.6.4 Otherstuff

This is a suggestion for later: We will calculate the on-site matrix

$$\begin{aligned}
X_{\alpha,\beta}(\vec{R}_2 - \vec{R}_1) &= \sum_{\gamma} \mathbf{Z}_{\alpha,\gamma}(\vec{R}_2 - \vec{R}_1) \mathbf{Z}_{\gamma,\beta}(\vec{R}_1 - \vec{R}_2) \\
&= \sum_{\eta,\mu \in R_1} \sum_{\gamma,\gamma' \in R_2} \left[\langle p|\phi^H\rangle^{-1,\dagger} \mathbf{c}^\dagger \right]_{\alpha,\eta} S_{R_\eta,L_\eta,R_\gamma,L_\gamma} \\
&\quad \times \left[\langle p|\phi^J\rangle^\dagger \langle p|\phi^H\rangle^{-1,\dagger} \mathbf{c}^\dagger \right]_{\gamma,\gamma'} S_{R_{\gamma'},L_{\gamma'},R_\mu,L_\mu} \left[\langle p|\phi^J\rangle^\dagger \right]_{\mu,\beta} \\
&= \sum_{\eta,\mu \in R_1} \sum_{\gamma,\gamma' \in R_2} \left[\mathbf{c} \langle p|\phi^H\rangle^{-1} \right]_{\alpha,\eta}^\dagger S_{R_\eta,L_\eta,R_\gamma,L_\gamma} \\
&\quad \times \left[\mathbf{c} \langle p|\phi^H\rangle^{-1} \langle p|\phi^J\rangle \right]_{\gamma,\gamma'}^\dagger S_{R_{\gamma'},L_{\gamma'},R_\mu,L_\mu} \left[\langle p|\phi^J\rangle \right]_{\mu,\beta}^\dagger
\end{aligned} \tag{6.45}$$

$$\begin{aligned}
S_{RL,R'L'} &\stackrel{\text{Eq. 2.7}}{=} (-1)^{\ell'+1} 4\pi \sum_{L''} C_{L,L',L''} H_{L''}(\vec{R}' - \vec{R}) \begin{cases} (-ik)^{\ell'+\ell''-\ell''} & \text{for } k^2 > 0 \\ \delta_{\ell'+\ell''-\ell''} & \text{for } k^2 = 0 \\ \kappa^{\ell'+\ell''-\ell''} & \text{for } k^2 = -\kappa^2 < 0 \end{cases} \\
S_{R'L',RL} &= S_{RL,R'L'} \\
S_{RL,R'L'} S_{R'L',RL''} &= (-1)^{\ell'+\ell''} (4\pi)^2 \sum_{\bar{L}} \sum_{\bar{L}'} C_{L,L',\bar{L}} C_{L',L'',\bar{L}'} H_{\bar{L}}(\vec{R}' - \vec{R}) H_{\bar{L}'}(\vec{R} - \vec{R}') \\
&\quad \times \begin{cases} (-ik)^{2\ell+2\ell'-\bar{\ell}-\bar{\ell}'} & \text{for } k^2 > 0 \\ \delta_{\ell+\ell'-\bar{\ell}} \delta_{\ell'+\ell''-\bar{\ell}'} & \text{for } k^2 = 0 \\ \kappa^{2\ell+2\ell'-\bar{\ell}-\bar{\ell}'} & \text{for } k^2 = -\kappa^2 < 0 \end{cases} \\
&= (-1)^{\ell'+\ell''} (4\pi)^2 \sum_{\bar{L}} \sum_{\bar{L}'} C_{L,L',\bar{L}} C_{L',L'',\bar{L}'} H_{\bar{L}}(\vec{R}' - \vec{R}) H_{\bar{L}'}(\vec{R} - \vec{R}') \\
&\quad \times \begin{cases} (-ik)^{2\ell+2\ell'-\bar{\ell}-\bar{\ell}'} & \text{for } k^2 > 0 \\ \delta_{\ell+\ell'-\bar{\ell}} \delta_{\ell'+\ell''-\bar{\ell}'} & \text{for } k^2 = 0 \\ \kappa^{2\ell+2\ell'-\bar{\ell}-\bar{\ell}'} & \text{for } k^2 = -\kappa^2 < 0 \end{cases}
\end{aligned} \tag{6.46}$$

Let me denote the radial part of $H_L(\vec{R}) = h_\ell(|\vec{r}|) Y_L(\vec{R})$.

$$\begin{aligned}
S_{RL,R'L'} S_{R'L',RL''} &= (-1)^{\ell'+\ell''} (4\pi)^2 \sum_{\bar{\ell},\bar{\ell}'} h_{\bar{\ell}}(|\vec{R}' - \vec{R}|) h_{\bar{\ell}'}(|\vec{R}' - \vec{R}|) \\
&\quad \times \sum_{\bar{m},\bar{m}'} C_{L,L',\bar{L}} C_{L',L'',\bar{L}'} Y_{\bar{L}}(\vec{R}' - \vec{R}) (-1)^{\bar{\ell}} Y_{\bar{L}'}(\vec{R}' - \vec{R}) \\
&\quad \times \begin{cases} (-ik)^{2\ell+2\ell'-\bar{\ell}-\bar{\ell}'} & \text{for } k^2 > 0 \\ \delta_{\ell+\ell'-\bar{\ell}} \delta_{\ell'+\ell''-\bar{\ell}'} & \text{for } k^2 = 0 \\ \kappa^{2\ell+2\ell'-\bar{\ell}-\bar{\ell}'} & \text{for } k^2 = -\kappa^2 < 0 \end{cases}
\end{aligned} \tag{6.47}$$

6.7 Analysis of the code structure in paw_lmto

PAW_WAVES

```
CALL LMTO$PROJTONTBO_NEW('FWRD',XK(:,IKPT),NDIM,NBH,NPRO,THIS\%PROJ &
&                                ,NORB,THIS\%TBC_NEW)
```

The subroutine lmto\$etot is called from the paw_waves1 object.

```
LMTO$ETOT(LMNXX_,NDIMD_,NAT_,DENMAT_,DH_)
  LMTO_HYBRID()
    CALL LMTO_NTBO_DENMAT_NEW()
    CALL LMTO_HYBRIDENERGY()
    CALL LMTO_NTBO_DENMAT_DER_NEW()
    CALL LMTO_CLEAN_DENMAT_NEW()
```

6.8 New code

6.8.1 Determine off-site density matrix in partial-wave representation

With a routine derived from LMTO_NTBO_DENMAT_NEW, I determine the density matrix in terms of partial-wave projector functions.

$$\rho_{\alpha,\vec{0},\beta,\vec{t}} = \frac{1}{\sum_k} \sum_k \sum_{n=1}^{nb} \langle \tilde{p}_{\alpha,0} | \psi_{n,k} \rangle f_{n,k} \langle \psi_{n,k} | \tilde{p}_{\beta,\vec{0}} \rangle e^{i\vec{k}\vec{t}} \quad (6.48)$$

There is a data type for matrices represented by a neighborlist. The third dimension describes the spin indices.¹

```
TYPE NGBRMAT_TYPE
  INTEGER(4)      :: IAT1  ! FIRST ATOM (LINKED TO THE LEFT INDEX OF MAT)
  INTEGER(4)      :: IAT2  ! SECOND ATOM (LINKED TO THE RIGHT INDEX OF MAT)
  INTEGER(4)      :: IT(3) ! LATTICE TRANSLATIONS TO BE ADDED TO ATOM 2
  INTEGER(4)      :: N1    ! RIGHT DIMENSION OF MAT
  INTEGER(4)      :: N2    ! LEFT DIMENSION OF MAT
  INTEGER(4)      :: N3    ! THIRD DIMENSION OF MAT
  REAL(8),POINTER :: MAT(:, :, :) ! (N1,N2,N3)
END TYPE NGBRMAT_TYPE
```

In a closed-packed sphere packing there are 12 nearest neighbors.

¹

- non spin-polarized, $N3 = 1$.
- spin polarized, $N3 = 2$. The two matrices are $\sigma = \uparrow, \downarrow$
- non-collinear, $N3 = 3$. The three matrices are the projections onto S_x, S_y, S_z .

6.8.2 Density matrix in orbital representation

The orbital projections are evaluated using an onsite transformation

$$\langle \tilde{\pi}_\alpha | = \sum_{j \in R_\alpha} c_{\alpha,j} \langle \tilde{p}_j | \quad \text{with} \quad \langle \tilde{\pi}_\alpha | \tilde{\chi}_\beta \rangle = \sum_{j \in R_\alpha} c_{\alpha,j} \langle \tilde{p}_j | \tilde{\chi}_\beta \rangle \quad (6.49)$$

When the orbitals are frozen, the matrix \mathbf{c} is calculated once and for all with singular value decomposition. (Note that \mathbf{c} is not square.)

The restriction to onsite terms requires that the orbitals fall off with distance sufficiently fast, that the offsite terms of $\langle \tilde{p}_j | \tilde{\chi}_\alpha \rangle$ are negligible.

6.8.3 Bare structure constants

```
LM1X=(L1X+1)**2
LM2X=(L2X+1)**2
CALL LMT0$STRUCTURECONSTANTS(R2-R1,K2,L1X,L2X,S(:LM1X,:LM2X))
```

6.8.4 Routine radial\$helmholtz

The routine radial\$helmholtz shall act like radial\$poisson but with a Yukawa potential.

6.9 Helmholtz equation

<http://fortranwiki.org/fortran/show/Bessel+function>

Yukawa Potential obeys the Helmholtz equation: In RPA without local field effects, the Energy looks like that of a Hartree Fock calculation, but with a modified exchange term. In the exchange, the interaction is changed to the screened interaction, which is represented by a Yukawa potential.

$$v(\vec{r}) = \int d^3 r' \frac{e^2 n(\vec{r}')}{4\pi\epsilon_0 |\vec{r} - \vec{r}'| \exp(\lambda |\vec{r} - \vec{r}'|)} \quad (6.50)$$

The Yukawa potential satisfies the Helmholtz equation ²

$$\left(\vec{\nabla}^2 - \lambda^2\right) \frac{e^{-\lambda|\vec{r}|}}{|\vec{r}|} = -4\pi\delta(\vec{r}) \quad (6.53)$$

The Helmholtz equation has the form

$$\begin{aligned} \left(\nabla^2 - \lambda^2\right) v(\vec{r}) &= \int d^3 r' \frac{e^2 n(\vec{r}')}{\epsilon_0} \left(\nabla^2 - \lambda^2\right) \frac{1}{4\pi |\vec{r} - \vec{r}'| \exp(\lambda |\vec{r} - \vec{r}'|)} \\ &= - \int d^3 r' \frac{e^2 n(\vec{r}')}{\epsilon_0} \delta(\vec{r} - \vec{r}') \\ &= - \frac{e^2}{\epsilon_0} n(\vec{r}) \end{aligned} \quad (6.54)$$

2

$$\begin{aligned} \vec{\nabla} \frac{1}{r} e^{-\lambda r} &= \left(-\frac{1}{r^2} - \frac{1}{r} \lambda\right) e^{-\lambda r} \frac{\vec{r}}{r} = -(\lambda r + 1) \frac{\vec{r}}{r^3} e^{-\lambda r} \\ \vec{\nabla}^2 \frac{1}{|\vec{r}|} e^{-\lambda|\vec{r}|} &= \vec{r} \left[-\lambda \frac{1}{r^3} e^{-\lambda r} + (\lambda r + 1) \frac{3}{r^4} e^{-\lambda r} + (\lambda r + 1) \frac{1}{r^3} \lambda e^{-\lambda r} \right] \frac{\vec{r}}{r} - (\lambda r + 1) \frac{3}{r^3} e^{-\lambda r} \\ &= \left(-\lambda r + 3(\lambda r + 1) + (\lambda r + 1) \lambda r - (\lambda r + 1) 3\right) \frac{1}{r^3} e^{-\lambda r} \\ &= \left(-\lambda r + 3\lambda r + 3 + (\lambda r)^2 + \lambda r - 3\lambda r - 3\right) \frac{1}{r^3} e^{-\lambda r} = \lambda^2 \frac{1}{r} e^{-\lambda r} \end{aligned} \quad (6.51)$$

In order to verify that the inhomogeneity is a delta function we consider the integral

$$\begin{aligned} \int_{r < r_c} d^3 r \left(\vec{\nabla}^2 - \lambda^2\right) \frac{1}{r} e^{-\lambda r} &= \int_{r=r_c} d\vec{A} \vec{\nabla} \frac{1}{r} e^{-\lambda r} - \lambda^2 4\pi \int_0^{r_c} dr r^2 \frac{1}{r} e^{-\lambda r} \\ &= - \underbrace{\int_{r=r_c} d\vec{A} \frac{\vec{r}}{r^3} (\lambda r + 1) e^{-\lambda r}}_{4\pi} - \lambda^2 4\pi \int_0^{r_c} dr \underbrace{r e^{-\lambda r}}_{\frac{-1}{\lambda^2} \partial_r (\lambda r + 1) e^{-\lambda r}} \\ &= -4\pi (\lambda r_c + 1) e^{-\lambda r_c} + 4\pi \left((\lambda r_c + 1) e^{-\lambda r_c} - 1\right) = -4\pi \end{aligned} \quad (6.52)$$

Radial Green function: In order to obtain the radial Green's function, we expand the density and potential in spherical harmonics

$$\begin{aligned} (\nabla^2 - \lambda^2) v_L(|\vec{r}|) Y_L(\vec{r}) &= -\frac{e^2}{\epsilon_0} n_L(|\vec{r}|) Y_L(\vec{r}) \\ \left(\frac{1}{r} \partial_r^2 r - \frac{\ell(\ell+1)}{r^2} - \lambda^2 \right) v_L(|\vec{r}|) &= -\frac{e^2}{\epsilon_0} n_L(|\vec{r}|) \end{aligned} \quad (6.55)$$

Determine a radial Green's function

$$\left(\frac{1}{r} \partial_r^2 r - \frac{\ell(\ell+1)}{r^2} - \lambda^2 \right) g_\ell(r, r') = \delta(r - r') \quad (6.56)$$

and make the following ansatz for the potential.

$$v_L(r) = \int_0^\infty dr' g_\ell(r, r') \frac{e^2 n_L(r')}{\epsilon_0} \quad (6.57)$$

Let $j_\ell(r)$ be the regular solution and $h_\ell(r)$ the irregular solution, which goes to zero at infinity. The $j_\ell(r)$ is, up to a constant factor, equal to the modified spherical Bessel function of the first kind with argument κr and $h_\ell(r)$ is, up to a constant factor, the modified bessel function of the third kind with argument κr . The factor is chosen so that they have the κ -independent limit for small arguments.

$$j_\ell(r) = \frac{1}{(2\ell+1)!!} r^\ell (1 + O(r^2)) \quad \text{and} \quad h_\ell(r) = (2\ell-1)!! r^{-\ell-1} (1 + O(r^2)) \quad (6.58)$$

In the following, I will refer to the $j_\ell(r)$ as Bessel functions and to the $h_\ell(r)$ as Hankel functions.

The Green's function must solve the homogeneous equation for $r \neq r'$, and it must be continuous but not differentiable for $r = r'$. This is accomplished with the ansatz

$$g_\ell(r, r') = A(r') \left[\theta(r' - r) \frac{j_\ell(r)}{j_\ell(r')} + \theta(r - r') \frac{h_\ell(r)}{h_\ell(r')} \right] \quad (6.59)$$

The factor $A(r)$ is then determined by

$$\lim_{\epsilon \rightarrow 0} \int_{r'-\epsilon}^{r'+\epsilon} dr \frac{1}{r} \partial_r^2 r g_L(r, r') = 1 \quad (6.60)$$

which leads to the following condition

$$\begin{aligned} 1 &= \lim_{\epsilon \rightarrow 0} \frac{1}{r'} \left[\partial_r r g_L(r, r') \right]_{r'-\epsilon}^{r'+\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{1}{r'} \left[g_L(r, r') + r \partial_r g_L(r, r') \right]_{r'-\epsilon}^{r'+\epsilon} \\ &= \partial_r|_{r'+\epsilon} g_L(r, r') - \partial_r|_{r'-\epsilon} g_L(r, r') \\ &= A(r') \left[\delta(r - r') \frac{h_\ell(r)}{h_\ell(r')} + \theta(r' - r) \frac{\partial_r h_\ell(r')}{h_\ell(r')} - \delta(r' - r) \frac{j_\ell(r)}{j_\ell(r')} - \theta(r' - r') \partial_r \frac{j_\ell(r')}{j_\ell(r')} \right] \\ &= A(r') \left[\frac{\partial_r h_\ell(r')}{h_\ell(r')} - \frac{\partial_r j_\ell(r')}{j_\ell(r')} \right] \\ &= A(r') \left[\frac{j_\ell(r') \partial_r h_\ell(r') - h_\ell(r') \partial_r j_\ell(r')}{j_\ell(r') h_\ell(r')} \right] \end{aligned} \quad (6.61)$$

The numerator can be simplified as follows: We can use the Helmholtz equation to show that the following integral vanishes. The origin has been excluded, because K_L is not differentiable, so that Gauss theorem cannot be applied there

$$\begin{aligned}
0 &= \int_{r_1 < |\vec{r}| < r_2} d^3r \left(J_L \vec{\nabla}^2 K_L - K_L \vec{\nabla}^2 J_L \right) \\
&= \int_{r_1 < |\vec{r}| < r_2} d^3r \vec{\nabla} \left(J_L \vec{\nabla} K_L - K_L \vec{\nabla} J_L \right) \\
&= \int_{|\vec{r}|=r_2} d\vec{A} \left(J_L \vec{\nabla} K_L - K_L \vec{\nabla} J_L \right) - \int_{|\vec{r}|=r_1} d\vec{A} \left(J_L \vec{\nabla} K_L - K_L \vec{\nabla} J_L \right) \quad (6.62)
\end{aligned}$$

Thus the surface integral is independent of the radius. I name its value C

$$\begin{aligned}
C &= \int_{|\vec{r}|=r_2} d\vec{A} \left(J_L \vec{\nabla} K_L - K_L \vec{\nabla} J_L \right) \\
&= \int_{|\vec{r}|=r_2} d\vec{A} \left[j_\ell Y_L \left(h_\ell \vec{\nabla} Y_L + Y_L \vec{\nabla} h_\ell \right) - h_\ell Y_L \left(j_\ell \vec{\nabla} Y_L + Y_L \vec{\nabla} j_\ell \right) \right] \\
&= \int_{|\vec{r}|=r_2} d\vec{A} \underbrace{Y_L Y_L}_{\sum_{L''} C_{L'',L,L} Y_{L''}} \left[j_\ell \vec{\nabla} h_\ell - h_\ell \vec{\nabla} j_\ell \right] \\
&= r^2 \left(j_\ell \partial_r h_\ell - h_\ell \partial_r j_\ell \right) \quad (6.63)
\end{aligned}$$

The Hankel and Bessel functions are defined³ such that

$$\begin{aligned}
j_\ell(r) &= \frac{1}{(2\ell+1)!!} r^\ell + \dots \\
h_\ell(r) &= (2\ell-1)!! r^{-\ell-1} + \dots \quad (6.64)
\end{aligned}$$

so that

$$C = r^2 \frac{1}{2\ell+1} \left((-\ell-1)r^{-2} - \ell r^{-2} \right) = -1 \quad (6.65)$$

$$A(\vec{r}') = \frac{j(\vec{r}') h_\ell(r')}{j(\vec{r}') \partial_r h_\ell(r') - h_\ell(r') \partial_r j_\ell(r')} = -r^2 j(\vec{r}') h_\ell(r') \quad (6.66)$$

We insert this result into the expression for the Green function

$$\begin{aligned}
g_\ell(r, r') &= A(r') \left[\theta(r' - r) \frac{j_\ell(r)}{j(\vec{r}')} + \theta(r - r') \frac{h_\ell(r)}{h_\ell(r')} \right] \\
&= -r'^2 \left[\theta(r' - r) j_\ell(r) h(\vec{r}') + \theta(r - r') h_\ell(r) j_\ell(r') \right] \quad (6.67)
\end{aligned}$$

³Methods, Section: Working with spherical Hankel and Bessel functions

In practice we calculate the following steps

$$\begin{aligned}
v_l(r) &= \int_0^\infty dr' g_l(r, r') \frac{e^2 n_L(r')}{\epsilon_0} \\
&= -4\pi \frac{e^2}{4\pi\epsilon_0} \int_0^\infty dr' r'^2 \left[j_l(r) \theta(r' - r) h_l(\vec{r}') + h_l(r) \theta(r - r') j_l(r') \right] n_L(r') \\
&= j_l(r) \left[-4\pi \frac{e^2}{4\pi\epsilon_0} \int_0^\infty dr' r'^2 \theta(r' - r) h_l(\vec{r}') \frac{e^2 n_L(r')}{\epsilon_0} \right] \\
&\quad + h_l(r) \left[-4\pi \frac{e^2}{4\pi\epsilon_0} \int_0^\infty dr' r'^2 \theta(r - r') j_l(r') n_L(r') \right] \\
&= j_l(r) \left[-4\pi \frac{e^2}{4\pi\epsilon_0} \int_0^r dr' r'^2 h_l(\vec{r}') n_L(r') \right] + h_l(r) \left[-4\pi \frac{e^2}{4\pi\epsilon_0} \int_r^\infty dr' r'^2 j_l(r') n_L(r') \right] \\
&= j_l(r) \left[a(\infty) - a(r) \right] + h_l(r) b(r) \\
a(r) &= -4\pi \frac{e^2}{4\pi\epsilon_0} \int_0^r dr' r'^2 h_l(\vec{r}') n_L(r') \\
b(r) &= -4\pi \frac{e^2}{4\pi\epsilon_0} \int_0^r dr' r'^2 j_l(r') n_L(r') \tag{6.68}
\end{aligned}$$

I use the routines SPFUNCTION\$MODBESSEL(L,K*X,Y,DYDX) (ABRAMOWITZ 10.2.4) and SPFUNCTION\$MODHANKEL(L,K*X,Y,DYDX) (ABRAMOWITZ 10.2.4) (which is taken from routine LMT0\\$\solidbessel LMT0\\$\solidhankel in paw_lmtobasics

Chapter 7

Off-site matrix elements NDDO

7.1 Exchange energy and U-tensor

The exchange energy with a screened interaction has the form

$$\begin{aligned}
 E_{xc} &= -\frac{1}{2} \sum_{n,m} f_n f_m \int d^3 r \int d^3 r' \psi_n^*(\vec{r}) \psi_m(\vec{r}) \frac{e^2 e^{-\lambda|\vec{r}-\vec{r}'|}}{4\pi|\vec{r}-\vec{r}'|} \psi_m^*(\vec{r}') \psi_n(\vec{r}') \\
 &= -\frac{1}{2} \sum_{n,m} f_n f_m \sum_{\alpha,\beta,\gamma,\delta} \langle \psi_n | \pi_\alpha \rangle \langle \pi_\gamma | \psi_m \rangle \langle \psi_m | \pi_\beta \rangle \langle \pi_\delta | \psi_n \rangle \\
 &\quad \times \underbrace{\int d^3 r \int d^3 r' \chi_\alpha^*(\vec{r}) \chi_\gamma(\vec{r}) \frac{e^2 e^{-\lambda|\vec{r}-\vec{r}'|}}{4\pi|\vec{r}-\vec{r}'|} \chi_\beta^*(\vec{r}') \chi_\delta(\vec{r}')}_{U_{\alpha,\beta,\gamma,\delta}} \\
 &= -\frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} \underbrace{\sum_n \langle \pi_\delta | \psi_n \rangle f_n \langle \psi_n | \pi_\alpha \rangle}_{\rho_{\delta,\alpha}} \underbrace{\sum_m \langle \pi_\gamma | \psi_m \rangle f_m \langle \psi_m | \pi_\beta \rangle}_{\rho_{\gamma,\beta}} U_{\alpha,\beta,\gamma,\delta} \\
 &= -\frac{1}{2} \sum_{\alpha,\beta,\gamma,\delta} \rho_{\delta,\alpha} \rho_{\gamma,\beta} U_{\alpha,\beta,\gamma,\delta}
 \end{aligned} \tag{7.1}$$

with the density matrix

$$\rho_{\alpha,\beta} = \sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle \psi_n | \pi_\beta \rangle \tag{7.2}$$

and the screened U-tensor

$$U_{\alpha,\beta,\gamma,\delta} = \int d^3 r \int d^3 r' \chi_\alpha^*(\vec{r}) \chi_\beta^*(\vec{r}') \frac{e^2 \cdot e^{-\lambda|\vec{r}'-\vec{r}|}}{4\pi\epsilon_0|\vec{r}'-\vec{r}|} \chi_\gamma(\vec{r}) \chi_\delta(\vec{r}') \tag{7.3}$$

7.1.1 Division of exchange energy

The goal is to limit the set of matrix elements for which the Coulomb interaction is considered to onsite terms and terms which are restricted to a pair of atoms.

The exchange energy can be divided into terms for which the density connected to one node of the interaction, i.e. either \vec{r} or \vec{r}' , are centered on the same atom or on different atoms.

If the two orbitals are centered on two atoms, the density is located in the bond center and its contribution vanishes quickly with distance.

When we exclude terms for which one node is connected to orbitals from different sites there are only in-site and NDDO terms left. NDDO stands for "Neglect of diatomic differential overlap" and has been invented by Pople.

$$E_{xc} = \underbrace{-\frac{1}{2} \sum_R \sum_{\alpha, \beta, \gamma, \delta \in R} \rho_{\delta, \alpha} \rho_{\gamma, \beta} U_{\alpha, \beta, \gamma, \delta}}_{\text{onsite}} - \underbrace{\frac{1}{2} \sum_{R \neq R'} \sum_{\alpha, \gamma \in R} \sum_{\delta, \beta \in R'} \rho_{\delta, \alpha} \rho_{\gamma, \beta} U_{\alpha, \beta, \gamma, \delta}}_{\text{NDDO-terms}} + \text{terms with off-site densities} \quad (7.4)$$

If we include all terms centered on a bond, there is one contribution with one off-site density, the 31-term, and one with two off-site densities, which I call bond exchange "bondx".

$$\Delta E_{xc} = \underbrace{-\frac{1}{2} \sum_{R \neq R'} 4 \sum_{\alpha \in R} \sum_{\beta, \gamma, \delta \in R'} \rho_{\delta, \alpha} \rho_{\gamma, \beta} U_{\alpha, \beta, \gamma, \delta}}_{\text{31-terms}} - \underbrace{\frac{1}{2} \sum_{R \neq R'} \sum_{\alpha, \beta \in R} \sum_{\delta, \gamma \in R'} \rho_{\delta, \alpha} \rho_{\gamma, \beta} U_{\alpha, \beta, \gamma, \delta}}_{\text{Bondx-terms}} + \text{terms on at least three sites} \quad (7.5)$$

For the time being we concentrate on the NDDO terms.

7.1.2 Interaction

Let me abstract from the specific form of the interaction and describe it as $w(r)$. Later, we will require its value and derivatives which are computed here.

$$\begin{aligned}
w(r) &= \frac{e^2}{4\pi\epsilon_0 r} \underbrace{e^{-\lambda r}}_{\text{screening}} \\
\partial_r w(r) &= \frac{-e^2}{4\pi\epsilon_0 r^2} \underbrace{(1 + \lambda r)}_{\text{screening}} e^{-\lambda r} \\
\partial_r^2 w(r) &= \frac{-e^2}{4\pi\epsilon_0 r^2} (1 + \lambda r) e^{-\lambda r} \underbrace{\left(-\frac{2}{r} - \lambda + \frac{\lambda}{1 + \lambda r}\right)}_{-\lambda^2 r / (1 + \lambda r)} = \frac{+2e^2}{4\pi\epsilon_0 r^3} \underbrace{(1 + \lambda r) \left(1 + \frac{1}{2} \frac{\lambda^2 r^2}{1 + \lambda r}\right)}_{\text{screening}} e^{-\lambda r} \\
&= \frac{+2e^2}{4\pi\epsilon_0 r^3} \underbrace{\left(1 + \lambda r + \frac{1}{2} \lambda^2 r^2\right)}_{\text{screening}} e^{-\lambda r} \tag{7.6}
\end{aligned}$$

7.2 Long-range expansions

When the Coulomb potential is placed on a radial grid, it is important to remove the long-ranged tail, which extends beyond the grid. This is important, in particular if the potential is interpolated.

7.2.1 Long-range expansion of the potential

Let me start with the potential of an electron density

$$\begin{aligned}
v(\vec{r}) &= \int d^3 r' w(|\vec{r} - \vec{r}'|) n(\vec{r}') \\
&= \int d^3 r' n(\vec{r}') \left[w(|\vec{r} - \vec{R}'|) + (\vec{r} - \vec{R}') \cdot \vec{\nabla}_{r'} w(|\vec{r} - \vec{r}'|) \Big|_{\vec{r}' = \vec{R}'} + \dots \right] \\
&= \int d^3 r' n(\vec{r}') \left[w(|\vec{r} - \vec{R}'|) - (\vec{r} - \vec{R}') \cdot \vec{\nabla}_{r'} w(|\vec{r} - \vec{R}'|) + \dots \right] \\
&= w(|\vec{r} - \vec{R}|) \underbrace{\left[\int d^3 r' n(\vec{r}') \right]}_Q - \left(\vec{\nabla} w(|\vec{r} - \vec{R}'|) \right) \underbrace{\left[\int d^3 r' n(\vec{r}') (\vec{r} - \vec{R}') \right]}_{\vec{D}} + \dots \\
&= \int d^3 r' \delta(\vec{r} - \vec{R}') \left[Q w(|\vec{r} - \vec{r}'|) - \vec{D} \cdot \vec{\nabla} w(|\vec{r} - \vec{r}'|) + \dots \right] \\
&= \int d^3 r' w(|\vec{r} - \vec{r}'|) \left[Q \delta(\vec{r} - \vec{R}') - \vec{D} \cdot \vec{\nabla} \delta(\vec{r} - \vec{R}') + \dots \right] \tag{7.7}
\end{aligned}$$

This provides me with the multipole expansion of the potential, as well as a multipole representation of the density.

$$P(\vec{r}) = \sum_R Q \delta(\vec{r} - \vec{R}) - \vec{D} \cdot \vec{\nabla} \delta(\vec{r} - \vec{R}) + \dots \tag{7.8}$$

The multipole representation of the density is a density that vanishes everywhere else than a point, but produces the same multipole expansion of the potential as the original density.

The long-ranged potential $v^{lr}(\vec{r})$ is the potential created by the multipole expansion.

$$v^{lr}(\vec{r}) = \int d^3r' w(|\vec{r} - \vec{r}'|) P(\vec{r}') = (Q - \vec{D} \cdot \vec{\nabla}) w(|\vec{r} - \vec{r}'|) \quad (7.9)$$

It can be expressed in terms of real spherical harmonics as

$$v^{lr}(\vec{r}) = \left[\sqrt{4\pi} Q w(|\vec{r} - \vec{r}'|) \right] Y_s(\vec{r} - \vec{r}') + \sum_i \left[\sqrt{\frac{4\pi}{3}} D_i \left(\partial_r w|_{|\vec{r} - \vec{r}'|} \right) \right] Y_{p_i}(\vec{r} - \vec{r}') \quad (7.10)$$

7.2.2 Long-range expansion of the energy

$$\begin{aligned} E^{lr} &= \int d^3r \int d^3r' P_1(\vec{r}) w(|\vec{r} - \vec{r}'|) P_2(\vec{r}') \\ &= \int d^3r \left[Q_1 \delta(\vec{r} - \vec{R}_1) - \vec{D}_1 \cdot \vec{\nabla}_r \delta(\vec{r} - \vec{R}_1) \right] \\ &\quad \times \int d^3r' w(|\vec{r} - \vec{r}'|) \left[Q_2 \delta(\vec{r}' - \vec{R}_2) - \vec{D}_2 \cdot \vec{\nabla}_{r'} \delta(\vec{r}' - \vec{R}_2) \right] \\ &= \int d^3r \left[Q_1 \delta(\vec{r} - \vec{R}_1) + \delta(\vec{r} - \vec{R}_1) \vec{D}_1 \cdot \vec{\nabla}_r \right] \left[Q_2 w(|\vec{r} - \vec{R}_2|) - \vec{D}_2 \cdot \vec{\nabla}_r w(|\vec{r} - \vec{R}_2|) \right] \\ &= Q_1 Q_2 w(|\vec{R}_1 - \vec{R}_2|) + \vec{D}_1 Q_2 \vec{\nabla}_{R_1} w(|\vec{R}_1 - \vec{R}_2|) - Q_1 \vec{D}_2 \vec{\nabla}_{R_1} w(|\vec{R}_1 - \vec{R}_2|) \\ &\quad - \left(\vec{D}_1 \vec{\nabla}_{R_1} \right) \left(\vec{D}_2 \vec{\nabla}_{R_1} \right) w(|\vec{R}_1 - \vec{R}_2|) \\ &= Q_1 Q_2 w(|\vec{R}_1 - \vec{R}_2|) + \left(\vec{D}_1 Q_2 - Q_1 \vec{D}_2 \right) \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{R}_1 - \vec{R}_2}{|\vec{R}_1 - \vec{R}_2|} \\ &\quad - \left(\vec{D}_1 \vec{\nabla}_{R_1} \right) \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \\ &= Q_1 Q_2 w(|\vec{R}_1 - \vec{R}_2|) + \left(\vec{D}_1 Q_2 - Q_1 \vec{D}_2 \right) \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{R}_1 - \vec{R}_2}{|\vec{R}_1 - \vec{R}_2|} \\ &\quad - \partial_r^2 w \Big|_{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_1 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} - \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \left[\left(\vec{D}_1 \vec{\nabla}_{R_1} \right) \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \right] \\ &= Q_1 Q_2 w(|\vec{R}_1 - \vec{R}_2|) + \left(\vec{D}_1 Q_2 - Q_1 \vec{D}_2 \right) \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{R}_1 - \vec{R}_2}{|\vec{R}_1 - \vec{R}_2|} \\ &\quad - \partial_r^2 w \Big|_{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_1 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \\ &\quad - \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \left[\frac{\vec{D}_1 \vec{D}_2}{|\vec{R}_1 - \vec{R}_2|} - \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|^2} \frac{\vec{D}_1 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \right] \\ &= Q_1 Q_2 w(|\vec{R}_1 - \vec{R}_2|) + \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \left[\left(\vec{D}_1 Q_2 - Q_1 \vec{D}_2 \right) \frac{\vec{R}_1 - \vec{R}_2}{|\vec{R}_1 - \vec{R}_2|} - \frac{\vec{D}_1 \vec{D}_2}{|\vec{R}_1 - \vec{R}_2|} \right] \\ &\quad + \left[-\partial_r^2 w \Big|_{|\vec{R}_1 - \vec{R}_2|} + \frac{1}{|\vec{R}_1 - \vec{R}_2|} \partial_r w \Big|_{|\vec{R}_1 - \vec{R}_2|} \right] \frac{\vec{D}_1 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \quad (7.11) \end{aligned}$$

For the screened potential, we obtain:

$$\begin{aligned}
 E^{lr} &= \frac{e^2}{4\pi\epsilon_0} e^{-\lambda|\vec{R}_1 - \vec{R}_2|} \left\{ \frac{Q_1 Q_2}{|\vec{R}_1 - \vec{R}_2|} - \frac{1 + \lambda|\vec{R}_1 - \vec{R}_2|}{|\vec{R}_1 - \vec{R}_2|^2} \left[\left(\vec{D}_1 Q_2 - Q_1 \vec{D}_2 \right) \frac{\vec{R}_1 - \vec{R}_2}{|\vec{R}_1 - \vec{R}_2|} - \frac{\vec{D}_1 \vec{D}_2}{|\vec{R}_1 - \vec{R}_2|} \right] \right. \\
 &\quad \left. + \left[-2 \frac{1 + \lambda|\vec{R}_1 - \vec{R}_2| + \frac{1}{2} \lambda^2 |\vec{R}_1 - \vec{R}_2|^2}{|\vec{R}_1 - \vec{R}_2|^3} - \frac{1 + \lambda|\vec{R}_1 - \vec{R}_2|}{|\vec{R}_1 - \vec{R}_2|^3} \right] \frac{\vec{D}_1 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \right\} \\
 &= \frac{e^2}{4\pi\epsilon_0} e^{-\lambda|\vec{R}_1 - \vec{R}_2|} \left\{ \frac{Q_1 Q_2}{|\vec{R}_1 - \vec{R}_2|} - \frac{1 + \lambda|\vec{R}_1 - \vec{R}_2|}{|\vec{R}_1 - \vec{R}_2|^2} \left[\left(\vec{D}_1 Q_2 - Q_1 \vec{D}_2 \right) \frac{\vec{R}_1 - \vec{R}_2}{|\vec{R}_1 - \vec{R}_2|} - \frac{\vec{D}_1 \vec{D}_2}{|\vec{R}_1 - \vec{R}_2|} \right] \right. \\
 &\quad \left. - \frac{3(1 + \lambda|\vec{R}_1 - \vec{R}_2|) + \lambda^2 |\vec{R}_1 - \vec{R}_2|^2}{|\vec{R}_1 - \vec{R}_2|^3} \frac{\vec{D}_1 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \frac{\vec{D}_2 (\vec{R}_1 - \vec{R}_2)}{|\vec{R}_1 - \vec{R}_2|} \right\} \quad (7.12)
 \end{aligned}$$

With $\vec{R}_2 = \vec{R}_1 + \Delta \vec{e}_z$, we obtain

$$\begin{aligned}
 E^{lr} &= \frac{e^2}{4\pi\epsilon_0} e^{-\lambda\Delta} \left\{ \frac{Q_1 Q_2}{\Delta} + \frac{1 + \lambda\Delta}{\Delta^2} \left[\left(\vec{e}_z \vec{D}_1 Q_2 - Q_1 \vec{e}_z \vec{D}_2 \right) + \frac{\vec{D}_1 \vec{D}_2}{\Delta} \right] \right. \\
 &\quad \left. - \frac{3(1 + \lambda\Delta) + \lambda^2 \Delta^2}{\Delta^3} \left(\vec{e}_z \vec{D}_1 \right) \left(\vec{e}_z \vec{D}_2 \right) \right\} \quad (7.13)
 \end{aligned}$$

The potential of atom two and its gradient at \vec{R}_1 is obtained as

$$\begin{aligned}
 V(\vec{R}_1) &= \frac{\partial E^{lr}}{\partial Q_1} \\
 \vec{\nabla}_{R_1} V(\vec{R}_1) &= \vec{\nabla}_{D_1} E^{lr} \quad (7.14)
 \end{aligned}$$

7.2.3 Division of U-tensor into short- and long-ranged parts

In order to perform the integrations efficiently, we need to remove the long-ranged tails from the potential.

Let me make the following definitions.

$$\begin{aligned}
 n_{\alpha,\beta}(\vec{r}) &= \chi_{\alpha}^*(\vec{r}) \chi_{\beta}(\vec{r}) \\
 n_{\gamma,\delta}(\vec{r}) &= \chi_{\gamma}^*(\vec{r}) \chi_{\delta}(\vec{r}) \\
 v_{\alpha,\beta}(\vec{r}) &= \int d^3 r' w(|\vec{r} - \vec{r}'|) n_{\alpha,\beta}(\vec{r}') \\
 v_{\gamma,\delta}(\vec{r}) &= \int d^3 r' w(|\vec{r} - \vec{r}'|) n_{\gamma,\delta}(\vec{r}') \quad (7.15)
 \end{aligned}$$

Let me furthermore introduce the multipole representation of the density $n_{\alpha,\beta}(\vec{r})$ and $n_{\gamma,\delta}(\vec{r})$

$$\begin{aligned}
 \rho_{\alpha,\beta}(\vec{r}) &= Q_{\alpha,\beta} \delta(\vec{r} - \vec{R}) - \vec{D}_{\alpha,\beta} \vec{\nabla} \delta(\vec{r} - \vec{R}) \\
 \rho_{\gamma,\delta}(\vec{r}) &= Q_{\gamma,\delta} \delta(\vec{r} - \vec{R}') - \vec{D}_{\gamma,\delta} \vec{\nabla} \delta(\vec{r} - \vec{R}') \quad (7.16)
 \end{aligned}$$

The multipole representation has the same multipoles as the original density, but it is non-zero only at a point.

Then, the U-tensor matrix elements are

$$\begin{aligned}
U_{\alpha,\beta,\gamma,\delta} &= \int d^3r \int d^3r' n_{\alpha,\beta}(\vec{r}) w(|\vec{r} - \vec{r}'|) n_{\gamma,\delta}(\vec{r}') \\
&= \int d^3r' v_{\alpha,\beta}(\vec{r}') n_{\gamma,\delta}(\vec{r}') \\
&= \int d^3r' \left[v_{\alpha,\beta}(\vec{r}') - \int d^3r p_{\alpha,\beta}(\vec{r}) w(|\vec{r} - \vec{r}'|) \right] n_{\gamma,\delta}(\vec{r}') \\
&\quad + \int d^3r p_{\alpha,\beta}(\vec{r}) v_{\gamma,\delta}(\vec{r}) \\
&= \int d^3r' \left[v_{\alpha,\beta}(\vec{r}') - \int d^3r p_{\alpha,\beta}(\vec{r}) w(|\vec{r} - \vec{r}'|) \right] n_{\gamma,\delta}(\vec{r}') \\
&\quad + \int d^3r p_{\alpha,\beta}(\vec{r}) \left[v_{\gamma,\delta}(\vec{r}) - \int d^3r' w(|\vec{r} - \vec{r}'|) p_{\gamma,\delta}(\vec{r}') \right] \\
&\quad + \int d^3r \int d^3r' p_{\alpha,\beta}(\vec{r}) w(|\vec{r} - \vec{r}'|) p_{\gamma,\delta}(\vec{r}') \\
&= \int d^3r' v_{\alpha,\beta}^{sr}(\vec{r}') n_{\gamma,\delta}(\vec{r}') + \int d^3r p_{\alpha,\beta}(\vec{r}) v_{\gamma,\delta}^{sr}(\vec{r}) + U_{\alpha,\beta,\gamma,\delta}^{lr} \quad (7.17)
\end{aligned}$$

OFFSITE NDDO U-TENSOR IN TERMS OF SHORT AND LONG-RANGED CONTRIBUTIONS

The following division is exact, even if the multipole expansion of the point representation $p(\vec{r})$ differs from that of the corresponding densities $n(\vec{r})$. It is useful, when $n_{\alpha,\beta}$ and $n_{\gamma,\delta}$ have rapidly converging multipole expansions, which is well represented by the corresponding point representation.

$$U_{\alpha,\beta,\gamma,\delta} = \int d^3r' v_{\alpha,\beta}^{sr}(\vec{r}') n_{\gamma,\delta}(\vec{r}') + \int d^3r p_{\alpha,\beta}(\vec{r}) v_{\gamma,\delta}^{sr}(\vec{r}) + U_{\alpha,\beta,\gamma,\delta}^{lr} \quad (7.18)$$

with the short-ranged potentials

$$\begin{aligned}
v_{\alpha,\beta}^{sr}(\vec{r}') &\stackrel{\text{def}}{=} v_{\alpha,\beta}(\vec{r}') - \int d^3r w(|\vec{r}' - \vec{r}|) p_{\alpha,\beta}(\vec{r}) \\
v_{\gamma,\delta}^{sr}(\vec{r}) &\stackrel{\text{def}}{=} v_{\gamma,\delta}(\vec{r}) - \int d^3r' w(|\vec{r} - \vec{r}'|) p_{\gamma,\delta}(\vec{r}') \quad (7.19)
\end{aligned}$$

and the long-ranged part of the U-tensor

$$U_{\alpha,\beta,\gamma,\delta}^{lr} \stackrel{\text{def}}{=} \int d^3r \int d^3r' p_{\alpha,\beta}(\vec{r}) w(|\vec{r} - \vec{r}'|) p_{\gamma,\delta}(\vec{r}') \quad (7.20)$$

7.3 Expansion of the density in angular momenta

In `simplelmtto_onecenterqln`, we evaluate the quantities $q_{s,\alpha,\beta}$ and $q_{p,\alpha,\beta}$, which can be used to evaluate monopoles and dipoles. We divide an orbital into radial and angular part, as

$$\chi_\alpha(\vec{r}) = f_\alpha(|\vec{r} - \vec{R}_\alpha|) Y_{L_\alpha}(\vec{r} - \vec{R}_\alpha) \quad (7.21)$$

Thus, we obtain the densities of two orbitals centered at the same site $\vec{R} = \vec{R}_\alpha = \vec{R}_\beta$

$$\begin{aligned} n_{\alpha,\beta}(\vec{r}) &= \chi_\alpha^*(\vec{r}) \chi_\beta(\vec{r}) \\ &= f_\alpha(|\vec{r} - \vec{R}|) f_\beta(|\vec{r} - \vec{R}|) Y_{L_\alpha}^*(\vec{r} - \vec{R}) Y_{L_\beta}(\vec{r} - \vec{R}) \\ &= \sum_L C_{L,L_\alpha,L_\beta} f_\alpha(|\vec{r} - \vec{R}|) f_\beta(|\vec{r} - \vec{R}|) Y_L(\vec{r} - \vec{R}) \end{aligned} \quad (7.22)$$

with the Gaunt coefficients $C_{L,L',L''} = \int d\Omega Y_L(\vec{r}) Y_{L'}(\vec{r}) Y_{L''}(\vec{r})$

$$\begin{aligned} Q_{\alpha,\beta} &= \int d^3r \chi_\alpha^*(\vec{r}) \chi_\beta(\vec{r}) = \int d^3r \chi_\alpha^*(\vec{r}) \chi_\beta(\vec{r}) \underbrace{\sqrt{4\pi} |\vec{r} - \vec{R}|^0 Y_s(\vec{r} - \vec{R})}_{=1} \\ &= C_{s,L_\alpha,L_\beta} \underbrace{\sqrt{4\pi} \int_0^\infty dr r^2 f_\alpha(r) f_\beta(r)}_{q_{s,\alpha,\beta}} \end{aligned} \quad (7.23)$$

$$\begin{aligned} D_{j,\alpha,\beta} &= \int d^3r \chi_\alpha^*(\vec{r}) \chi_\beta(\vec{r}) (r_j - R_j) = \int d^3r \chi_\alpha^*(\vec{r}) \chi_\beta(\vec{r}) |\vec{r} - \vec{R}|^1 \underbrace{\sqrt{\frac{4\pi}{3}} Y_{p_j}(\vec{r} - \vec{R})}_{r_j} \\ &= C_{p_j,L_\alpha,L_\beta} \underbrace{\sqrt{\frac{4\pi}{3}} \int_0^\infty dr r^3 f_\alpha(r) f_\beta(r)}_{q_{p,\alpha,\beta}} \end{aligned} \quad (7.24)$$

The variables $q_{s,\alpha,\beta}$ and $q_{p,\alpha,\beta}$ are stored as `QLN(1, LN1, LN2)` and `QLN(2, LN1, LN2)`, respectively.

Short-ranged potentials

$$\begin{aligned}
v_{\gamma,\delta}^{sr}(\vec{r}) &= \sum_L v_{L,\gamma,\delta}^{sr}(|\vec{r} - \vec{R}_2|) Y_L(\vec{r} - \vec{R}_2) \\
&= \sum_L C_{L,\gamma,L,\delta,L} \int d^3r' w(|\vec{r} - \vec{r}'|) f_\gamma(|\vec{r}' - \vec{R}_2|) f_\gamma(|\vec{r}' - \vec{R}_2|) Y_L(\vec{r}' - \vec{R}_2) \\
&\quad - \underbrace{C_{S,L,\gamma,L,\delta} q_{S,\gamma,\delta}}_{Q_{\gamma,\delta}} \sqrt{4\pi} w(|\vec{r} - \vec{R}_2|) Y_S(\vec{r} - \vec{R}_2) \\
&\quad - \sum_j \underbrace{C_{p_j,L,\gamma,L,\delta} q_{p,\gamma,\delta}}_{D_{j,\gamma,\delta}} \sqrt{\frac{4\pi}{3}} \left(\partial_r w|_{|\vec{r}-\vec{R}_2|} \right) Y_{p_j}(\vec{r} - \vec{R}_2) \\
&= \sum_L C_{L,\gamma,L,\delta,L} \left\{ \int d^3r' w_\ell(|\vec{r} - \vec{r}'|) f_\gamma(|\vec{r}' - \vec{R}_2|) f_\gamma(|\vec{r}' - \vec{R}_2|) \right. \\
&\quad \left. - \left[\delta_{L,S} q_{S,\gamma,\delta} \sqrt{4\pi} w(|\vec{r} - \vec{R}_2|) + \sum_j \delta_{L,p_j} q_{p,\gamma,\delta} \sqrt{\frac{4\pi}{3}} \left(\partial_r w|_{|\vec{r}-\vec{R}_2|} \right) \right] \right\} Y_L(\vec{r}' - \vec{R}_2) \\
&= \sum_L C_{L,\gamma,L,\delta,L} \left\{ \int d^3r' w_\ell(|\vec{r} - \vec{r}'|) f_\gamma(|\vec{r}' - \vec{R}_2|) f_\gamma(|\vec{r}' - \vec{R}_2|) \right. \\
&\quad \left. - \left[\delta_{\ell,0} q_{S,\gamma,\delta} \sqrt{4\pi} w(|\vec{r} - \vec{R}_2|) + \delta_{\ell,1} q_{p,\gamma,\delta} \sqrt{\frac{4\pi}{3}} \left(\partial_r w|_{|\vec{r}-\vec{R}_2|} \right) \right] \right\} Y_L(\vec{r}' - \vec{R}_2) \quad (7.25)
\end{aligned}$$

This expression is used to construct the short-ranged potentials.

Correction

The middle term in Eq. 7.18, which I denote as X , is written as

$$\begin{aligned}
X &\stackrel{\text{def}}{=} \int d^3r \, p_{\alpha,\beta}(\vec{r}) v_{\gamma,\delta}^{sr}(\vec{r}) \\
&= \int d^3r \left[Q_{\alpha,\beta} \delta(\vec{r} - \vec{R}_1) - \vec{D}_{\alpha,\beta} \vec{\nabla} \delta(\vec{r} - \vec{R}_1) \right] v_{\gamma,\delta}^{sr}(\vec{r}) \\
&= \int d^3r \, \delta(\vec{r} - \vec{R}_1) \left[Q_{\alpha,\beta} + \vec{D}_{\alpha,\beta} \vec{\nabla} \right] v_{\gamma,\delta}^{sr}(\vec{r}) \\
&= \left[Q_{\alpha,\beta} + \vec{D}_{\alpha,\beta} \vec{\nabla}_{R_1} \right] v_{\gamma,\delta}^{sr}(\vec{R}_1) \\
&= \underbrace{C_{L\alpha,L\beta,s} q_{s,\alpha,\beta}}_{Q_{\alpha,\beta}} + \sum_{j \in \{x,y,z\}} \underbrace{C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta}}_{D_{j,\alpha,\beta}} \frac{\partial}{\partial R_{1j}} \Bigg] \\
&\quad \times \sum_L v_{L,\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) Y_L(\vec{R}_1 - \vec{R}_2) \\
&= \sum_L C_{L\alpha,L\beta,s} q_{s,\alpha,\beta} v_{L,\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) Y_L(\vec{R}_1 - \vec{R}_2) \\
&\quad + \sum_L \sum_{j \in \{x,y,z\}} C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta} \left(\partial_r v_{L,\gamma,\delta}^{sr} \Big|_{|\vec{R}_1 - \vec{R}_2|} \right) \frac{R_{1j} - R_{2j}}{|\vec{R}_1 - \vec{R}_2|} Y_L(\vec{R}_1 - \vec{R}_2) \\
&\quad + \sum_L \sum_{j \in \{x,y,z\}} C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta} v_{L,\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) \\
&\quad \times \underbrace{\sqrt{\frac{4\pi}{3}} \sum_{L'} C_{j,L,L'} \frac{1}{|\vec{R}_1 - \vec{R}_2|} [-\ell + (2\ell + 1)\delta_{\ell',\ell-1}] Y_{L'}(\vec{R}_1 - \vec{R}_2)}_{\frac{\partial}{\partial R_{1j}} Y_L(\vec{R}_1 - \vec{R}_2)} \tag{7.26}
\end{aligned}$$

We used the expression for the gradient of spherical harmonics (Section 49.10 “Gradients of spherical harmonics” of Methods)

$$\vec{\nabla}_j Y_L(\vec{r}) = \sqrt{\frac{4\pi}{3}} \sum_{L'} C_{j,L,L'} \frac{1}{|\vec{r}|} [-\ell + (2\ell + 1)\delta_{\ell',\ell-1}] Y_{L'}(\vec{r}) \tag{7.27}$$

In the following step I use $\vec{R}_1 - \vec{R}_2 = -\vec{e}_z|\vec{R}_1 - \vec{R}_2|$.

$$\begin{aligned}
X &= \sum_L C_{L\alpha,L\beta,s} q_{s,\alpha,\beta} v_{L,\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) Y_L(\vec{R}_1 - \vec{R}_2) \\
&+ \sum_L \sum_{j \in \{x,y,z\}} C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta} \left(\partial_r v_{L,\gamma,\delta}^{sr} \Big|_{|\vec{R}_1 - \vec{R}_2|} \right) \underbrace{\frac{R_{1,j} - R_{2,j}}{|\vec{R}_1 - \vec{R}_2|}}_{-\delta_{j,z}} \underbrace{Y_L(\vec{R}_1 - \vec{R}_2)}_{(-1)^\ell Y_L(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) \delta_{m,0}} \\
&+ \sum_L \sum_{j \in \{x,y,z\}} C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta} v_{L,\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) \\
&\times \sqrt{\frac{4\pi}{3}} \sum_{L'} C_{j,L,L'} \frac{1}{|\vec{R}_1 - \vec{R}_2|} [-\ell + (2\ell + 1)\delta_{\ell',\ell-1}] \underbrace{Y_{L'}(\vec{R}_1 - \vec{R}_2)}_{\delta_{m',0}(-1)^{\ell'} Y_{L'}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|)} \\
&= \sum_\ell C_{L\alpha,L\beta,s} q_{s,\alpha,\beta} (-1)^\ell Y_{\ell,0}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) v_{(\ell,0),\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) \\
&+ \sum_\ell C_{L\alpha,L\beta,p_z} q_{p,\alpha,\beta} \left(\partial_r v_{(\ell,0),\gamma,\delta}^{sr} \Big|_{|\vec{R}_1 - \vec{R}_2|} \right) (-1)^{\ell+1} Y_{(\ell,0)}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) \\
&+ \sum_L \sum_{j \in \{x,y,z\}} C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta} \frac{v_{L,\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|)}{|\vec{R}_1 - \vec{R}_2|} \\
&\times \sqrt{\frac{4\pi}{3}} \sum_{\ell'} C_{p_j,L,(\ell',0)} [-\ell + (2\ell + 1)\delta_{\ell',\ell-1}] (-1)^{\ell'} Y_{(\ell',0)}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|)
\end{aligned} \tag{7.28}$$

Now we exploit that $C_{p_j,L,(\ell',0)}$ vanishes except when $m_{p_j} = m_L$ and $\ell_L - \ell' = \pm 1$.

$$\begin{aligned}
X &= \sum_\ell C_{L\alpha,L\beta,s} q_{s,\alpha,\beta} v_{(\ell,0),\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|) (-1)^\ell Y_{\ell,0}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) \\
&+ \sum_\ell C_{L\alpha,L\beta,p_z} q_{p,\alpha,\beta} \left(\partial_r v_{(\ell,0),\gamma,\delta}^{sr} \Big|_{|\vec{R}_1 - \vec{R}_2|} \right) (-1)^{\ell+1} Y_{(\ell,0)}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) \\
&+ \sum_\ell \sum_{j \in \{x,y,z\}} C_{L\alpha,L\beta,p_j} q_{p,\alpha,\beta} \frac{v_{(\ell,m_j),\gamma,\delta}^{sr}(|\vec{R}_1 - \vec{R}_2|)}{|\vec{R}_1 - \vec{R}_2|} \sqrt{\frac{4\pi}{3}} \\
&\times \left\{ C_{p_j,(\ell,m_j),(\ell-1,0)} [\ell + 1] (-1)^{\ell-1} Y_{(\ell-1,0)}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) \right. \\
&\quad \left. + C_{p_j,(\ell,m_j),(\ell+1,0)} [-\ell] (-1)^{\ell+1} Y_{(\ell+1,0)}(+\vec{e}_z|\vec{R}_1 - \vec{R}_2|) \right\}
\end{aligned} \tag{7.29}$$

7.3.1 Tabulating data

$$\begin{aligned}
 U_{\alpha,\beta,\gamma,\delta} &= \int d^3r \int d^3r' n_{\alpha,\gamma}(\vec{r}) w(\vec{r} - \vec{r}') n_{\beta,\delta}(\vec{r}) \\
 &= \int d^3r n_{\alpha,\gamma}(\vec{r}) v(\vec{r}) \\
 &= \sum_{L,L'} \sum_{L_\alpha,L_\gamma} \sum_{L_\beta,L_\delta} \delta_{m,m'} C_{L,L_\alpha,L_\gamma} C_{L',L_\beta,L_\delta} \int dr f_\alpha(\vec{r} - \vec{R}) f_\gamma(\vec{r} - \vec{R}) Y_L(\vec{r} - \vec{R}) \\
 &\quad \times \underbrace{\int dr' w(\vec{r} - \vec{r}') f_\beta(\vec{r}' - \vec{R}') f_\delta(\vec{r}' - \vec{R}') Y_{L'}(\vec{r}' - \vec{R}')}_{v_{\ell',\beta,\delta}(|\vec{r} - \vec{R}'|) Y_{L'}(\vec{r}' - \vec{R}')} \quad (7.30)
 \end{aligned}$$

The integral is performed in `simplelmtwo_center`.

$$\begin{aligned}
 I &= \int_{-\infty}^{\infty} dz \int_0^{2\pi} d\phi \int_0^{\infty} d\rho F(|\vec{r}|) Y_{\ell,m}(\vec{r}) G(|\vec{r} - d\vec{e}_z|) Y_{\ell',m'}(\vec{r} - d\vec{e}_z) \\
 &= 2\pi \delta_{m,m'} \sqrt{\frac{2\ell+1}{4\pi} \frac{(\ell-m)!}{(\ell+m)!}} \sqrt{\frac{2\ell'+1}{4\pi} \frac{(\ell'-m)!}{(\ell'+m)!}} \\
 &\quad \times \int_{-\infty}^{\infty} dz \int_0^{\infty} d\rho P_\ell^m\left(\frac{z}{\sqrt{\rho^2 + z^2}}\right) P_{\ell'}^m\left(\frac{z-d}{\sqrt{\rho^2 + (z-d)^2}}\right) F(|\vec{r}|) G(|\vec{r} - d\vec{e}_z|) \quad (7.31)
 \end{aligned}$$

The z-axis is chosen along the interatomic distance vector. Thus the integral gives a non-zero result if the magnetic quantum numbers have the same value. ρ is the distance from the bond axis.

The Legendre polynomials are, with $m > 0$,

$$\begin{aligned}
 P_\ell^m(z) &= \frac{(-1)^m}{\ell! 2^\ell} (\sqrt{1-x^2})^m \left(\frac{d}{dx}\right)^{\ell+m} (1-x^2)^\ell \\
 P_\ell^{-m}(z) &= (-1)^m \frac{(\ell-m)!}{(\ell+m)!} P_\ell^m(x) \quad (7.32)
 \end{aligned}$$

The method is taken from Genz and Malik[?], who use a variant of the method of van Dooren and de Ridder[?].

For comparison, see also the work of Romanowski and Jalbout. [? ? ? ?]

The radial functions of the local orbitals do not depend on the magnetic quantum number. Hence, the integral depends on $(\ell, \ell', m, \ell_\alpha, n_\alpha, \ell_\gamma, n_\gamma, \ell_\beta, n_\beta, \ell_\delta, n_\delta)$ with the same m . These integrals are calculated on a distance grid and then fitted by a set of interpolating exponentials.

The U-tensor is

$$\begin{aligned}
 U_{\alpha,\beta,\gamma,\delta} &= \sum_{\ell,\ell'} \sum_m \sum_{\ell_\alpha,\ell_\gamma} \sum_{\ell_\beta,\ell_\delta} \left(\sum_{m_\alpha,m_\gamma} \sum_{m_\beta,m_\delta} C_{\ell,m,L_\alpha,L_\gamma} C_{\ell',m,L_\beta,L_\delta} \right) \\
 &\quad \times I_{\ell,\ell',m,\ell_\alpha,n_\alpha,\ell_\gamma,n_\gamma,\ell_\beta,n_\beta,\ell_\delta,n_\delta} \left([n_{\ell,\alpha,\gamma}], [v_{\ell',\beta,\delta}], d \right) \quad (7.33)
 \end{aligned}$$

7.3.2 Integrals with the point density

Let me consider the two-center integral Eq. 7.31, with the density $n_{\alpha,\beta}$ replaced by its point-density representation.

$$\begin{aligned} n_{\alpha,\beta}(\vec{R} + \vec{r}) &= \sum_L C_{L,L_\alpha,L_\beta} f_\alpha(|\vec{r}|) f_\beta(|\vec{r}|) Y_L(\vec{r}) \\ \rightarrow p_{\alpha,\beta}(\vec{R} + \vec{r}) &= C_{s,L_\alpha,L_\beta} q_{s,\alpha,\beta} \delta(\vec{r}) - \sum_j C_{p_j,L_\alpha,L_\beta} q_{p,\alpha,\beta} \partial_{r_j} \delta(\vec{r}) \end{aligned} \quad (7.34)$$

This implies that we need to replace

$$\begin{aligned} n_{\ell,\alpha,\beta}(\vec{R} + \vec{r}) &= f_\alpha(|\vec{r}|) f_\beta(|\vec{r}|) \\ \rightarrow p_{(\ell,m),\alpha,\beta}(\vec{R} + \vec{r}) &= \delta_{(\ell,m),s} q_{s,\alpha,\beta} \delta(\vec{r}) + \sum_j \delta_{(\ell,m),p_j} q_{p,\alpha,\beta} \partial_{r_j} \delta(\vec{r}) \\ &= \delta(\vec{r}) \left[\delta_{(\ell,m),s} q_{s,\alpha,\beta} + \sum_j \delta_{(\ell,m),p_j} q_{p,\alpha,\beta} \partial_{r_j} \right] \end{aligned} \quad (7.35)$$

The resulting integral is

$$\begin{aligned}
& I_{\ell, \ell', m, \ell, \alpha, n_\alpha, \ell, \gamma, n_\gamma, \ell, \beta, n_\beta, \ell, \delta, n_\delta} \\
\text{Eq. 7.31} \quad & \int d^3r \left(\delta_{(\ell, m), s} q_{s, \alpha, \beta} \delta(\vec{r}) - \sum_j \delta_{(\ell, m), p_j} q_{p, \alpha, \beta} \vec{\nabla} \delta(\vec{r}) \right) v_{(\ell', m), \gamma, \delta}(\vec{r} - (\vec{R}' - \vec{R})) Y_{\ell', m}(\vec{r} - (\vec{R}' - \vec{R})) \\
& = \delta_{(\ell, m), s} q_{s, \alpha, \beta} \left(v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') \right) Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \delta_{(\ell, m), p_z} q_{p, \alpha, \beta} \left(\partial_r v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') \right) Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \sum_{p_j \in p_x, p_y, p_z} \delta_{(\ell, m), p_j} q_{p, \alpha, \beta} \left(v_{(\ell', m), \gamma, \delta}(\vec{R} - \vec{R}') \right) \\
& \quad \times \underbrace{\sqrt{\frac{4\pi}{3}} \frac{1}{|\vec{R} - \vec{R}'|} \sum_{L''} C_{p_j, (\ell', m), L''} \left(-\ell' + (2\ell' + 1) \delta_{L'', \ell' - 1} \right) Y_{L''}(\vec{R} - \vec{R}')}_{\vec{\nabla}_j Y_{\ell', m}(\vec{R} - \vec{R}'); \text{ see Eq. 7.37}} \\
& = \delta_{(\ell, m), s} \frac{q_{s, \alpha, \beta}}{\sqrt{4\pi}} v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \delta_{(\ell, m), p_z} q_{p, \alpha, \beta} \left(\partial_r v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') \right) Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \sum_{p_j \in p_x, p_y, p_z} \delta_{(\ell, m), p_j} \sqrt{\frac{4\pi}{3}} q_{p, \alpha, \beta} \frac{v_{(\ell', m), \gamma, \delta}(\vec{R} - \vec{R}')}{|\vec{R} - \vec{R}'|} \\
& \quad \times \underbrace{\sum_{\ell''} C_{p_j, (\ell', m), (\ell'', 0)} \left(-\ell' + (2\ell' + 1) \delta_{\ell'', \ell' - 1} \right) Y_{\ell'', 0}(\vec{R} - \vec{R}')}_{-\ell' C_{p_j, (\ell', m), (\ell' + 1, 0)} Y_{\ell' + 1, 0}(\vec{R} - \vec{R}') + C_{p_j, (\ell', m), (\ell' - 1, 0)} (\ell' + 1) Y_{\ell' - 1, 0}(\vec{R} - \vec{R}')} \\
& = \delta_{(\ell, m), s} \frac{q_{s, \alpha, \beta}}{\sqrt{4\pi}} v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \delta_{(\ell, m), p_z} q_{p, \alpha, \beta} \left(\partial_r v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') \right) Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \sum_{m'} \delta_{(\ell, m), (1, m')} \sqrt{\frac{4\pi}{3}} q_{p, \alpha, \beta} \frac{v_{(\ell', m), \gamma, \delta}(\vec{R} - \vec{R}')}{|\vec{R} - \vec{R}'|} \\
& \quad \times \left[-\ell' C_{(1, m'), (\ell', m), (\ell' + 1, 0)} Y_{\ell' + 1, 0}(\vec{R} - \vec{R}') + C_{(1, m'), (\ell', m), (\ell' - 1, 0)} (\ell' + 1) Y_{\ell' - 1, 0}(\vec{R} - \vec{R}') \right] \\
& = \delta_{(\ell, m), s} \frac{q_{s, \alpha, \beta}}{\sqrt{4\pi}} v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \delta_{(\ell, m), p_z} q_{p, \alpha, \beta} \left(\partial_r v_{(\ell', 0), \gamma, \delta}(\vec{R} - \vec{R}') \right) Y_{\ell', 0}(\vec{R} - \vec{R}') \\
& + \delta_{\ell, 1} \sqrt{\frac{4\pi}{3}} q_{p, \alpha, \beta} \frac{v_{(\ell', m), \gamma, \delta}(\vec{R} - \vec{R}')}{|\vec{R} - \vec{R}'|} \\
& \quad \times \left[-\ell' C_{(1, m), (\ell', m), (\ell' + 1, 0)} Y_{\ell' + 1, 0}(\vec{R} - \vec{R}') + C_{(1, m), (\ell', m), (\ell' - 1, 0)} (\ell' + 1) Y_{\ell' - 1, 0}(\vec{R} - \vec{R}') \right]
\end{aligned} \tag{7.36}$$

(Section 49.10 “Gradients of spherical harmonics” of Methods)

$$\vec{\nabla}_j Y_L(\vec{r}) = \sqrt{\frac{4\pi}{3}} \sum_{L'} C_{j,L,L'} \frac{1}{|\vec{r}|} [-\ell + (2\ell + 1)\delta_{\ell',\ell-1}] Y_{L'}(\vec{r}) \quad (7.37)$$

7.3.3 Tests

- calculate U^{lr} as function of distance, once with the routine MONOANDDIPOLE and once with the code in SIMPLELMT0_OFFSITEX22SETUP, which is embedded in the routine.
- check that long range potentials and numerical potentials agree for long distances. Then, replace the potentials by the long range part.
- The term X (“correction”) gives the same result as U^{lr} .
- the two-center term agrees with U^{lr} in a certain distance range.

During the tests all contributions to the density and the potential with $\ell, \ell' > 0$ have been set to zero.

7.3.4 Complete long-range expansion

We use

$$\begin{aligned} Q_{\alpha,\beta} &\stackrel{\text{Eq. 7.23}}{=} C_{s,L\alpha,L\beta} q_{s,\alpha,\beta} \\ D_{j,\alpha,\beta} &\stackrel{\text{Eq. 7.24}}{=} C_{p_j,L\alpha,L\beta} q_{p,\alpha,\beta} \\ n_{\alpha,\beta} &= \sum_L n_{L,\alpha,\beta}(|\vec{r} - \vec{R}|) Y_L(\vec{r} - \vec{R}) \\ n_{L,\alpha,\beta}(\vec{r}) &= f_\alpha(|\vec{r} - \vec{R}|) f_\beta(|\vec{r} - \vec{R}|) C_{L,L\alpha,L\beta} \\ n_{\gamma,\delta} &= \sum_L n_{L,\alpha,\beta}(|\vec{r} - \vec{R}|) Y_L(\vec{r} - \vec{R}) \\ n_{L,\alpha,\beta}(\vec{r}) &= f_\alpha(|\vec{r} - \vec{R}|) f_\beta(|\vec{r} - \vec{R}|) C_{L,L\alpha,L\beta} \end{aligned} \quad (7.38)$$

$$\begin{aligned} v_{\gamma,\delta}^{sr}(\vec{r}) &= \int d^3r' w(\vec{r} - \vec{r}') \left[\chi_\gamma(\vec{r}') \chi_\delta(\vec{r}') - Q_{\gamma,\delta} \delta(\vec{r}' - \vec{R}') + D_{\gamma,\delta} \vec{\nabla}' \delta(\vec{r}' - \vec{R}') \right] \\ &= \int d^3r' w(\vec{r} - \vec{r}') f_\gamma(|\vec{r}' - \vec{R}'|) f_\delta(|\vec{r}' - \vec{R}'|) \sum_L C_{L\gamma,L\delta,L} Y_L(\vec{r}' - \vec{R}') \\ &\quad - w(\vec{r} - \vec{R}') C_{s,L\gamma,L\delta} q_{s,\gamma,\delta} - \sum_{j=1}^3 C_{p_j,L\gamma,L\delta,L} q_{p,\gamma,\delta} \vec{\nabla}_j w(\vec{r} - \vec{R}') \\ &= \sum_L C_{L\gamma,L\delta,L} Y_L(\vec{r} - \vec{R}') \left[v_{L,\gamma,\delta}(|\vec{r} - \vec{R}'|) \right. \\ &\quad \left. - q_{s,\gamma,\delta} \sqrt{4\pi} w(|\vec{r} - \vec{R}'|) \delta_{L,s} - q_{p,\gamma,\delta} \sqrt{\frac{4\pi}{3}} \partial_r w|_{|\vec{r}-\vec{R}'|} \delta_{L,p_j} \right] \end{aligned} \quad (7.39)$$

$$\begin{aligned}
\int d^3 r' v_{\alpha,\beta}^{sr}(\vec{r}') n_{\gamma,\delta}(\vec{r}') &= \sum_{L,L'} \int d^3 r' Y_L(\vec{r}' - \vec{R}) Y_{L'}(\vec{r}' - \vec{R}') v_{L,\alpha,\beta}^{sr}(\vec{r}' - \vec{R}) n_{L',\gamma,\delta}(\vec{r}' - \vec{R}') \\
&= \sum_{L,L'} C_{L,L\alpha,L\beta} C_{L',L\gamma,L\delta} \int d^3 r' Y_L(\vec{r}' - \vec{R}) Y_{L'}(\vec{r}' - \vec{R}') v_{L,\alpha,\beta}^{sr}(\vec{r}' - \vec{R}) n_{L',\gamma,\delta}(\vec{r}' - \vec{R}')
\end{aligned} \tag{7.40}$$

To complete the short-ranged part of the U-tensor matrix element, we need to evaluate

$$\begin{aligned}
(Q_{\alpha,\beta} + \vec{D}_{\alpha,\beta} \vec{\nabla}_R) v_{\gamma,\delta}^{sr}(\vec{R}) &= (Q_{\alpha,\beta} + \vec{D}_{\alpha,\beta} \vec{\nabla}_R) \sum_L C_{L\gamma,L\delta,L} Y_L(\vec{R} - \vec{R}') v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}') \\
&= \sum_L C_{L\gamma,L\delta,L} \underbrace{Y_L(\vec{R} - \vec{R}')}_{\delta_{m,0} Y_{\ell,0}(-\vec{e}_z)} Q_{\alpha,\beta} v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}') \\
&\quad + \sum_L C_{L\gamma,L\delta,L} v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}') \vec{D}_{\alpha,\beta} \vec{\nabla}_R Y_L(\vec{R} - \vec{R}') \\
&\quad + \sum_L C_{L\gamma,L\delta,L} Y_L(\vec{R} - \vec{R}') \vec{D}_{\alpha,\beta} \vec{\nabla}_R v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}') \\
&= \sum_L C_{L\gamma,L\delta,L} \underbrace{Y_L(\vec{R} - \vec{R}')}_{\delta_{m,0} Y_{\ell,0}(-\vec{e}_z)} Q_{\alpha,\beta} v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}') \\
&\quad + \sum_L C_{L\gamma,L\delta,L} v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}') \vec{D}_{\alpha,\beta} \vec{\nabla}_R Y_L(\vec{R} - \vec{R}') \\
&\quad + \sum_L C_{L\gamma,L\delta,L} \underbrace{Y_L(\vec{R} - \vec{R}')}_{\delta_{m,0} Y_{\ell,0}(-\vec{e}_z)} D_{z,\alpha,\beta} \partial_r v_{L,\gamma,\delta}^{sr}(\vec{R} - \vec{R}')
\end{aligned} \tag{7.41}$$

Appendix A

Definition of solid Hankel functions

The solid Hankel function has the form

$$H_L(\vec{R}) = Y_L(\vec{R}) \begin{cases} n_\ell(\sqrt{k^2} \cdot |\vec{R}|) \cdot \sqrt{k^2}^{\ell+1} & \text{for } k^2 > 0 \text{ (Abramowitz 10.1.26)} \\ m_\ell(\sqrt{-k^2} \cdot |\vec{R}|) \cdot \sqrt{\frac{2}{\pi}} \sqrt{-k^2}^{\ell+1} & \text{for } k^2 < 0 \text{ (Abramowitz 10.2.4)} \\ (2\ell - 1)!! |\vec{R}|^{-\ell-1} & \text{for } k^2 = 0 \text{ (Abramowitz 10.2.5)} \end{cases} \quad (\text{A.1})$$

The solid Hankel function is defined such that the boundary conditions at the origin are independent of k^2 .

- the function

$$n_\ell(r) = r^\ell \left(-\frac{1}{r} \partial_r \right)^\ell \frac{1}{r} \cos(r) \quad (\text{A.2})$$

is the spherical Neumann function (see Eq. 8.175 of Cohen Tannoudhi Band 2), which is also called the spherical Bessel function of the second kind. Abramowitz defines $n_\ell(r) = -y_\ell(r)$ (compare Abramowitz Eq. 10.1.26)

The spherical Neumann function obeys the radial Helmholtz equation (Abramowitz Eq. 10.1.1) for positive kinetic energy

$$\begin{aligned} r^2 \partial_r^2 n_\ell + 2r \partial_r n_\ell + (r^2 - \ell(\ell + 1)) n_\ell &= 0 \\ \Rightarrow \left[-\frac{1}{r} \partial_r r + \frac{\ell(\ell + 1)}{r^2} \right] n_\ell(r) &= +n_\ell(r) \end{aligned} \quad (\text{A.3})$$

Note that the subroutine SPFUNCTION\$NEUMANN returns the Neumann function with the opposite sign, namely what Abramowitz defines as Bessel function of the second kind. The minus sign is added in the calling routine.

- The function

$$m_\ell(r) = r^\ell \left(-\frac{1}{r} \partial_r \right)^\ell \frac{1}{r} e^{-r} \quad (\text{A.4})$$

used for $k^2 < 0$ is obeys the radial Helmholtz equation (Abramowitz Eq. 10.2.1) for negative kinetic energy

$$\begin{aligned} r^2 \partial_r^2 m_\ell + 2r \partial_r m_\ell - (r^2 + \ell(\ell + 1)) m_\ell &= 0 \\ \Rightarrow \left[-\frac{1}{r} \partial_r r + \frac{\ell(\ell + 1)}{r^2} \right] m_\ell(r) &= -m_\ell(r) \end{aligned} \quad (\text{A.5})$$

They are solutions for negative energy and therefore they fall off exponentially. The solution $m_\ell(r)$ is proportional to the modified spherical Bessel functions of the third kind as defined by Abramowitz[?] in their Eq. 10.2.4.

$$m_\ell(r) = \frac{2}{\pi} \left[\sqrt{\frac{\pi}{2r}} K_{\ell+1}(r) \right] \quad (\text{A.6})$$

which can be verified by comparing the defining equation Eq. A.4 with equations 10.2.24-25 and the definition Eq. 10.2.4 of Abramowitz.

A.1 Modified spherical Bessel functions

The modified spherical Bessel functions of the first kind obey (Abramowitz 10.2.1)

$$\begin{aligned} x^2 \partial_x^2 + 2x \partial_x - (x^2 + \ell(\ell+1)) w(x) &= 0 \\ \Rightarrow \left(\frac{1}{x} \partial_x^2 x - \frac{\ell(\ell+1)}{x^2} - 1 \right) w(x) &= 0 \end{aligned} \quad (\text{A.7})$$

The regular solutions are modified bessel functions of the first kind. (abramowitz 10.2.2, abramowitz 10.2.13)

$$\begin{aligned} j_0(x) &= \sqrt{\frac{\pi}{2x}} I_{\frac{1}{2}}(x) = \frac{\sinh(x)}{x} \\ j_1(x) &= \sqrt{\frac{\pi}{2x}} I_{\frac{3}{2}}(x) = \frac{x \cosh(x) - \sinh(x)}{x^2} \\ j_2(x) &= \sqrt{\frac{\pi}{2x}} I_{\frac{5}{2}}(x) = \frac{x^2 \sinh(x) - 3x \cosh(x) + 3 \sinh(x)}{x^3} \end{aligned} \quad (\text{A.8})$$

They have the leading order (abramowitz 10.2.5)

$$j_\ell(x) = \frac{x^\ell}{(2\ell+1)!!} + O(x^{\ell+2}) \quad (\text{A.9})$$

The irregular solutions are the **modified Bessel functions of the third kind** (abramowitz 10.2.4, abramowitz 10.2.17) $\frac{\pi}{2} h_\ell(x)$. We will name $h_\ell(x)$ Hankel function.

$$\begin{aligned} \frac{\pi}{2} h_0(x) &= \sqrt{\frac{\pi}{2x}} K_{\frac{1}{2}}(x) = \frac{\pi}{2} e^{-x} \frac{1}{x} \\ \frac{\pi}{2} h_1(x) &= \sqrt{\frac{\pi}{2x}} K_{\frac{3}{2}}(x) = \frac{\pi}{2} e^{-x} \frac{1+x}{x^2} \\ \frac{\pi}{2} h_2(x) &= \sqrt{\frac{\pi}{2x}} K_{\frac{5}{2}}(x) = \frac{\pi}{2} e^{-x} \frac{x^2 + 3x + 3}{x^3} \end{aligned} \quad (\text{A.10})$$

They have the leading order (abramowitz 10.2.6, abramowitz 10.2.4))

$$\frac{\pi}{2} h_\ell(x) = \frac{\pi (2\ell-1)!!}{2 x^{\ell+1}} + \dots \quad (\text{A.11})$$

A.2 Bare structure constants

This section is copied from *Methods-book*, Section “Working with spherical Hankel and Bessel functions”, Peter Blöchl, private communication.

The bare structure constants have been determined first by Segall[?] . He uses the theorem[?] that supposedly goes back to Kasterin (N. Kasterin, Proc. Acad. Sci Amsterdam 6, 460 (1897/98)); see Seegall[?] , Eq. B4)

$$h_\ell^{(1)}(\kappa r)Y_{\ell,m}(\vec{r}) = i^{-\ell}\mathcal{Y}_{\ell,m}(\vec{\nabla}_r)h_0^{(1)}(\kappa r) \quad (\text{A.12})$$

where $h_\ell^{(1)}(x)$ is the spherical Hankel function of the first kind (see Eq. A.18 below) and where (Eq. B5 of Segall[?])

$$\mathcal{Y}_{\ell,m}(\vec{\nabla}) = \sqrt{\frac{2\ell+1}{4\pi} \frac{(\ell-m)!}{(\ell+m)!}} \left(\frac{1}{ik}\right)^{|m|} \left(\partial_x \pm i\partial_y\right) \mathcal{P}_\ell^{|m|} \left(\frac{1}{ik}\partial_z\right) \quad (\text{A.13})$$

where the positive sign applies for nonzero m and the negative sign for negative m . Furthermore (see Segall[?] Eq. B5)

$$\mathcal{P}_\ell^{|m|}(z) = \frac{d^{|m|}P_\ell(z)}{dz^{|m|}}$$

where $P_\ell(z)$ is the conventional Legendre polynomial.

In addition Segall[?] refers in his Eq. B7 to Morse and Feshbach[?] (part II, p. 1574) for

$$h_0^{(1)}(\kappa|\vec{r}-\vec{r}'|) = 4\pi \sum_L \left(h_\ell^{(1)}(\kappa|\vec{r}'|)Y_L(\vec{r}') \right) j_\ell(\kappa|\vec{r}|)Y_L^*(\vec{r}) \quad (\text{A.14})$$

which is valid for $|\vec{r}'| > |\vec{r}|$.

The two equations, Eq. A.12 and Eq. A.14, can be combined into

$$\begin{aligned} h_\ell^{(1)}(\kappa|\vec{r}|)Y_{\ell,m}(\vec{r}) &\stackrel{\text{Eq. A.12}}{=} i^{-\ell}\mathcal{Y}_{\ell,m}(\vec{\nabla}_r)h_0^{(1)}(\kappa|\vec{r}|) \\ &= i^{-\ell}\mathcal{Y}_{\ell,m}(\vec{\nabla}_r)h_0^{(1)}(\kappa|\vec{r}-\vec{R}|) \\ &\stackrel{\text{Eq. A.14}}{=} i^{-\ell}\mathcal{Y}_{\ell,m}(\vec{\nabla}_r) \left[4\pi \sum_{L'} \left(h_{\ell'}^{(1)}(\kappa|\vec{R}|)Y_{L'}(-\vec{R}) \right) j_{\ell'}(\kappa|\vec{r}-\vec{R}|)Y_{L'}^*(\vec{r}-\vec{R}) \right] \\ &\stackrel{\text{Eq. A.16}}{=} 4\pi \sum_{L'} \left(i^{-\ell}\mathcal{Y}_{\ell,m}(\vec{\nabla}_R)h_{\ell'}^{(1)}(\kappa|\vec{R}|)Y_{L'}(-\vec{R}) \right) j_{\ell'}(\kappa|\vec{r}-\vec{R}|)Y_{L'}^*(\vec{r}-\vec{R}) \end{aligned}$$

Here we used that

$$\vec{\nabla}_r[f(\vec{R})g(\vec{r}-\vec{R})] = f(\vec{R})\vec{\nabla}_r g(\vec{r}-\vec{R}) = -f(\vec{R})\vec{\nabla}_R g(\vec{r}-\vec{R}) \quad (\text{A.15})$$

$$= -\underbrace{\vec{\nabla}_R \left[\overbrace{f(\vec{R})g(\vec{r}-\vec{R})}^{h(\vec{r})} \right]}_{=0} + \left[\vec{\nabla}_R f(\vec{R}) \right] g(\vec{r}-\vec{R}) \quad (\text{A.16})$$

We summarize the final result

CONDITION FOR STRUCTURE CONSTANTS (POSITIVE ENERGIES)

$$h_\ell^{(1)}(\kappa|\vec{r}|)Y_{\ell,m}(\vec{r}) = 4\pi \sum_{L'} \left(i^{-\ell} \mathcal{Y}_{\ell,m}(\vec{\nabla}_R) h_{\ell'}^{(1)}(\kappa|\vec{R}|) Y_{L'}(-\vec{R}) \right) j_{\ell'}(\kappa|\vec{r} - \vec{R}|) Y_{L'}^*(\vec{r} - \vec{R}) \quad (\text{A.17})$$

where $h^{(1)}(x)$ is the spherical Hankel function of the first kind defined in Abramowitz and Stegun (AS)[?]]

$$h_\ell^{(1)}(x) \stackrel{\text{AS10.1.1}}{=} j_\ell(x) + i y_\ell(x) \stackrel{\text{AS10.1.26}}{=} x^\ell \left(-\frac{1}{x} \partial_x \right)^\ell \frac{\sin(x) - i \cos(x)}{x} \quad (\text{A.18})$$

Expression for the structure constants

By comparing our notation to that of Daniel Grieger and using his expression for the Structure constants, we arrive at the following expression for the structure constants in our notation.

There was a misunderstanding with the sign of the structure constants. Here I follow the signconvention $K = -\sum JS$, which is opposite to the one I and Daniel had earlier.

$$S_{R',L',R,L} = -4\pi \sum_{L''} H_{L''}^B(\vec{R}' - \vec{R}) C_{L,L'',L} \left\{ \begin{array}{l} (-1)^{\ell'} (-ik)^{\ell+\ell'-\ell''} \\ (-1)^{\ell'} \delta_{\ell+\ell',\ell''} \\ (-1)^{\ell'} \kappa^{\ell+\ell'-\ell''} \end{array} \right\} \quad (\text{A.19})$$

A.3 Consistency checks

We consider the case with $\kappa = 0$, for which the solid Bessel and Hankel functions are

$$K_{0,L}^\infty(\vec{r}) = (2\ell - 1)!! \frac{1}{|\vec{r}|^{\ell+1}} Y_L(\vec{r}) \quad (\text{A.20})$$

$$J_{0,L}(\vec{r}) = \frac{1}{(2\ell + 1)!!} |\vec{r}|^\ell Y_L(\vec{r}) \quad (\text{A.21})$$

The explicit form of the first few is

$$K_{0,s}^\infty(\vec{r}) = \frac{1}{\sqrt{4\pi}} \frac{1}{|\vec{r}|} \quad (\text{A.22})$$

$$K_{0,p_x}^\infty(\vec{r}) = \sqrt{\frac{3}{4\pi}} \frac{x}{|\vec{r}|^3} \quad (\text{A.23})$$

$$J_{0,s}(\vec{r}) = \frac{1}{\sqrt{4\pi}} \quad (\text{A.24})$$

$$J_{0,p_x}(\vec{r}) = \frac{1}{3} \sqrt{\frac{3}{4\pi}} x \quad (\text{A.25})$$

Now we extract the structure constants from the off-site expansion

$$K_{\vec{0},s}^{\infty}(\vec{r}) = -S_{\vec{0},s;\vec{R},s}J_s(\vec{r}-\vec{R}) - S_{\vec{0},s;\vec{R},p_x}J_{p_x}(\vec{r}-\vec{R}) - S_{\vec{0},s;\vec{R},p_y}J_{p_y}(\vec{r}-\vec{R}) - S_{\vec{0},s;\vec{R},p_z}J_{p_z}(\vec{r}-\vec{R}) \quad (\text{A.26})$$

which allows us to evaluate the structure constants directly calculating value and derivatives at the second center and by exploiting selection rules¹

$$K_{\vec{0},s}^{\infty}(\vec{R}) = \frac{1}{\sqrt{4\pi}} \frac{1}{|\vec{R}|} = - \underbrace{\left(-\frac{1}{|\vec{R}|}\right)}_{S_{\vec{0},s;\vec{R},s}} \underbrace{\frac{1}{\sqrt{4\pi}}}_{J_{\vec{R},s}(\vec{R})} \quad (\text{A.27})$$

$$\partial_x|_{\vec{R}} K_{\vec{0},s}^{\infty} = -\frac{1}{\sqrt{4\pi}} \frac{X}{|\vec{R}|^3} = - \underbrace{\sqrt{3} \frac{X}{|\vec{R}|^3}}_{S_{\vec{0},s;\vec{R},p_x}} \underbrace{\frac{1}{3} \sqrt{\frac{3}{4\pi}}}_{\partial_x J_{\vec{R},p_x}(\vec{R})} \quad (\text{A.28})$$

$$K_{\vec{0},p_x}(\vec{R}) = \sqrt{\frac{3}{4\pi}} \frac{X}{|\vec{R}|^3} = - \underbrace{\left(-\sqrt{3} \frac{X}{|\vec{R}|^3}\right)}_{S_{\vec{0},p_x;\vec{R},s}} \underbrace{\frac{1}{\sqrt{4\pi}}}_{J_{\vec{R},s}(\vec{R})}$$

$$\partial_x|_{\vec{R}} K_{\vec{0},p_x}^{\infty} = \sqrt{\frac{3}{4\pi}} \left(\frac{1}{|\vec{R}|^3} - 3 \frac{X^2}{|\vec{R}|^5} \right) = - \underbrace{3 \frac{3X^2 - \vec{R}^2}{|\vec{R}|^2}}_{S_{\vec{0},p_x;\vec{R},p_x}} \underbrace{\frac{1}{3} \sqrt{\frac{3}{4\pi}} |\vec{R}|^{-3}}_{\partial_x|_{\vec{R}} J_{\vec{R},p_x}} \quad (\text{A.29})$$

Thus, the matrix of structure constants in the (s,p_x) subspace is

$$\mathbf{S}_{\vec{0},\vec{R}} = \begin{pmatrix} -|\vec{R}|^{-1} & \sqrt{3}X/|\vec{R}|^3 \\ -\sqrt{3}X/|\vec{R}|^3 & 3[3X^2/R^2 - 1] \end{pmatrix} \quad (\text{A.30})$$

We compare this result now for the one obtained from the direct formula for $\kappa = 0$. These structure constants have the form

$$S_{RL,R'L'} = (-1)^{\ell'+1} 4\pi \sum_{L''} C_{L,L',L''} H_{L''}(\vec{R}' - \vec{R}) \delta^{\ell+\ell'-\ell''} \quad (\text{A.31})$$

¹Only an s-function has a finite value at the origin, only a p-function has a finite first derivative at the center, etc.

The structure constants obtained from this equation are

$$\begin{aligned}
 S_{\vec{0},s;\vec{R},s} &= (-1)4\pi \underbrace{\frac{1}{\sqrt{4\pi}}}_{C_{sss}} \cdot \underbrace{\frac{1}{\sqrt{4\pi}} \frac{1}{|\vec{R}|}}_{H_s(\vec{R})} = -\frac{1}{|\vec{R}|} \\
 S_{\vec{0},s;\vec{R},p_x} &= 4\pi \underbrace{\frac{1}{\sqrt{4\pi}}}_{C_{p_x,s,p_x}} \underbrace{\sqrt{\frac{3}{4\pi}} \frac{X}{|\vec{R}|^3}}_{H_{p_x}(\vec{R})} = \sqrt{3} \frac{X}{|\vec{R}|^3} \\
 S_{\vec{0},p_x;\vec{R},s} &= (-1)4\pi \underbrace{\frac{1}{\sqrt{4\pi}}}_{C_{p_x,s,s}} \sqrt{\frac{3}{4\pi}} \frac{X}{|\vec{R}|^3} = -\sqrt{3} \frac{X}{|\vec{R}|^3} \\
 S_{\vec{0},p_x;\vec{R},p_x} &= 4\pi \underbrace{\frac{1}{\sqrt{5\pi}}}_{C_{p_x,p_x,d_{3x^2-r^2}}} \underbrace{\overbrace{\sqrt{\frac{5}{16\pi}} \frac{3X^2-R^2}{|\vec{R}|^2}}^{Y_{3x^2-r^2}} \frac{3}{|\vec{R}|^3}}_{H_{3x^2-r^2}(\vec{R})} = 3 \frac{3X^2-R^2}{|\vec{R}|^5} \quad (\text{A.32})
 \end{aligned}$$

For Gaunt coefficients see footnote.²

2

$$\begin{aligned}
 Y_{p_x} Y_{p_x} &= \frac{3}{4\pi} \frac{x^2}{r^2} = \frac{1}{4\pi} \frac{x^2}{r^2} + \frac{1}{4\pi} \frac{3x^2-r^2}{r^2} = \frac{1}{\sqrt{4\pi}} Y_s + \frac{1}{4\pi} \sqrt{\frac{16\pi}{5}} Y_{3x^2-r^2} = \frac{1}{\sqrt{4\pi}} Y_s + \sqrt{\frac{1}{5\pi}} Y_{3x^2-r^2} \\
 &\Rightarrow C_{p_x,p_x,s} = \frac{1}{\sqrt{4\pi}} \quad \text{and} \quad C_{p_x,p_x,d_{3x^2-r^2}} = \frac{1}{\sqrt{5\pi}} \quad (\text{A.33})
 \end{aligned}$$

Appendix B

Bloch theorem revisited

The Bloch states are eigenstates of the discrete lattice translation

$$\hat{S}(\vec{t}) = \int d^3r |\vec{r} + \vec{t}\rangle \langle \vec{r}| \quad (\text{B.1})$$

for the discrete lattice vectors \vec{t} . The eigenvalue equation has the form

$$\hat{S}(\vec{t})|\psi_{\vec{k}}\rangle = |\psi_{\vec{k}}\rangle e^{i\vec{k}\vec{t}} \quad (\text{B.2})$$

This eigenvalue equation can be recast into the form

$$\langle \vec{r} - \vec{t} | \psi_{\vec{k}} \rangle = \langle \vec{r} | \psi_{\vec{k}} \rangle e^{i\vec{k}\vec{t}} \quad (\text{B.3})$$

This implies that the states can be written as product of a periodic function and a phase factor

$$\langle \vec{r} | \psi_{\vec{k}} \rangle = u_{\vec{k}}(\vec{r}) e^{i\vec{k}\vec{r}} \quad (\text{B.4})$$

with

$$u_{\vec{k}}(\vec{r}) = u_{\vec{k}}(\vec{r} + \vec{t}) \quad (\text{B.5})$$

Bloch theorem in a local orbital basis

With $q_{\alpha} \stackrel{\text{def}}{=} \langle \pi_{\alpha} | \psi \rangle$, we obtain

$$\begin{aligned} \hat{S}(\vec{t}) \sum_{\alpha} |\chi_{\alpha}\rangle q_{\alpha,n} &= \sum_{\alpha} |\chi_{\alpha}\rangle q_{\alpha,n} e^{i\vec{k}_n \vec{t}} \\ \int d^3r |\vec{r} + \vec{t}\rangle \langle \vec{r}| \sum_{\alpha} |\chi_{\alpha}\rangle q_{\alpha,n} &= \int d^3r |\vec{r}\rangle \langle \vec{r}| \sum_{\alpha} |\chi_{\alpha}\rangle q_{\alpha,n} e^{i\vec{k}_n \vec{t}} \\ \sum_{\alpha} \langle \vec{r} - \vec{t} | \chi_{\alpha} \rangle q_{\alpha,n} &= \sum_{\alpha} \langle \vec{r} | \chi_{\alpha} \rangle q_{\alpha,n} e^{i\vec{k}_n \vec{t}} \\ \sum_{\alpha} \langle \vec{r} | \chi_{\alpha+\vec{t}} \rangle q_{\alpha,n} &= \sum_{\alpha} \langle \vec{r} | \chi_{\alpha} \rangle q_{\alpha,n} e^{i\vec{k}_n \vec{t}} \\ \sum_{\alpha'} \langle \vec{r} | \chi_{\alpha'} \rangle q_{\alpha'+\vec{t},n} &= \sum_{\alpha} \langle \vec{r} | \chi_{\alpha} \rangle q_{\alpha,n} e^{i\vec{k}_n \vec{t}} \\ q_{\alpha+\vec{t},n} &= q_{\alpha,n} e^{-i\vec{k}_n \vec{t}} \end{aligned} \quad (\text{B.6})$$

Density matrix

$$\rho_{\alpha,\beta+\vec{t}} = \sum_n \langle \pi_\alpha | \psi_n \rangle f_n \langle \psi_n | \pi_\beta \rangle e^{+i\vec{k}_n \vec{t}} \quad (\text{B.7})$$

Appendix C

Offsite matrix elements using Gaussian integrals

In LMTO_INITIALIZE there are non-functional calls for doing the integrations in a representation of Gauss orbitals. The routines are no more present, except for

- LMTO_TAILEDPRODUCTS. They have been removed in 7501a0g from Feb.2, 2013 (svn revision 1116 from Nov. 20, 2011). LMTO_TAILEDPRODUCTS has been removed after 9803b02 on from Mar. 1, 2014.
- LMTO_EXPANDPRODS

The routines related to Gaussians have been moved into the file `paw_lmto_stuffwithgaussians.f90`

```
!!$      CALL LMTO_TAILEDPRODUCTS()  
          gaussian_fitgauss  
!!$      CALL LMTO_GAUSSFITKPRIME()  
!!$      CALL LMTO_GAUSSFITKAUGMENT()  
!!$      CALL LMTO_GAUSSFITKJTails()  
!!$      CALL LMTO_ONSITEOVERLAP()
```

Appendix D

Double counting

D.1 Other double-counting schemes

An excellent review of double-counting terms has been discussed by Nekrasov et al.[?].

Around-mean-field (AMF) and fully-localized limit (FLL)

In LDA+U, two double-counting schemes are in common use: one is called around mean-field (AMF)[?] and the other is called fully localized limit (FLL)[?].

In the Hubbard model, the two approximations have the form[?].

$$\begin{aligned}\hat{H}_{AMF}^{dc} &= \frac{1}{2}U \sum_{\sigma} \hat{n}_{d,\sigma}(\hat{n}_d - n_{\sigma}^0) - \frac{1}{2}J \sum_{\sigma} \hat{n}_{d,\sigma}(\hat{n}_{d,\sigma} - n_{\sigma}^0) \\ \hat{H}_{FLL}^{dc} &= \frac{1}{2}U n_d(n_d - 1) - \frac{1}{2}J \sum_{\sigma} \hat{n}_{d,\sigma}(\hat{n}_{d,\sigma} - 1)\end{aligned}\tag{D.1}$$

with

$$\begin{aligned}n_{d,\sigma} &= \sum_m \langle \hat{n}_{m,\sigma} \rangle \\ n_d &= \sum_{\sigma} n_{d,\sigma} \\ n_{\sigma}^0 &= \frac{1}{2(2\ell + 1)} \sum_{m,\sigma} n_{m,\sigma} \\ n^0 &= \sum_{\sigma} n_{\sigma}^0\end{aligned}\tag{D.2}$$

It is still unclear what are operators and what are numbers, but I believe all are numbers.

Nekrasov, Pavlov Sadovskii scheme

Nekrasov et al.[?] proposes a new scheme which amounts to removing the density of the correlated orbitals from the integral for the exchange-correlation term. The corresponding double-

counting energy would be

$$E^{DC} = E_{xc}[n_t - n_{corr}] - E_{xc}[n_t] \quad (D.3)$$

where n_t is the density of all orbitals, while n_{corr} is the density of the correlated orbitals only.

Blöchl-Walther-Pruschke scheme

Our scheme[?] differs in that it divides the DFT exchange correlation energy based on a partitioning of the two-particle density, where $n(\vec{r}) = \sum_R n_R(\vec{r})$

$$\begin{aligned} E_{xc} &= \int d^3r n(\vec{r}) \frac{1}{2} \int d^3r' \frac{e^2 \int_0^1 d\lambda h_\lambda(\vec{r}, \vec{r}')}{4\pi\epsilon|\vec{r} - \vec{r}'|} \\ &= \frac{1}{2} \sum_{R,R'} \int d^3r \int d^3r' \left(\frac{n_R(\vec{r})}{n(\vec{r})} \right) \left(n(\vec{r}) \frac{e^2 \int_0^1 d\lambda h_\lambda(\vec{r}, \vec{r}')}{4\pi\epsilon|\vec{r} - \vec{r}'|} \right) \left(\frac{n_{R'}(\vec{r}')}{n(\vec{r}')} \right) \end{aligned} \quad (D.4)$$

This formulation takes into account that the electrons are indistinguishable and that each electron contributes equally to the exchange correlation energy. Furthermore, the partitions of the expression add up to the total exchange-correlation energy (when also the inter-site terms are included).

The model rests on a well defined expression for the two-particle density, namely

$$n^{(2)}(\vec{r}, \vec{r}') = n^{(1)}(\vec{r})n^{(1)}(\vec{r}') + \frac{1}{2} \sum_{R,R'} \left(\frac{n_R(\vec{r})}{n(\vec{r})} \right) \left(n(\vec{r}) h_{\lambda=1}(\vec{r}, \vec{r}') \right) \left(\frac{n_{R'}(\vec{r}')}{n(\vec{r}')} \right) \quad (D.5)$$

and an expression for the kinetic-energy correction

$$T - T_s = \frac{1}{2} \sum_{R,R'} \int d^3r \int d^3r' \left(\frac{n_R(\vec{r})}{n(\vec{r})} \right) \left(n(\vec{r}) \frac{e^2 \int_0^1 d\lambda \left(h_\lambda(\vec{r}, \vec{r}') - h_{\lambda=1}(\vec{r}, \vec{r}') \right)}{4\pi\epsilon|\vec{r} - \vec{r}'|} \right) \left(\frac{n_{R'}(\vec{r}')}{n(\vec{r}')} \right) \quad (D.6)$$

as well as the potential energy of exchange and correlation

$$U_{xc} = \frac{1}{2} \sum_{R,R'} \int d^3r \int d^3r' \left(\frac{n_R(\vec{r})}{n(\vec{r})} \right) \left(n(\vec{r}) \frac{e^2 \int_0^1 d\lambda h_{\lambda=1}(\vec{r}, \vec{r}')}{4\pi\epsilon|\vec{r} - \vec{r}'|} \right) \left(\frac{n_{R'}(\vec{r}')}{n(\vec{r}')} \right) \quad (D.7)$$

Note, however, that only the total exchange-correlation energy including Coulomb and kinetic energy contributions is a density functional, whereas both contributions can be expressed individually as density-matrix functional.

Only local terms

The equations above reproduce the full exchange correlation energy, respectively the kinetic energy correction as double sum over local terms.

If we employ the local approximation, only the diagonal terms in R, R' need to be considered. Thus, a double-counting correction for a set of correlated orbitals, which contribute n_R to the total density would be

$$E_{dc} = -\frac{1}{2} \sum_R \int d^3r \int d^3r' \left(\frac{n_R(\vec{r})}{n(\vec{r})} \right) \left(n(\vec{r}) \frac{e^2 \int_0^1 d\lambda h_\lambda(\vec{r}, \vec{r}')}{4\pi\epsilon |\vec{r} - \vec{r}'|} \right) \left(\frac{n_R(\vec{r}')}{n(\vec{r}')} \right) \quad (D.8)$$

This integral can be approximated further as

$$E_{xc} = -\frac{1}{2} \sum_R \int d^3r n(\vec{r}) \epsilon[n(\vec{r})] \left(\frac{n_R(\vec{r})}{n(\vec{r})} \right)^2 \quad (D.9)$$

D.1.1 Around mean field (AMF)

The original expression, AMF, for the double-counting term is derived from the mean-field approximation of the Hubbard model.

This section is unfinished. It is an attempt to derive the AMF limit consistent with our U-tensor We start out from the interaction

$$\hat{W} = \frac{1}{2} \sum_{i,j,k,l} W_{i,j,l,k} \hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l \quad (D.10)$$

with

$$\begin{aligned} W_{i,j,k,l} &= \int d^4x \int d^4x' \frac{e^2 \chi_i^*(\vec{x}) \chi_j^*(\vec{x}') \chi_k(\vec{x}) \chi_l(\vec{x}')}{4\pi\epsilon_0 |\vec{r} - \vec{r}'|} \\ &= \delta_{\sigma_i, \sigma_k} \delta_{\sigma_j, \sigma_l} \int d^3r \int d^3r' \frac{e^2 \phi_i^*(\vec{r}) \phi_j^*(\vec{r}') \phi_k(\vec{r}) \phi_l(\vec{r}')}{4\pi\epsilon_0 |\vec{r} - \vec{r}'|} = \delta_{\sigma_i, \sigma_k} \delta_{\sigma_j, \sigma_l} U_{i,j,k,l} \end{aligned} \quad (D.11)$$

where the spin-orbitals $\chi_i(\vec{r}, \sigma) = \phi_i(\vec{r}) \delta_{\sigma, \sigma_i}$ are decomposed into a outer product of a spatial orbital $\phi_i(\vec{r})$ and a spinor. The interaction has the internal symmetry

$$W_{i,j,k,l} = W_{j,i,l,k} = W_{k,l,i,j}^* = W_{l,k,j,i}^* \quad (D.12)$$

In order to arrive at the mean-field Hamiltonian, we express the wave function in terms of wave function of a non-interacting system, i.e. by Slater determinants. This non-interacting system is then optimized to minimize the expectation value of the Hamiltonian containing the interaction. (See Bogoljubov inequality) The result is the Hartree-Fock approximation.

The ground state is

$$|\Phi_0\rangle = \prod_n (\hat{a}_n^\dagger)^{\sigma_n} |\mathcal{O}\rangle \quad (D.13)$$

where $\vec{\sigma} \in \{0, 1\}^\infty$ is the occupation number representation of the Slater determinant. The orbitals making up the Slater determinant are $|\phi_n\rangle$

$$\hat{a}_n^\dagger = \sum_j \hat{c}_j^\dagger \langle \pi_j | \psi_n \rangle \quad (D.14)$$

with a complete set of one-particle orbitals and the bi-orthogonality condition this relationship can be inverted

$$\sum_n \hat{a}_n^\dagger \langle \psi_n | \chi_i \rangle = \sum_j \hat{c}_j^\dagger \langle \pi_j | \underbrace{\sum_n |\psi_n\rangle \langle \psi_n|}_{=1} | \chi_i \rangle = \hat{c}_i^\dagger \quad (\text{D.15})$$

$\underbrace{\hspace{10em}}_{\delta_{i,j}}$

$$\begin{aligned} \langle \Phi | \hat{a}_m^\dagger \hat{a}_n^\dagger \hat{a}_o \hat{a}_p | \Phi \rangle &= \delta_{m,o} \delta_{n,p} \langle \Phi | \hat{a}_m^\dagger \underbrace{\hat{a}_n^\dagger \hat{a}_m}_{\delta_{m,n} - \hat{a}_m \hat{a}_n^\dagger} \hat{a}_n | \Phi \rangle + \delta_{m,p} \delta_{n,o} \langle \Phi | \hat{a}_m^\dagger \underbrace{\hat{a}_n^\dagger \hat{a}_n \hat{a}_m}_{-(\delta_{m,n} - \hat{a}_m \hat{a}_n^\dagger) \hat{a}_n} | \Phi \rangle \\ &\quad - \delta_{m,p} \delta_{n,o} \delta_{m,n} \langle \Phi | \hat{a}_m^\dagger \hat{a}_m^\dagger \underbrace{\hat{a}_m \hat{a}_m}_{=0} | \Phi \rangle \\ &= \delta_{m,o} \delta_{n,p} \left[\delta_{m,n} \langle \Phi | \hat{a}_m^\dagger \hat{a}_n | \Phi \rangle - \langle \Phi | \hat{a}_m^\dagger \hat{a}_m | \Phi \rangle \langle \Phi | \hat{a}_n^\dagger \hat{a}_n | \Phi \rangle \right] \\ &\quad - \delta_{m,p} \delta_{n,o} \left[\delta_{m,n} \langle \Phi | \hat{a}_m^\dagger \hat{a}_n | \Phi \rangle - \langle \Phi | \hat{a}_m^\dagger \hat{a}_m | \Phi \rangle \langle \Phi | \hat{a}_n^\dagger \hat{a}_n | \Phi \rangle \right] \\ &= -\delta_{m,o} \delta_{n,p} \langle \Phi | \hat{a}_m^\dagger \hat{a}_m | \Phi \rangle \langle \Phi | \hat{a}_n^\dagger \hat{a}_n | \Phi \rangle + \delta_{m,p} \delta_{n,o} \langle \Phi | \hat{a}_m^\dagger \hat{a}_m | \Phi \rangle \langle \Phi | \hat{a}_n^\dagger \hat{a}_n | \Phi \rangle \\ &= f_m f_n \left[\delta_{m,o} \delta_{n,p} - \delta_{m,p} \delta_{n,o} \right] \quad (\text{D.16}) \end{aligned}$$

Here the occupations are those of the reference Slater determinant, the mean field.

Now we can evaluate the matrix elements for the interaction as

$$\begin{aligned} \langle \Phi_0 | \hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l | \Phi_0 \rangle &= \sum_{m,n,o,p} \langle \psi_m | \chi_i \rangle \langle \psi_n | \chi_j \rangle \langle \Phi_0 | \hat{a}_m^\dagger \hat{a}_n^\dagger \hat{a}_o \hat{a}_p | \Phi_0 \rangle \langle \chi_k | \psi_o \rangle \langle \chi_l | \psi_p \rangle \\ &= \sum_{m,n,o,p} \langle \psi_m | \chi_i \rangle \langle \psi_n | \chi_j \rangle \left[f_m f_n \left[\delta_{m,o} \delta_{n,p} - \delta_{m,p} \delta_{n,o} \right] \right] \langle \chi_k | \psi_o \rangle \langle \chi_l | \psi_p \rangle \\ &= \sum_{m,n} f_m f_n \langle \psi_m | \chi_i \rangle \langle \psi_n | \chi_j \rangle \langle \chi_k | \psi_m \rangle \langle \chi_l | \psi_n \rangle - \sum_{m,n} f_m f_n \langle \psi_m | \chi_i \rangle \langle \psi_n | \chi_j \rangle \langle \chi_k | \psi_n \rangle \langle \chi_l | \psi_m \rangle \\ &= \underbrace{\sum_m \langle \chi_k | \psi_m \rangle f_m \langle \psi_m | \chi_i \rangle}_{\rho_{k,i}} \underbrace{\sum_m \langle \chi_l | \psi_m \rangle f_m \langle \psi_m | \chi_j \rangle}_{\rho_{l,j}} - \underbrace{\sum_m \langle \chi_l | \psi_m \rangle f_m \langle \psi_m | \chi_i \rangle}_{\rho_{l,i}} \underbrace{\sum_n \langle \chi_k | \psi_n \rangle f_n \langle \psi_n | \chi_j \rangle}_{\rho_{k,j}} \\ &= \rho_{k,i} \rho_{l,j} - \rho_{l,i} \rho_{k,j} \\ &= \langle \Phi_0 | \hat{c}_i^\dagger \hat{c}_k | \Phi_0 \rangle \langle \Phi_0 | \hat{c}_j^\dagger \hat{c}_l | \Phi_0 \rangle - \langle \Phi_0 | \hat{c}_i^\dagger \hat{c}_l | \Phi_0 \rangle \langle \Phi_0 | \hat{c}_j^\dagger \hat{c}_k | \Phi_0 \rangle \quad (\text{D.17}) \end{aligned}$$

This is the expectation value of a Slater determinant which is the zero'th order term for an expansion in the deviation from these mean values.

The density matrix has the form

$$\rho_{i,j} = \langle \Phi_0 | \hat{c}_j^\dagger \hat{c}_i | \Phi_0 \rangle = \sum_{m,n} \langle \psi_m | \chi_j \rangle \langle \chi_i | \psi_n \rangle \underbrace{\langle \Phi_0 | \hat{a}_m^\dagger \hat{c}_n | \Phi_0 \rangle}_{\delta_{m,n} f_n} = \sum_n \chi_i | \psi_n \rangle f_n \langle \psi_n | \chi_j \rangle \quad (\text{D.18})$$

$$\begin{aligned} \hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l &= \hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l \\ &= \langle \Phi_0 | \hat{c}_i^\dagger \hat{c}_k | \Phi_0 \rangle \langle \Phi_0 | \hat{c}_j^\dagger \hat{c}_l | \Phi_0 \rangle - \langle \Phi_0 | \hat{c}_i^\dagger \hat{c}_l | \Phi_0 \rangle \langle \Phi_0 | \hat{c}_j^\dagger \hat{c}_k | \Phi_0 \rangle \quad (\text{D.19}) \end{aligned}$$

interaction in terms of products of one-particle operators $\hat{c}_i^\dagger \hat{c}_j$. There are two equivalent forms for this, namely

$$\begin{aligned}\hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l &= \delta_{j,k} \hat{c}_i^\dagger \hat{c}_l - \hat{c}_i^\dagger \hat{c}_k \hat{c}_j^\dagger \hat{c}_l \\ &= \delta_{j,l} \hat{c}_i^\dagger \hat{c}_k - \hat{c}_i^\dagger \hat{c}_l \hat{c}_j^\dagger \hat{c}_k\end{aligned}\quad (\text{D.20})$$

Similarly the products of the field operators can be rewritten in a number of different ways

$$\begin{aligned}\hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_k \hat{c}_l &= -\hat{c}_i^\dagger \hat{c}_j^\dagger \hat{c}_l \hat{c}_k = -\hat{c}_j^\dagger \hat{c}_i^\dagger \hat{c}_k \hat{c}_l = \hat{c}_j^\dagger \hat{c}_i^\dagger \hat{c}_l \hat{c}_k \\ &= \delta_{j,k} \hat{c}_i^\dagger \hat{c}_l - \hat{c}_i^\dagger \hat{c}_k \hat{c}_j^\dagger \hat{c}_l =\end{aligned}\quad (\text{D.21})$$

$$\begin{aligned}&= \frac{1}{2} \sum_{i,j,k,l} W_{i,j,l,k} \frac{1}{2} \left[-\left(\hat{c}_i^\dagger \hat{c}_k \hat{c}_j^\dagger \hat{c}_l - \delta_{k,j} \hat{c}_i^\dagger \hat{c}_l \right) + \left(\hat{c}_i^\dagger \hat{c}_l \hat{c}_j^\dagger \hat{c}_k - \delta_{j,l} \hat{c}_i^\dagger \hat{c}_k \right) \right] \\ &\stackrel{i \leftrightarrow l}{=} \frac{1}{2} \sum_{i,j,k,l} \frac{W_{i,j,k,l} - W_{i,j,l,k}}{2} \left(\hat{n}_{i,k} \hat{n}_{j,l} - \delta_{k,j} \hat{n}_{i,l} \right)\end{aligned}\quad (\text{D.22})$$

In the mean-field approximation, terms that are quadratic in $\hat{n} - \langle \hat{n} \rangle$ are ignored.

$$\begin{aligned}\hat{W}_{MF} &= \frac{1}{2} \sum_{i,j,k,l} \frac{W_{i,j,k,l} - W_{i,j,l,k}}{2} \left(\langle \hat{n}_{i,k} \rangle \langle \hat{n}_{j,l} \rangle - \delta_{k,j} \langle \hat{n}_{i,l} \rangle \right) \\ &+ \frac{1}{2} \sum_{i,j,k,l} \frac{W_{i,j,k,l} - W_{i,j,l,k}}{2} \left([\hat{n}_{i,k} - \langle \hat{n}_{i,k} \rangle] \langle \hat{n}_{j,l} \rangle + \langle \hat{n}_{i,k} \rangle [\hat{n}_{j,l} - \langle \hat{n}_{j,l} \rangle] - \delta_{k,j} [\hat{n}_{i,l} - \langle \hat{n}_{i,l} \rangle] \right) \\ &= \frac{1}{2} \sum_{i,j,k,l} \frac{W_{i,j,k,l} - W_{i,j,l,k}}{2} \left(\hat{n}_{i,k} \langle \hat{n}_{j,l} \rangle + \langle \hat{n}_{i,k} \rangle \hat{n}_{j,l} - \delta_{k,j} \hat{n}_{i,l} - \langle \hat{n}_{i,k} \rangle \langle \hat{n}_{j,l} \rangle \right)\end{aligned}\quad (\text{D.23})$$

The resulting Hamiltonian is a one-particle hamiltonian, which depends parametrically on the occupations., i.e the density matrix.

This Hamiltonian can be derived consistently¹ from the density-matrix functional with the density matrix $\rho_{i,j} = \langle \hat{c}_i^\dagger \hat{c}_j \rangle$

$$E_{AMF}^{DC} = \frac{1}{2} \sum_{i,j,k,l} \frac{W_{i,j,k,l} - W_{i,j,l,k}}{2} \left(\rho_{i,k} \rho_{j,l} - \delta_{k,j} \rho_{i,l} \right) \quad (\text{D.24})$$

In order to arrive at the common expression, we consider the **density-density approximation**, that is we consider only density matrices that are diagonal in the orbital and spin coordinates

$$\begin{aligned}E_{AMF}^{DC} &\approx \frac{1}{2} \sum_{\sigma,\sigma'} \sum_{i,j} \frac{W_{i,j,i,j} - W_{i,j,j,i}}{2} \rho_{i,i} \rho_{j,j} - \frac{1}{2} \sum_{i,j} \frac{W_{i,j,i,j} - W_{i,j,j,i}}{2} \rho_{i,i} \\ &= \frac{1}{2} \sum_{i \neq j} \frac{W_{i,j,j,i} - W_{i,j,i,j}}{2} \rho_{i,i} (\rho_{j,j} + 1)\end{aligned}\quad (\text{D.25})$$

¹Both, the the total energy and the derived one-particle hamiltonian is consistent with the mean-field form of the interaction.

The terms with $i = j$ cancel.

$$\begin{aligned}
E_{AMF}^{DC} &\approx \frac{1}{2} \sum_{m,m'} \frac{U_{m,m',m,m'} - U_{m,m',m',m} \delta_{\sigma,\sigma'}}{2} n_{m,\sigma} n_{m',\sigma'} \\
&\quad - \frac{1}{2} \sum_i \left(\sum_{m',\sigma'} \frac{U_{m,m',m',m} - U_{m,m',m,m'} \delta_{\sigma,\sigma'}}{2} \right) n_{m,\sigma} \\
&= \frac{1}{2} \sum_{\sigma} \sum_m \frac{U_{m,m,m,m}}{2} n_{m,\sigma} n_{m,\bar{\sigma}} \\
&\quad + \frac{1}{2} \sum_{m \neq m'} \sum_{\sigma,\sigma'} \frac{U_{m,m',m,m'} - U_{m,m',m',m} \delta_{\sigma,\sigma'}}{2} n_{m,\sigma} n_{m',\sigma'} \\
&\quad - \frac{1}{2} \sum_i \left(\sum_{m'} \frac{2U_{m,m',m',m} - U_{m,m',m,m'}}{2} \right) n_{m,\sigma}
\end{aligned} \tag{D.26}$$

D.1.2 Definition of U and J parameters

The U- and J-parameters for one angular-momentum shell are defined by the U-tensor as follows[?] (see Eq. 5 of Shick et al.)

$$U := \frac{1}{(2\ell + 1)^2} \sum_{m_1, m_2} U_{m_1, m_2, m_1, m_2} \tag{D.27}$$

$$J := U - \frac{1}{2\ell(2\ell + 1)} \sum_{m_1, m_2} (U_{m_1, m_2, m_1, m_2} - U_{m_1, m_2, m_2, m_1}) \tag{D.28}$$

where the Coulomb matrix elements are

$$U_{i,j,k,l} = \int d^3r \int d^3r' \frac{e^2 \chi_i^*(\vec{r}) \chi_j^*(\vec{r}') \chi_k(\vec{r}) \chi_l(\vec{r}')}{4\pi\epsilon_0 |\vec{r} - \vec{r}'|} \tag{D.29}$$

The indices used here are spatial-orbital indices.

Appendix E

Changelog, Bugfixes

- the core-valence exchange contribution differs from the old version, because it also includes the projection on the phidot functions.
- there has been a bug in `lmtoscreen`, which has been fixed with version 3. It may be better to rewrite all structure constants routines with the transposed structure constants.
- in `lmtomakestructureconstants`, the structure constants have not been calculated because the parallelization was wrong. in 13403f6.

```
-      IF (MOD(IAT1-1,NTASKS) .NE. THISTASK-1) THEN  
+      IF (MOD(IAT1-1,NTASKS) .eq. THISTASK-1) THEN
```