

## Foundations of Data Science, Fall 2020

### 3. Linear Regression

Prof. Dan Olteanu

**DaST**  
Data • (Systems+Theory)


Sept 22, 2020

D S  
a T



<https://lms.uzh.ch/url/RepositoryEntry/16830890400>

<https://uzh.zoom.us/j/96690150974?pwd=cnZmMTduWUtCeWoxYW85Z3RMfnpTZz09>



#### Outline

##### Goals

- Review the supervised learning setting
- Describe the linear regression framework
- Apply the linear model to make predictions
- Derive the least squares estimate

##### Supervised Learning Setting

- Data consists of **input** and **output** pairs
- Inputs (also covariates, independent variables, predictors, features)
- Output (also variates, dependent variable, targets, labels)

2

#### Why study linear regression?

- **Least squares** is at least 200 years old going back to Legendre and Gauss
- Francis Galton (1886): "Regression to the mean"
- Often real processes can be **approximated** by linear models
- More complex models require understanding linear regression
- Closed form analytic solutions can be obtained
- Many **key notions** of machine learning can be introduced

3

#### Toy Example: Commute Times

Want to predict commute time into city centre

What variables would be useful?

- Distance to city centre
- Day of the week



##### Data

dist (km)	day	commute time (min)
2.7	fri	25
4.1	mon	33
1.0	sun	15
5.2	tue	45
2.8	sat	22



4

#### Linear Models

Suppose the input is a vector  $\mathbf{x} \in \mathbb{R}^D$  and the output is  $y \in \mathbb{R}$ .

We have data  $(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_N, y_N)$

Notation: data dimension  $D$ , size of dataset  $N$ , column vectors

##### Linear Model

$$y = w_0 + x_1 w_1 + \dots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

5

## Linear Models: Commute Time

### Linear Model

$$y = w_0 + x_1 w_1 + \dots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

Input encoding: mon-sun has to be converted to a number

- monday: 0, tuesday: 1, ..., sunday: 6 BAD encoding!
- One-hot encoding:** Use seven 0-1 features instead
- Simplifying example: 0 if weekend, 1 if weekday

Say  $x_1 \in \mathbb{R}$  (distance) and  $x_2 \in \{0, 1\}$  (weekend/weekday)

Linear model for commute time

$$y = w_0 + x_1 w_1 + x_2 w_2 + \epsilon$$

6

## Linear Model : Adding a feature for bias term

dist	day	commute time	one	dist	day	commute time
$x_1$	$x_2$	$y$	$x_0$	$x_1$	$x_2$	$y$
2.7	fri	25	1	2.7	fri	25
4.1	mon	33	1	4.1	mon	33
1.0	sun	15	1	1.0	sun	15
5.2	tue	45	1	5.2	tue	45
2.8	sat	22	1	2.8	sat	22

### Model

$$y = w_0 + x_1 w_1 + x_2 w_2 + \epsilon$$

### Model

$$y = w_0 x_0 + x_1 w_1 + x_2 w_2 + \epsilon$$

$$= \mathbf{x} \cdot \mathbf{w} + \epsilon$$


7

## Learning Linear Models

Data:  $(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_N, y_N)$ , where  $\mathbf{x}_i \in \mathbb{R}^D$  and  $y_i \in \mathbb{R}$

Model parameter  $\mathbf{w}$ , where  $\mathbf{w} \in \mathbb{R}^D$

Training phase: (learning/estimation  $\mathbf{w}$  from data)



Testing/Deployment phase: (predict  $\hat{y}_{\text{new}} = \mathbf{x}_{\text{new}} \cdot \mathbf{w}$ )

- How different is  $\hat{y}_{\text{new}}$  from  $y_{\text{new}}$  (actual observation)?
- We should keep some data aside for testing before deploying a model


8

## Least Squares Objective Function

$(\mathbf{x}_1, y_1), \dots, (\mathbf{x}_N, y_N)$ , where  $x_i, y_i \in \mathbb{R}$   $\hat{y}(x) = w_0 + x \cdot w_1$  (no noise term)

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\hat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

Predict commute time using only distance



Loss function  
Cost function  
Objective Function  
Energy Function  
Notation -  $\mathcal{L}, J, E, R$


This objective is known as the residual sum of squares or (RSS)

The estimate  $(w_0, w_1)$  is known as the least squares estimate

9

## Least Squares Objective Function

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$



10

$$\mathcal{L}(w_0, w_1) = \frac{1}{2N} \cdot \sum_i (w_0 + x_i \cdot w_1 - y_i)^2 \quad \text{argmin}_{w_0, w_1} \mathcal{L}(w_0, w_1)$$

Partial derivatives:

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{2N} \sum_i (w_0 + x_i \cdot w_1 - y_i) = 0$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{2N} \sum_i (w_0 + x_i \cdot w_1 - y_i) \cdot x_i = 0$$

Normal equations:

$$\begin{cases} w_0 \cdot \frac{\sum 1}{N} + w_1 \cdot \frac{\sum x_i}{N} - \frac{\sum y_i}{N} = 0 \\ w_0 \cdot \frac{\sum x_i}{N} + w_1 \cdot \frac{\sum x_i^2}{N} - \frac{\sum x_i y_i}{N} = 0 \end{cases}$$

$$\text{Var}(x) = \frac{\sum x_i^2}{N} - \bar{x}^2$$

$$\text{Covar}(x, y) = \frac{\sum x_i y_i}{N} - \bar{x} \cdot \bar{y}$$

11

$$\begin{aligned}
 & \begin{cases} w_0 + w_1 \bar{x} - \bar{y} = 0 \\ w_0 \bar{x} + w_1 \frac{\sum x_i^2}{N} - \frac{\sum x_i y_i}{N} = 0 \end{cases} \Rightarrow w_0 = \bar{y} - w_1 \bar{x} \\
 & \underbrace{\bar{y} \bar{x} - w_1 \bar{x}^2 + w_1 \frac{\sum x_i^2}{N} - \frac{\sum x_i y_i}{N}}_{w_1 \cdot \text{Var}(x)} = 0, \\
 & w_1 = \frac{\text{Cov}(x, y)}{\text{Var}(x)} \\
 & \begin{cases} w_1 = \frac{\text{Cov}(x, y)}{\text{Var}(x)} \\ w_0 = \bar{y} - w_1 \bar{x} \end{cases}
 \end{aligned}$$

12

## Computing the Model Parameters: Summary

$$\langle (x_i, y_i) \rangle_{i=1}^N, \text{ where } x_i, y_i \in \mathbb{R} \quad \hat{y}(x) = w_0 + x \cdot w_1 \text{ (no noise term)}$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\hat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i) x_i$$

We obtain the solution for  $(w_0, w_1)$  by setting the partial derivatives to 0 and solving the resulting system. (Normal Equations)

$$w_0 + w_1 \cdot \frac{\sum_i x_i}{N} = \frac{\sum_i y_i}{N} \quad (1)$$

$$w_0 \cdot \frac{\sum_i x_i}{N} + w_1 \cdot \frac{\sum_i x_i^2}{N} = \frac{\sum_i x_i y_i}{N} \quad (2)$$

$$\bar{x} = \frac{\sum_i x_i}{N}$$

$$\bar{y} = \frac{\sum_i y_i}{N}$$

$$\widehat{\text{var}}(x) = \frac{\sum_i x_i^2}{N} - \bar{x}^2$$

$$\widehat{\text{cov}}(x, y) = \frac{\sum_i x_i y_i}{N} - \bar{x} \cdot \bar{y}$$

$$w_1 = \frac{\widehat{\text{cov}}(x, y)}{\widehat{\text{var}}(x)}$$

$$w_0 = \bar{y} - w_1 \cdot \bar{x}$$

13

## Linear Regression : General Case

Recall that the linear model is

$$\hat{y}_i = \sum_{j=0}^D x_{ij} w_j$$

where we assume that  $x_{i0} = 1$  for all  $x_i$ , so that the bias term  $w_0$  does not need to be treated separately.

Expressing everything in matrix notation

$$\hat{\mathbf{y}} = \mathbf{X} \mathbf{w}$$

Here we have  $\hat{\mathbf{y}} \in \mathbb{R}^{N \times 1}$ ,  $\mathbf{X} \in \mathbb{R}^{N \times (D+1)}$  and  $\mathbf{w} \in \mathbb{R}^{(D+1) \times 1}$

$$\begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_N \end{bmatrix} = \begin{bmatrix} \mathbf{x}_1^T \\ \mathbf{x}_2^T \\ \vdots \\ \mathbf{x}_N^T \end{bmatrix} \begin{bmatrix} w_0 \\ \vdots \\ w_D \end{bmatrix} = \begin{bmatrix} x_{10} & \cdots & x_{1D} \\ x_{20} & \cdots & x_{2D} \\ \vdots & \ddots & \vdots \\ x_{N0} & \cdots & x_{ND} \end{bmatrix} \begin{bmatrix} w_0 \\ \vdots \\ w_D \end{bmatrix}$$

14

## Back to Toy Example

one	dist (km)	weekday?	commute time (min)
1	2.7	1 (fri)	25
1	4.1	1 (mon)	33
1	1.0	0 (sun)	15
1	5.2	1 (tue)	45
1	2.8	0 (sat)	22

We have  $N = 5$ ,  $D + 1 = 3$  and so we get

$$\mathbf{y} = \begin{bmatrix} 25 \\ 33 \\ 15 \\ 45 \\ 22 \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & 2.7 & 1 \\ 1 & 4.1 & 1 \\ 1 & 1.0 & 0 \\ 1 & 5.2 & 1 \\ 1 & 2.8 & 0 \end{bmatrix}, \quad \mathbf{w} = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix}$$

$$\text{For } \mathbf{w} = [6.09, 6.53, 2.11]^T, \text{ our predictions would be } \hat{\mathbf{y}} = \begin{bmatrix} 25.83 \\ 34.97 \\ 12.62 \\ 42.16 \\ 24.37 \end{bmatrix}$$

15

## Finding Optimal Solutions using Calculus

$$\begin{aligned}
 \mathcal{L}(\mathbf{w}) &= \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^T \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X} \mathbf{w} - \mathbf{y})^T (\mathbf{X} \mathbf{w} - \mathbf{y}) \\
 &= \frac{1}{2N} (\mathbf{w}^T (\mathbf{X}^T \mathbf{X}) \mathbf{w} - \mathbf{w}^T \mathbf{X}^T \mathbf{y} - \mathbf{y}^T \mathbf{X} \mathbf{w} + \mathbf{y}^T \mathbf{y}) \\
 &= \frac{1}{2N} (\mathbf{w}^T (\mathbf{X}^T \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^T \mathbf{X} \mathbf{w} + \mathbf{y}^T \mathbf{y}) \\
 &= \dots
 \end{aligned}$$

Then, write out all partial derivatives to form the gradient  $\nabla_{\mathbf{w}} \mathcal{L}$

$$\begin{aligned}
 \frac{\partial \mathcal{L}}{\partial w_0} &= \dots \\
 \frac{\partial \mathcal{L}}{\partial w_1} &= \dots \\
 &\vdots \\
 \frac{\partial \mathcal{L}}{\partial w_D} &= \dots
 \end{aligned}$$

Instead, we will use matrix calculus shortcuts to differentiate using matrix notation directly

16

## Differentiating Matrix Expressions

Rules (Tricks)

(i) Linear Form Expressions:  $\nabla_{\mathbf{w}} (\mathbf{c}^T \mathbf{w}) = \mathbf{c}$

$$\mathbf{c}^T \mathbf{w} = \sum_{j=0}^D c_j w_j$$

$$\frac{\partial (\mathbf{c}^T \mathbf{w})}{\partial w_j} = c_j, \quad \text{and so } \nabla_{\mathbf{w}} (\mathbf{c}^T \mathbf{w}) = \mathbf{c} \quad (3)$$

(ii) Quadratic Form Expressions:

$$\nabla_{\mathbf{w}} (\mathbf{w}^T \mathbf{A} \mathbf{w}) = \mathbf{A} \mathbf{w} + \mathbf{A}^T \mathbf{w} \quad (= 2\mathbf{A} \mathbf{w} \text{ for symmetric } \mathbf{A})$$

$$\mathbf{w}^T \mathbf{A} \mathbf{w} = \sum_{i=0}^D \sum_{j=0}^D w_i w_j A_{ij}$$

$$\frac{\partial (\mathbf{w}^T \mathbf{A} \mathbf{w})}{\partial w_k} = \sum_{i=0}^D w_i A_{ik} + \sum_{j=0}^D A_{kj} w_j = \mathbf{A}_{[:,k]}^T \mathbf{w} + \mathbf{A}_{[k,:]} \mathbf{w}$$

$$\nabla_{\mathbf{w}} (\mathbf{w}^T \mathbf{A} \mathbf{w}) = \mathbf{A}^T \mathbf{w} + \mathbf{A} \mathbf{w} \quad (4)$$

17

$$\sum_i \left( \begin{bmatrix} x_1^T \\ \vdots \\ x_N^T \end{bmatrix} w - y_i \right)^2$$

$$\begin{bmatrix} x_1^T \\ \vdots \\ x_N^T \end{bmatrix} w - \begin{bmatrix} y_1 \\ \vdots \\ y_N \end{bmatrix} = \begin{bmatrix} x_1 w - y_1 \\ \vdots \\ x_N w - y_N \end{bmatrix}$$

$$\begin{bmatrix} x_1 w - y_1 \\ \vdots \\ x_N w - y_N \end{bmatrix}^T \begin{bmatrix} x_1 w - y_1 \\ \vdots \\ x_N w - y_N \end{bmatrix} = \sum_i (x_i w - y_i)^2$$

18

$$\mathcal{L}(w) = \frac{1}{2N} (Xw - y)^T (Xw - y)$$

$$= \frac{1}{2N} (w^T X^T - y^T) (Xw - y)$$

$$= \frac{1}{2N} (w^T X^T X w - w^T X^T y - y^T X w + y^T y)$$

$$\mathcal{L}(w) = \frac{1}{2N} (w^T X^T X w - 2 y^T X w + y^T y)$$

$$\nabla_w \mathcal{L} = \frac{1}{N} (X^T X w - X^T y) = 0$$

$$X^T X w = X^T y$$

$$w = (X^T X)^{-1} X^T y$$

Predictions on  $x$ :  $\hat{y} = X w = X (X^T X)^{-1} X^T y$   
hat matrix

19

### Deriving the Least Squares Estimate: Summary

$$\mathcal{L}(w) = \frac{1}{2N} \sum_{i=1}^N (x_i^T w - y_i)^2 = \frac{1}{2N} (w^T (X^T X) w - 2 \cdot y^T X w + y^T y)$$

We compute the gradient  $\nabla_w \mathcal{L} = 0$  using the matrix differentiation rules,

$$\nabla_w \mathcal{L} = \frac{1}{N} ((X^T X) w - X^T y)$$

By setting  $\nabla_w \mathcal{L} = 0$  and solving we get,

$$(X^T X) w = X^T y$$

$$w = (X^T X)^{-1} X^T y \quad (\text{Assuming inverse exists})$$

The predictions made by the model on the data  $X$  are given by

$$\hat{y} = X w = X (X^T X)^{-1} X^T y$$

$X (X^T X)^{-1} X^T$  is called the "hat" matrix

20

### Complexity of Parameter Estimation

$$w = \underbrace{\begin{pmatrix} X^T & X \\ (D+1) \times N & N \times (D+1) \\ (D+1) \times (D+1) \end{pmatrix}^{-1}}_{(D+1) \times (D+1)} \underbrace{\begin{pmatrix} X^T & y \\ (D+1) \times N & N \times 1 \end{pmatrix}}_{(D+1) \times 1}$$

•  $Z = X^T X$  in  $O(D^2 N)$

• If  $D = O(N)$ , then the best known method (Le Gall) needs  $O(N^{2.37})$

•  $Z^{-1}$  in  $O(D^3)$

•  $A = X^T y$  in  $O(DN)$

•  $w = Z^{-1} A$  in  $O(D^2)$

Overall complexity for computing  $w$ :  $O(D^2 N + D^3)$

21

### Complexity of Parameter Estimation

What if  $X$  is defined by a join of several relations?

- The number of rows  $N$  may be exponential in the number of relations:

$$N = O(M^{\text{number relations}})$$

- $X$  is **sparse**, it can be represented in  $O(M)$  space losslessly for acyclic joins  
Acyclic joins are common in practice
- $w$  can be computed in  $O(D^2 M + D^3)$
- Find out more: <https://fdbresearch.github.io/>

22

### When Do We Expect $X^T X$ to be Invertible?

Matrix  $(X^T X) \in \mathbb{R}^{(D+1) \times (D+1)}$

•  $\text{rank}(X^T X) = \text{rank}(X) \leq \min\{D+1, N\}$

• It is invertible if  $\text{rank}(X) = D+1$

What if we use one-hot encoding for a feature like **day**?

•  $x_{\text{mon}}, \dots, x_{\text{sun}}$  stand for 0-1 valued variables in the one-hot encoding

• We always have  $x_{\text{mon}} + \dots + x_{\text{sun}} = 1$


• This introduces a linear dependence in the columns of  $X$  reducing the rank

• In this case, we can drop some features to adjust rank

We'll see alternative approaches later in the course


23

Least Squares Estimate in the Presence of Outliers




24

Least Squares Estimate in the Presence of Outliers




24

Least Squares Estimate in the Presence of Outliers



24

Least Squares Estimate in the Presence of Outliers



24