

Introduction to Spectral Methods

Numerical Solution of Poisson's Equation

Pablo Brubeck

Department of Physics
Tecnológico de Monterrey

October 7, 2016



Outline

Poisson's Equation

One dimension

Two Dimensions

Outline

Poisson's Equation

One dimension

Two Dimensions

Poisson's Equation

We are interested to find the function $u(x, y, z)$ that satisfies the boundary value problem

$$\nabla^2 u = f$$

Subject to boundary conditions $u(C) = g$.

Poisson's Equation

We are interested to find the function $u(x, y, z)$ that satisfies the boundary value problem

$$\nabla^2 u = f$$

Subject to boundary conditions $u(C) = g$.

Where the laplacian operator is $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$.

Poisson's Equation

We are interested to find the function $u(x, y, z)$ that satisfies the boundary value problem

$$\nabla^2 u = f$$

Subject to boundary conditions $u(C) = g$.

Where the laplacian operator is $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$.

When $f = 0$, we have Laplace's equation as a special case.

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$
- ▶ Newtonian gravity $\nabla^2 \phi = 4\pi G\rho$

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$
- ▶ Newtonian gravity $\nabla^2 \phi = 4\pi G\rho$
- ▶ Thermal equilibrium $\nabla^2 T = 0$

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$
- ▶ Newtonian gravity $\nabla^2 \phi = 4\pi G\rho$
- ▶ Thermal equilibrium $\nabla^2 T = 0$
- ▶ Incompressible fluid flow $\nabla^2 \nu = 0, \nabla^2 p = f(\nu, V)$

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$
- ▶ Newtonian gravity $\nabla^2 \phi = 4\pi G\rho$
- ▶ Thermal equilibrium $\nabla^2 T = 0$
- ▶ Incompressible fluid flow $\nabla^2 \nu = 0, \nabla^2 p = f(\nu, V)$
- ▶ Mechanical Engineering

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$
- ▶ Newtonian gravity $\nabla^2 \phi = 4\pi G\rho$
- ▶ Thermal equilibrium $\nabla^2 T = 0$
- ▶ Incompressible fluid flow $\nabla^2 \nu = 0, \nabla^2 p = f(\nu, V)$
- ▶ Mechanical Engineering
- ▶ Image processing

Poisson's Equation has many important applications.

- ▶ Electrostatics $\nabla^2 V = -\rho/\epsilon_0$
- ▶ Newtonian gravity $\nabla^2 \phi = 4\pi G\rho$
- ▶ Thermal equilibrium $\nabla^2 T = 0$
- ▶ Incompressible fluid flow $\nabla^2 \nu = 0, \nabla^2 p = f(\nu, V)$
- ▶ Mechanical Engineering
- ▶ Image processing
- ▶ And much more ...

Outline

Poisson's Equation

One dimension

Two Dimensions

Poisson's Equation in one dimension

$$\frac{d^2 u}{dx^2} = f(x)$$

Subject to $u(a) = g_1$ and $u(b) = g_2$.

It is not always possible to integrate twice to find the solution.

We must translate this continuous problem into a discrete one.

For the space coordinate $x \in [a, b]$, we choose n nodes $\{x_i\}_{i=1}^n$ on which we would like to know the value of $u(x_i) = u_i$.

$$\text{Input } u_1, u_n, \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}, \text{ and } \vec{f} = \begin{bmatrix} f(x_1) \\ f(x_2) \\ \vdots \\ f(x_n) \end{bmatrix}; \text{ output } \vec{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}.$$

The discrete solution must solve the continuous problem.

Is there a way to express $u(x)$ in terms of u_i ?

$$u(x) = \sum_j u_j \delta_j(x)$$

The discrete solution must solve the continuous problem.

Is there a way to express $u(x)$ in terms of u_i ?

$$u(x) = \sum_j u_j \delta_j(x)$$

We introduce the cardinal functions $\delta_j(x)$.

$$u(x_i) = \sum_j u_j \delta_j(x_i) = u_i$$

The discrete solution must solve the continuous problem.

Is there a way to express $u(x)$ in terms of u_i ?


$$u(x) = \sum_j u_j \delta_j(x)$$

We introduce the cardinal functions $\delta_j(x)$.

$$u(x_i) = \sum_j u_j \delta_j(x_i) = u_i$$

$$\delta_j(x_i) = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases}$$

This is how the cardinal functions look like.



Now differentiate the expansion in cardinal functions.

$$u'(x_i) = \sum_j u_j \delta'_j(x_i)$$

Now differentiate the expansion in cardinal functions.

$$u'(x_i) = \sum_j u_j \delta'_j(x_i)$$

We have found a matrix $D_{ij} = \delta'_j(x_i)$ that operates as a derivative.

$$\frac{d}{dx} \vec{u} = D \vec{u}$$

Now differentiate the expansion in cardinal functions.

$$u'(x_i) = \sum_j u_j \delta'_j(x_i)$$

We have found a matrix $D_{ij} = \delta'_j(x_i)$ that operates as a derivative.

$$\frac{d}{dx} \vec{u} = D \vec{u}$$

$$\frac{d^2}{dx^2} \vec{u} = D^2 \vec{u} = \vec{f}$$

Solve the system of linear equations.

$$D^2 \vec{u} = \vec{f}$$

Solve the system of linear equations.

$$D^2 \vec{u} = \vec{f}$$

Recall that u_1 and u_n were given as boundary conditions.

Solve the system of linear equations.

$$D^2 \vec{u} = \vec{f}$$

Recall that u_1 and u_n were given as boundary conditions.

$$\begin{bmatrix} D_{11}^2 & D_{1j}^2 & D_{1n}^2 \\ D_{i1}^2 & D_{ij}^2 & D_{in}^2 \\ D_{n1}^2 & D_{nj}^2 & D_{nn}^2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_j \\ u_n \end{bmatrix} = \begin{bmatrix} D_{11}^2 \\ D_{i1}^2 \\ D_{n1}^2 \end{bmatrix} u_1 + \begin{bmatrix} D_{1j}^2 \\ D_{ij}^2 \\ D_{nj}^2 \end{bmatrix} [u_j] + \begin{bmatrix} D_{1n}^2 \\ D_{in}^2 \\ D_{nn}^2 \end{bmatrix} u_n = \begin{bmatrix} f_1 \\ f_i \\ f_n \end{bmatrix}$$

With i and j ranging from 2 to $n - 1$.

We have $n - 2$ unknowns, but still n equations.

Discard the first and last rows.

$\nabla^2 u = f$ will not hold true at the boundary, but we will ensure the boundary conditions.

$$\begin{aligned} [D_{ij}^2] [u_j] &= [f_i] - u_1 [D_{i1}^2] - u_n [D_{in}^2] \\ [u_j] &= [D_{ij}^2]^{-1} [f_i - u_1 D_{i1}^2 - u_n D_{in}^2] \end{aligned}$$

With i and j ranging from 2 to $n - 1$.

Cardinal functions are obtained from a mother function.

Usually we chose an orthogonal polynomial. It all depends on the domain of your solution.

Chebysheb	$\Psi_n(x) = T_n(x)$	$x \in [-1, 1]$
Legendre	$\Psi_n(x) = P_n(x)$	$x \in [-1, 1]$
Hermite	$\Psi_n(x) = H_n(x)e^{-x^2/2}$	$x \in (-\infty, \infty)$
Laguerre	$\Psi_n(x) = xL_{n-1}^{(1)}(x)e^{-x/2}$	$x \in [0, \infty)$

Cardinal functions are obtained
from a mother function.

Let $\Psi_n(x)$ be a function with n roots on $\{x_i\}_{i=1}^n$, i. e. $\Psi_n(x_i) = 0$.
By expanding $\Psi_n(x)$ around its roots we can find a expression for $\delta_j(x)$.

Cardinal functions are obtained
from a mother function.

Let $\Psi_n(x)$ be a function with n roots on $\{x_i\}_{i=1}^n$, i. e. $\Psi_n(x_i) = 0$.
By expanding $\Psi_n(x)$ around its roots we can find a expression for $\delta_j(x)$.

$$\Psi_n(x) = 0 + \Psi'_n(x_j)(x - x_j) + \frac{\Psi''_n(x_j)}{2}(x - x_j)^2 + O((x - x_j)^3)$$

Cardinal functions are obtained from a mother function.

Let $\Psi_n(x)$ be a function with n roots on $\{x_i\}_{i=1}^n$, i. e. $\Psi_n(x_i) = 0$. By expanding $\Psi_n(x)$ around its roots we can find an expression for $\delta_j(x)$.

$$\Psi_n(x) = 0 + \Psi'_n(x_j)(x - x_j) + \frac{\Psi''_n(x_j)}{2}(x - x_j)^2 + O((x - x_j)^3)$$

$$\delta_j(x) = \frac{\Psi_n(x)}{(x - x_j)\Psi'_n(x_j)} = 1 + \frac{\Psi''_n(x_j)}{2\Psi'_n(x_j)}(x - x_j) + O((x - x_j)^2)$$

Differentiate to obtain the elements of D .

$$D_{ij} = \delta'_j(x_i) = \frac{1}{x_i - x_j} \frac{\Psi'_n(x_i)}{\Psi'_n(x_j)}, \quad i \neq j$$

The diagonal is a bit tricky.

$$D_{jj} = \delta'_j(x_j) = \frac{\Psi''_n(x_j)}{2\Psi'_n(x_j)}$$

MATLAB code

Solve Poisson's equation

$$\frac{d^2 u}{dx^2} = 1000 \cos(5\pi x) e^{-x^2}$$

with boundary conditions $u(-1) = 2$, $u(1) = -1$.

MATLAB code


Solve Poisson's equation

$$\frac{d^2 u}{dx^2} = 1000 \cos(5\pi x) e^{-x^2}$$

with boundary conditions $u(-1) = 2$, $u(1) = -1$.

```
1 [D,x]=chebD(n); D2=D*D; % Differentiation matrix and nodes
2 f=1000*cos(5*pi*x).*exp(-x.^2); % Source term
3 u=zeros(n,1); u([1,n])=[-1,2]; % Impose boundary conditions
4 % Solve and plot
5 u(2:n-1)=D2(2:n-1,2:n-1)\(f(2:n-1)-D2(2:n-1,[1,n])*u([1,n]));
6 plot(x,u);
```

Here is our solution.



Outline

Poisson's Equation

One dimension

Two Dimensions

Poisson's Equation in two dimensions

Our domain of solution will be the rectangle $[a, b] \times [c, d]$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f(x, y)$$

Subject to boundary conditions

$$u(x, y) = \begin{cases} g_1(y), & x = a \\ g_2(y), & x = b \\ h_1(x), & y = c \\ h_2(x), & y = d \end{cases}$$

In 2D we use matrices for u and f .

$$u(x_i, y_j) = u_{ij}$$

$$f(x_i, y_j) = f_{ij}$$

$$u(x, y) = \sum_{k,l} u_{kl} \delta_k(x) \delta_l(y)$$

Multiplying times D does the job of partial derivatives.

$$\frac{\partial u}{\partial x} = DU = [Du_{i1} \quad Du_{ij} \quad Du_{in}]$$

Multiplying times D does the job of partial derivatives.

$$\frac{\partial u}{\partial x} = DU = [Du_{i1} \quad Du_{ij} \quad Du_{in}]$$

$$\frac{\partial u}{\partial y} = UD^T = (DU^T)^T = \begin{bmatrix} (Du_{1i})^T \\ (Du_{ji})^T \\ (Du_{ni})^T \end{bmatrix}$$

Multiplying times D does the job of partial derivatives.

$$\frac{\partial u}{\partial x} = DU = [Du_{i1} \quad Du_{ij} \quad Du_{in}]$$

$$\frac{\partial u}{\partial y} = UD^T = (DU^T)^T = \begin{bmatrix} (Du_{1i})^T \\ (Du_{ji})^T \\ (Du_{ni})^T \end{bmatrix}$$

$$\nabla^2 u = D^2 U + U(D^2)^T$$

Now solve the system of linear equations.

$$D^2 U + U(D^2)^T = F$$

Boundary conditions are the tricky part.

$$U = \tilde{U} + U_B = \begin{bmatrix} 0 & 0 & 0 \\ 0 & u_{ij} & 0 \\ 0 & 0 & 0 \end{bmatrix} + \begin{bmatrix} u_{11} & u_{1j} & u_{1n} \\ u_{i1} & 0 & u_{in} \\ u_{n1} & u_{nj} & u_{nn} \end{bmatrix}$$

$$D^2 \tilde{U} + \tilde{U}(D^2)^T = F - D^2 U_B - U_B(D^2)^T = \tilde{F}$$

Discard first and last rows and columns.

$\nabla^2 u = f$ will not hold true at the boundary, but we will ensure the boundary conditions.

$$(D_{ij}^2)\tilde{U}_{ij} + \tilde{U}_{ij}(D_{ij}^2)^T = \tilde{F}_{ij}$$

With i and j ranging from 2 to $n - 1$.

Now we just have to solve $(n - 2)^2 \times (n - 2)^2$ system of linear equations.

Matrix inversion does not help.

A matrix equation of the form $AX + XB = C$ is known as a Sylvester equation.

There's a built-in MATLAB function that solves them.

```
X=sylvester(A,B,C)
```

This algorithm uses the Schur decomposition to solve a block-triangular matrix very efficiently.

Two dimensional example

Solve Laplace's equation on $(x, y) \in [-1, 1] \times [-1, 1]$

$$\nabla^2 u = 0$$


with boundary conditions

$$u(x, y) = \begin{cases} \sin^4(\pi x), & y = 1 \text{ and } -1 < x < 0, \\ \frac{1}{5} \sin(3\pi y), & x = 1, \\ 0, & \text{otherwise.} \end{cases}$$

MATLAB Code

```
1  [D,x]=chebD(n); D2=D*D; y=x';
2  [xx, yy]=ndgrid(x);
3
4  % Boundary conditions
5  g=[0.2*sin(3*pi*y); 0*y];
6  h=[(x<0).*sin(pi*x).^4, 0*x];
7  uu=zeros(n);
8  uu([1 n],:)=g;
9  uu(:,[1 n])=h;
10
11 % Solve Laplace's equation
12 F=zeros(n);
13 RHS=F-D2(:,[1 n])*g-h*D2(:,[1 n])';
14 uu(2:n-1, 2:n-1)=sylvestor(D2(2:n-1, 2:n-1), ...
15 D2(2:n-1, 2:n-1)', RHS(2:n-1, 2:n-1));
16
17 surf1(xx,yy,uu,'light'); colormap(jet(256));
18 shading interp; axis square;
```

Here is our solution.



References



W. H. Press, *Numerical Recipes*.

Cambridge University Press, 2007.



L. N. Trefethen, *Spectral Methods in MATLAB (Software, Environments, Tools)*.

SIAM: Society for Industrial and Applied Mathematics, 2001.