

Math 5050 – Special Topics: Manifolds– Fall 2025

w/Professor Berchenko-Kogan

Paul Carmody

Section 7: Quotients – May 17, 2025

Pg. 77: Exercise 7.11 (Real projective space as a quotient of a sphere).* For $x = (x^1, \dots, x^n) \in \mathbb{R}^n$, let $\|x\| = \sqrt{\sum_i (x^i)^2}$ be the modulus of x . Prove that the map $f : \mathbb{R}^{n+1} - \{0\} \rightarrow S^n$ given by

$$f(x) = \frac{x}{\|x\|}$$

induces a homeomorphism $\bar{f} : \mathbb{R}P^n \rightarrow S^n / \sim$. Where

$$x \sim y \iff x = \pm y, x, y \in S^n$$

(*Hint:* Find an inverse map

$$\bar{g} : S^n / \sim \rightarrow \mathbb{R}P^n$$

and show that both \bar{f} and \bar{g} are continuous.)

Given the relation above $x \sim y \iff x = \pm y, x, y \in S^n$. Define

$$\bar{g}([x]) = [x]$$

Note that on the left $[x] \in S^n / \sim$ and $[x] \in \mathbb{R}P^n$. For clarity,¹

$$[a] \in S^n / \sim \implies [a] = \{a, -a\} \text{ where } a \in S^n$$

$$[b] \in \mathbb{R}P^n \implies [b] = \{x \in \mathbb{R}^{n+1} \mid x = \alpha b, \forall \alpha \in \mathbb{R}\} \text{ where } b \in \mathbb{R}^{n+1}$$

Notice that

$$\begin{aligned} \bar{f}([b]) &= [f(b)] = \left[\frac{b}{\|b\|} \right] \in S^n / \sim \\ \bar{g} \left(\left[\frac{b}{\|b\|} \right] \right) &= [b] \\ \therefore \bar{g} \circ \bar{f} &= \mathbb{I} \\ \text{and } \bar{g}([a]) &= [a] \in \mathbb{R}P^n \\ \bar{f}(\bar{g}([a])) &= \left[\frac{[a]}{\|[a]\|} \right] = [f(a)] = [a] \in S^n / \sim \\ \therefore \bar{f} \circ \bar{g} &= \mathbb{I} \end{aligned}$$

\bar{f} is continuous because f is continuous. \bar{g} is continuous because it is a mapping of one identity to another. Therefore, \bar{f} is a homeomorphism.

$$\begin{array}{ccc} S & \xrightarrow{f} & Y \\ \pi \downarrow & \searrow \bar{f} & \\ S/\sim & & \end{array}$$

Problems

7.1. Image of the inverse image of a map

Let $f : X \rightarrow Y$ be a map of sets, and let $B \subset Y$. Prove that $f(f^{-1}(B)) = B \cap f(X)$. Therefore, if f is surjective, then $f(f^{-1}(B)) = B$.

\subseteq : Let $b \in B$ and $a \in X$ such that $f(a) = b$. Then, $a \in f^{-1}(b)$, thus a is an arbitrary point in $f^{-1}(B)$. We know that $f(a) \in f(X)$ and $f(a) \in B$, therefore $f(a) \in B \cap f(X)$ and $f(f^{-1}(B)) \subseteq B \cap f(X)$.

\supseteq : Let $b \in B \cap f(X)$. Since $b \in f(X)$, there exists $a \in X$ such that $f(a) = b$ and since $b \in B$ then $b = f(a) \in f(f^{-1}(b)) \subseteq f^{-1}(B)$. Therefore $b \in f(f^{-1}(B))$

¹ $\mathbb{R}P^n \equiv \mathbb{R}^{n+1} / r(x, y)$ where $r(x, y)$ is the relation that is true when x, y, p are colinear.

7.2. Real projective plane

Let H^2 be the closed upper hemisphere in the unit sphere S^2 , and let $i : H^2 \rightarrow S^2$ be the inclusion map. In the notation of example 7.13, prove that the induced map $f : H^2 / \sim \rightarrow S^2 / \sim$ is a homeomorphism. (*Hint*: Imitate Proposition 7.3.) Let H^2 be the upper hemisphere and S^2 be the unit sphere

$$H^2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1, z \geq 0\}$$

$$S^2 = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$$

These two are homeomorphic to each via

$$\varphi : S^2 \mapsto H^2$$

$$\varphi(x, y, z) = (x, y, |z|)$$

and its inverse

$$\psi : H^2 \mapsto S^2$$

$$\psi(x, y, z) = (x, y, z)$$

Define the relations

$$(x, y, z) \sim (-x, -y, -z) \rightarrow x^2 + y^2 = z^2, \forall (x, y, z) \in S^2$$

$$(x, y, z) \sim (x, y, z) \rightarrow \sqrt{x^2 + y^2} = z, \forall (x, y, z) \in H^2$$

Then we induce $f : H^2 / \sim \rightarrow S^2 / \sim$ as

$$f([(x, y, z)]) = [(x, y, z)]$$

7.3. Closedness of the diagonal of a Hausdorff Space

Deduce Theorem 7.7 from Corollary 7.8 (*Hint*: To prove that if S / \sim is Hausdorff, then the graph R of \sim is closed in $S \times S$, use the continuity of the projection map $\pi : S \rightarrow S / \sim$. To prove the reverse implication, use the openness of π .)

7.4. Quotient of a sphere with antipodal points identified

Let S^n be the unit sphere centered at the origin in \mathbb{R}^{n+1} . Define an equivalence relation \sim on S^n by indentifying antipodal points:

$$x \sim y \iff x = \pm y, x, y \in S^n$$

- Show that \sim is an open equivalence relation.
- Apply Theorem 7.7 and Corollary 7.8 to prove that the quotient space S^n / \sim is Hausdorff, without making use of the homeomorphism $\mathbb{R}P^n \cong S^n / \sim$.

7.5. Orbit space of a continuous group action

Suppose a right action of a topological group G on a topological space S is continuous, this simply means that the map $S \times G \rightarrow S$ describing the action is continuous. Define two points x, y of S to be equivalent if they are in the same orbit; i.e., there is an element $g \in G$ such that $y = xg$. Let S/G be the quotient space; it is called the *orbit space* of the action. Prove that the projection map $\pi : S \rightarrow S/G$ is an open map. (This problem generalizes Proposition 7.14, in which $G = R^\times = \mathbb{R} - \{0\}$ and $S = \mathbb{R}^{n+1} - \{0\}$. Because \mathbb{R}^\times is commutative, a left \mathbb{R}^\times -action becomes a right \mathbb{R}^\times -action if scalar multiplication is written on the right.)

7.6. Quotient of \mathbb{R} by $2\pi\mathbb{Z}$.

Let the additive group $2\pi\mathbb{Z}$ act on \mathbb{R} on the right by $x \cdot 2\pi n = x + 2\pi n$, where n is an integer. Show that the orbit space $\mathbb{R}/2\pi\mathbb{Z}$ is a smooth manifold.

7.7. The circle as a quotient space

- Let $\{(U_\alpha, \phi_\alpha)\}_{\alpha=1}^2$ be the atlas of circles S^1 in Example 5.7, and let $\bar{\phi}_\alpha$ be the map ϕ_α followed by the projection $\mathbb{R} \rightarrow \mathbb{R}/2\pi\mathbb{Z}$. On $U_1 \cap U_2 = A \sqcup B$, since ϕ_1 and ϕ_2 differ by an integer multiple of 2π , $\bar{\phi}_1 = \bar{\phi}_2$. Therefore, $\bar{\phi}_1$ and $\bar{\phi}_2$ piece together to give a well-defined map $\bar{\phi} : S^1 \rightarrow \mathbb{R}/2\pi\mathbb{Z}$. Prove that $\bar{\phi}$ is C^∞ .
- The complex exponential $\mathbb{R} \rightarrow S^1, t \mapsto e^{it}$, is constant on each orbit of the action of $2\pi\mathbb{Z}$ on \mathbb{R} . Therefore, there is an induced map $F : \mathbb{R}/2\pi\mathbb{Z} \rightarrow S^1, F([t]) = e^{it}$. Prove that F is C^∞ .
- Prove that $F : \mathbb{R}/2\pi\mathbb{Z} \rightarrow S^1$ is a diffeomorphism.

7.8. The Grassmannian $G(k, n)$

7.9. Compactness of real projective space

Show that the real projective space $\mathbb{R}P^n$ is compact. (*Hint*: Use Exercise 7.11.)