RSI Checklist

T-4

Check self, positions	Check
Consider SMR	Check
Supply oxygen (consider all: NP, HFM, BVM+PEEP)	Check
Activate suction, position	Check
Patient position	Check
IV Access points- identify, start fluid bolus	Check
Attach monitors, cycle BP, **Optimize CV**	Check

T-2

Geardump:	
ETT+10cc syringe, smaller ETT	Check
Laryngoscope, check light	Check
Bougie on chest	Check
CO2 device	Check
Tie device	Check
EGD on standby	Check
Surgical kit on standby, handshake larynx	Check

T-1

Brief drugs	Check
Brief plan A and Plan B	Check
Cycle BP	Check

T

Give drugs	Check
Verbalize 1st look	*
Pass tube	*
Confirm placement, **CO2+**	Check

T+1

Cycle BP	Check
Place OG	Check
Secure and package	Check

Tracheal intubation of critically ill adults

Pre-oxygenate and checklist

Position: head up if possible
Assess airway and identify cricothyroid membrane
Waveform capnograph
Pre-oxygenate: facemask / CPAP / NIV / nasal O_2
Optimise cardiovascular system
Share plan for failure

Note the time

Plan A: Tracheal intubation

Laryngoscopy
Maximum 3 attempts

Maintain oxygenation
- **Continuous nasal oxygenation**
- **Facemask ventilation between attempts**

Neuromuscular block
Video or direct laryngoscopy +/- bougie or stylet
External laryngeal manipulation
Remove cricoid

Succeed → Confirm with capnography

First failure → **Call HELP**
- Video laryngoscopy
- Get front of neck airway(FONA)set

Fail → **Declare "failed intubation"**

Plan B/C: Rescue oxygenation

2nd generation supraglottic airway ⇄ **Facemask**
- **2 person**
- **Adjuncts**

Maximum 3 attempts each
Change device / size / operator
Open front of neck airway set

Succeed → **Stop, think, communicate**

Options
- Wake patient if planned
- Wait for expert
- Intubate via supraglottic airway x1
- Front of neck airway

Fail → **Declare "can't intubate, can't oxygenate"**

Plan D: Front of neck airway: FONA

Use FONA set
Scalpel cricothyroidotomy

Extend neck
Neuromuscular blockade
Continue rescue oxygenation

Trained expert only

Other FONA techniques

Non-scalpel cricothyroidotomy
Percutaneous tracheostomy
Surgical tracheostomy

EXPERT: one extra attempt if appropriate
Video / direct laryngoscopy
Facemask or supraglottic airway
Front of neck airway

This flowchart forms part of the DAS, ICS, FICM, RCoA guideline for tracheal intubation in critically ill adults and should be used in conjunction with the text.