# Models for quantum memory enhanced protocols

- B&K
- Purification
- "Kwiat" scheme

## Models for quantum memory enhanced protocols

- B&K
- Purification
- "Kwiat" scheme

#### Key parameters

- **d** node sep.
- n no. of memories (including the electron spin as a memory)
- **0.3** freq. conversion efficiency
- **0.3** outcoupling efficiency (inc ZPL contrib.)
- **2 μs** entanglement generation clock rate
- 20 μs memory gate time
- 1/400 memory 1/e lifetime (in no. attempts)
- 10 μs purification gate time
- **0.1** SPDC source pair rate


### Models for quantum memory enhanced protocols


- B&K
- Purification
- "Kwiat" scheme


#### Key parameters


- d node sep.
- n no. of memories (including the electron spin as a memory)
- **0.3** freq. conversion efficiency
- **0.3** outcoupling efficiency (inc ZPL contrib.)
- **2 μs** entanglement generation clock rate
- 20 μs memory gate time
- 1/400 memory 1/e lifetime (in no. attempts)
- 10 μs purification gate time
- **0.1** SPDC source pair rate

Models shown assume that we are operating in the limit where p(click per round) << 1)











# Fix the distance, change the number of memories



# Fix the distance, change the number of memories



# Fix the distance, change the number of memories

